

Original Research Article

Effect of Supplementation of Ashwagandha (*Withania somnifera*) Root Powder on Body Weight and Egg Quality in Layers

Commented [RA1]: Most appropriate: Growth performance (as BW and BWG are measured)

ABSTRACT

Background: As the demand for natural and sustainable solutions in poultry production continues to rise, ashwagandha root powder presents itself as a promising avenue for improving egg quality while aligning with consumer preferences for wholesome and nutritious products. Therefore, research trial was conducted to study the effect of supplementation of ashwagandha root powder on body weight and egg quality in layers during a period of 10 weeks.

Methodology: One hundred and twenty white leghorn layers were randomly divided into four treatments and reared on standard managerial conditions. The Ashwagandha root powder was supplemented with the basal diet at 0.5% level (T₁), 1% (T₂), 1.5 % (T₃) while T₀ was kept as control i.e. basal diet without ashwagandha root powder.

Commented [RA2]: White Leghorn

Design used: Completely Randomized Design

Commented [RA3]: Explain in 1 or 2 sentences

Results: The result noted that the body weight change was not affected by ashwagandha root powder supplementation. Egg quality parameters viz., egg shape index, shell weight, yolk weight, yolk height, albumen weight and albumen index were not influenced by addition of ashwagandha root powder. However, significant ($p < 0.05$) effect was observed on yolk index. No significant difference ($p > 0.05$) was seen on sensory evaluation of egg.

Commented [RA4]: P capital used in the description). Follow one form..

KEYWORDS: layers, egg quality, yolk index, *Withania somnifera*, sensory evaluation

1. INTRODUCTION

In recent years, there has been a growing interest in utilizing herbal feed supplements to enhance the health and productivity of poultry. Among these supplements, ashwagandha (*Withania somnifera*) has gained attention for its potential benefits in improving egg quality in layer hens. Ashwagandha, also known as Indian ginseng or winter cherry, is an adaptogenic herb renowned in traditional Ayurvedic medicine for its various health-promoting properties. Ashwagandha contains bioactive compounds

such as withanolides (witanopherin A, withanolides A-Y, witanone), alkaloids (sominin, witanin, anaferin and somniferinin) and flavonoids (3-O-rutinoside, quercetin) and its glycosidic derivatives [13] which possess antioxidant, anti-bacterial, anti-inflammatory and immunomodulatory properties [6]. Ashwagandha is loaded with minerals such as Ca, Cd, Al, K, Mn, Fe, Ni, Cu and Zn. Moreover, it also contains crude fibre (21-25 %), starch (6.09-9.46 mg/g), tannins (0.39-0.82), reducing and non-reducing sugars [10]. These compounds are believed to contribute to ashwagandha's potential to positively impact health and egg quality in layers. Understanding the effects of ashwagandha on layer egg quality is not only pertinent for the poultry industry's quest for sustainable and natural solutions but also holds significance for consumers who prioritize the nutritional quality of eggs. Therefore, we delve the effect of supplementation of Ashwagandha (*Withania somnifera*) root powder on body weight and egg quality in layers.

2. MATERIAL AND METHODS

The study was conducted at Poultry unit, Mahatma Phule Krishi Vidyapeeth, Rahuri, Ahmednagar (Maharashtra). The research trial was carried out on 120 white leghorn layers of 40 weeks of age for period of 10 weeks (40-50 weeks) and randomly assigned into four treatment groups. Four treatment includes T₀ (control): fed basal diet; T₁: fed basal diet incorporated with 0.5 percent ashwagandha root powder; T₂: fed basal diet incorporated with 1 percent ashwagandha root powder; T₃: fed basal diet incorporated with 1.5 percent ashwagandha root powder. The basal diet of laying hens was formulated as per BIS (2007). To calculate body weight gain of birds initial and final body weights of the laying hens were recorded. The maximum length and maximum width or breadth of egg was measured by vernier calliper and shape index was express as ratio of maximum width and maximum length of eggs in percentage.

$$\text{Shape index} = \frac{\text{Width of egg in mm}}{\text{Length of egg in mm}} \times 100$$

The weight of shell along with shell membrane was measured in grams on an electronic balance after proper drying of egg shell. Specific gravity was tested using the floatation method [11]. Yolk and albumen were separated with a separating spoon and weighed with an electronic balance in grams. The length of albumen and yolk diameter was measured with the help of Vernier Calliper in

Commented [RA5]: Should include some other references..

Commented [RA6]: I suggest: « contains»

Commented [RA7]: What about other zootechnical performances? Need to mention some..

Commented [RA8]: What about the housing type. Missing information on «standard managemental conditions” including rearing condition, lighting schedule etc etc...

Commented [RA9]: Include 1 or 2 line description on ethical treatment

Commented [RA10]: Overall experimental design is not clear.

millimetres. The height of yolk and albumen were recorded with spherometer. Albumen index was measured using formula,

$$\text{Albumin index} = \frac{\text{Average height of albumin in mm}}{\text{Albumin width of albumin in mm}} \times 100$$

Yolk index was calculated as per the formula given by ~~Card and Nesheim (1972)~~ [7]. The data obtained on various egg quality parameters during experiment were statistically analyzed using Completely Randomized Design (CRD) as per procedure described by Snedecor and Cochran (1994).

Formatted: Strikethrough

Commented [RA11]: Need to include some information on «Statistical analysis»

Commented [RA12]: Missing info: sensory evaluation procedure...

3. RESULTS AND DISCUSSION

3.1 Effect on body weight gain

The average body weight gain in T₀, T₁, T₂ and T₃ treatment groups were presented in Table 1. The average values of body weight changes of laying hens were 52.06, 51.51, 51.14 and 52.20 gm in T₀, T₁, T₂ and T₃ treatments, respectively. There was no significant difference was noted in the gain in body weight of white leghorn layers in different treatment groups. The results of body weight gain are similar with the findings [1,2] noted that no significant effect of garlic feeding on gain in body weight in laying hens.

Commented [RA13]: How it was defined? Clarify the method for determining significant differences (eg. post-hoc tests).

Commented [RA14]: Why garlic? There are lots of study available on effects of WS in layer.. for example: [S-10-6-54-602.pdf \(thejournal.com\)](https://doi.org/10.654602/thejournal.com)

Table 1. Body weight changes of layers

Treatment	Initial Body weight	Final Body weight	Body Weight Gain
T ₀	1307.77	1359.83	52.06
T ₁	1307.06	1358.57	51.51
T ₂	1312.63	1363.77	51.14
T ₃	1312.03	1364.23	52.20

Commented [RA15]: Statistical analysis is not clear, what about the P value ???

3.2 Effect of feeding ashwagandha root powder on egg quality parameters of white leghorn layers

Commented [RA16]: White Leghorns

The mean values of egg shape index, egg shell weight and specific gravity were depicted in Table 2. Numerically higher average egg shape index was reported in group T₂ (1 % ARP) followed by T₃, T₁ and T₀ however, the statistical analysis revealed that no significant differences (p value?) in all

treatment groups over entire period of experiment. The results are in close agreement [15] examined egg shape index were not significantly differ between treatment groups.

Table 2. Effect of ashwagandha root powder on external egg qualities of white leghorn layers

Egg Shape index (%)	Weeks	Treatment				SE (±)	CD @ 5
		T ₀	T ₁	T ₂	T ₃		%
Egg Shape index (%)	40-42 Weeks	73.34	75.33	76.97	73.97	0.81	NS
	42-44 Weeks	73.84	74.94	75.94	74.73	0.71	NS
	44-46 Weeks	71.76	73.91	74.25	73.60	0.91	NS
	46-48 Weeks	72.62	75.75	77.37	75.82	1.08	NS
	48-50 Weeks	72.90	74.62	75.23	74.72	0.61	NS
Shell weight (g)	40-42 Weeks	6.15	6.28	6.34	6.36	0.10	NS
	42-44 Weeks	6.29	6.23	6.41	6.33	0.15	NS
	44-46 Weeks	6.25	6.31	6.36	6.28	0.17	NS
	46-48 Weeks	6.42	6.34	6.30	6.35	0.13	NS
	48-50 Weeks	6.33	6.37	6.45	6.35	0.10	NS
Specific gravity	40-42 Weeks	1.089	1.088	1.090	1.086	0.0011	NS
	42-44 Weeks	1.090	1.091	1.085	1.087	0.0018	NS
	44-46 Weeks	1.091	1.088	1.088	1.090	0.0012	NS
	46-48 Weeks	1.088	1.089	1.090	1.089	0.0011	NS
	48-50 Weeks	1.091	1.088	1.085	1.090	0.0020	NS

The mean values in same row with different superscripts differ significantly (P< 0.05)

Statistical analysis of data on shell weight was shown in Table (2) were noted no significant difference (p value) among the treatment throughout the trial period. -Adding of herbal extracts of garlic, thyme and caraway to drinking water of quails were no significant effect on shell weight [3]. The Uttara layers fed 1gm, 2gm and 4gm/kg arjun bark powder produced eggs with unchanged egg shape index and shell weight [16]. Regarding egg specific gravity values, no significant differences (p value) were found in all treatment groups throughout the experimental period. These results are parallel to

Commented [RA18]: Idem

Commented [RA17]: What is SE?? Write below the table as footnote

Commented [RA19]: Why to compare with other extracts? Why not with WS extracts in another study?

	40-42 Weeks	17.96	17.88	17.64	17.88	0.30	NS
	42-44 Weeks	18.19	17.94	18.51	18.15	0.25	NS
	44-46 Weeks	18.49	18.38	18.82	18.59	0.26	NS
	46-48 Weeks	17.85	18.39	18.47	17.92	0.28	NS
	48-50 Weeks	18.01 ^b	18.26 ^b	19.12 ^a	18.61 ^{ab}	0.19	0.60
Yolk diameter (mm)	40-42 Weeks	36.32	38.81	40.19	39.92	1.18	NS
	42-44 Weeks	39.64	38.38	40.34	40.14	0.89	NS
	44-46 Weeks	39.25	38.54	39.61	39.15	0.44	NS
	46-48 Weeks	37.74	38.65	38.67	37.88	0.69	NS
	48-50 Weeks	39.15	37.45	39.49	39.71	0.78	NS
Yolk index (%)	40-42 Weeks	46.50	45.84	44.92	46.52	0.77	NS
	42-44 Weeks	47.69	46.92	45.83	45.33	1.38	NS
	44-46 Weeks	46.98 ^b	47.58 ^a	48.03 ^a	47.78 ^a	0.18	0.60
	46-48 Weeks	48.14 ^b	48.63 ^{ab}	49.05 ^a	48.89 ^a	0.20	0.65
	48-50 Weeks	46.11 ^b	47.03 ^b	48.52 ^a	46.91 ^b	0.33	1.06

The mean values in same row with different superscripts differ significantly ($P < 0.05$)

Biweekly data pertaining to albumen weight, height and albumen index are set out in Table 4.

The average albumen height and albumen index did not differ significantly among treatment groups during progressive weeks of age of white leghorn layers as well as with respect to whole period. Results are in agreement [14] who discovered that albumen index had not been affected significantly due to dietary supplementation of ARP. Also, [12] examined *Withania somnifera* supplements had non-significant effect on albumen height.

Data related to albumen weight of eggs of white leghorn layers are represented in Table 4. It was notified that the difference in albumen weight was non-significant during different weeks of age during period of trial except at beginning of experiment. At first bi-weekly (40-42 weeks) of experiment, significant difference observed in albumen weight. The mean significant higher value observed in T₃ (35.22 g) treatment group followed by T₁ (34.39), T₂ (33.86) and T₀ (32.67), respectively. It can be concluded; dietary supplementation of ashwagandha root powder at different levels had no effect on the albumen weight. The findings of study were parallel with [4] demonstrated

Commented [RA21]: Write signification of superscript in footnote

albumen weight and albumen index did not significantly affected on addition of *Withania somnifera* to Japanese quail ration. Additionally, [21] determined that egg albumen had not influenced on the addition of garlic powder at 5 and 10 g/kg of feed garlic in laying hens.

Table 4. Effect of ashwagandha root powder on egg albumen height, weight and albumen index of white leghorn layers

Albumen height (mm)	Weeks	Treatment				SE (±)	CD @ 5%
		T ₀	T ₁	T ₂	T ₃		
Albumen height (mm)	40-42 Weeks	7.25	7.53	7.54	7.73	0.28	NS
	42-44 Weeks	7.60	7.96	7.75	7.80	0.24	NS
	44-46 Weeks	7.61	7.83	8.06	7.50	0.28	NS
	46-48 Weeks	7.65	7.76	7.80	7.87	0.14	NS
	48-50 Weeks	7.29	7.73	7.40	7.23	0.37	NS
Albumen weight (g)	40-42 Weeks	32.67 ^c	34.39 ^{ab}	33.86 ^{bc}	35.22 ^a	0.39	1.27
	42-44 Weeks	35.04	33.05	32.80	32.75	0.87	NS
	44-46 Weeks	34.42	32.85	33.58	33.26	0.78	NS
	46-48 Weeks	33.09	33.76	34.15	33.94	1.19	NS
	48-50 Weeks	33.95	34.08	33.38	34.30	0.84	NS
Albumen index (%)	40-42 Weeks	12.19	13.33	12.79	12.57	0.65	NS
	42-44 Weeks	12.16	12.67	12.07	11.58	0.47	NS
	44-46 Weeks	12.69	12.61	12.98	12.51	0.41	NS
	46-48 Weeks	13.12	13.60	12.93	13.07	0.70	NS
	48-50 Weeks	11.41	12.55	11.57	11.18	0.60	NS

The mean values in same row with different superscripts differ significantly (P < 0.05)

3.3 Sensory evaluation of eggs

Regarding sensory attributes of eggs sample viz., their appearance & colour, flavour, aroma and overall acceptability showed in Table 5. It was observed that there was no significant distinction in appearance & colour, flavour, aroma and overall acceptability of eggs among the treatment groups of

white leghorn layers fed diets supplemented with Ashwagandha root powder at different levels. Likely, [17] revealed that addition of Phyto herbal feed additives in laying hens didn't impact on egg aroma, flavour and all eggs were accepted by panelists. Inclusion of herbal feed additives (black cumin, garlic and turmeric) with feed in laying hens did not affect on organoleptic parameters of eggs [19].

Commented [RA22]: Contain?? Mention the % or inclusion ratio of the plants

Table. 5 Effect of dietary supplementation of ashwagandha root powder on sensory score of eggs

Treatments	Sensory evaluation of Eggs			
	Appearance & Colour	Flavour	Aroma	Overall acceptability
T ₀	7.89	7.89	8.07	7.95
T ₁	7.83	8.04	8.06	7.98
T ₂	7.84	7.99	7.98	7.93
T ₃	7.80	8.02	7.81	7.88
SE (±)	0.12	0.09	0.08	0.06
CD @ 5 %	NS	NS	NS	NS

Commented [RA23]: What is SE?? Include footnote

Commented [RA24]: Footnote..

CONCLUSION

The statistical analysis of data revealed that utilization of ashwagandha root powder at different levels with basal diet of layer had no significant effect on egg shape index, shell weight, egg specific gravity, albumen height, index and yolk height. However, yolk index showed significant effect on supplementation ashwagandha root powder.

Commented [RA25]: Missing info: potential practical applications, brief description of future research needs.

REFERENCES

- Asrat M., Zeryehun T., Amha N. and Urge M. (2018). Effects of supplementation of different levels of garlic (*Allium sativum*) on egg production, egg quality and hatchability of White leghorn chicken. tesfahiwotzerihun@yahoo.com. 11 may 2020.

Commented [RA26]: ??? Follow one style

2. Ayed M.H., Aissa A. and Noumi M. A comparative study between the effects of feed inclusion with garlic (*Allium sativum*), cloves and turmeric (*Curcuma longa*) rhizome powder on laying hens' performance and egg quality. *Iran. J. Appl. Anim. Sci.* 2018;19(8):693-701.
3. Behnamifar A., Rahimi S., Karimi-Torshizi M.A., Hasanpour S. and Mohammadzade Z. Effect of thyme, garlic and caraway herbal extracts on blood parameters, productivity, egg quality, hatchability and intestinal bacterial population of laying Japanese quail. *Iran. J. Vet. Med.* 2015; 9(3): 179-187.
4. Bhardwaj R.K. Study on the efficiency of satavari and ashwagandha root powder supplementation on production, reproduction and carcass traits of Japanese quails. Ph. D. thesis submitted to Govind Ballabj Pant University of Agriculture & Technology, Pantnagar, Uttarakhand (2009).
5. BIS. 2007. Requirement for chicken feeds.IS: 1374-2007, New Delhi, India.
6. Budhiraja R.D., Krishan P. and Sudhir S. Biological activity of withanolides. *J. Sci. Industr. Res.* 2000 59:904-911.
7. Card, L.E. and Nesheim, M.C. *Poultry Production* 11th Edition Lea and Febiger, Philadelphia, USA (1972).
8. Curvelo E.R., Geraldo A., Silva L.M., Santos T.A. and Vieira-Filho. Levels of inclusion of curcumin extract and turmeric in diets for semidried hens and their effects on performance and egg yolk coloration. *Proceedings of II IFMG Science Technology Week* (2009).
9. Dilawar M.A., Mun H.S., Rathnayake D., Yang E.J., Seo Y.S., Park H.S. and Yang C.J. Egg quality parameters, production performance and immunity of laying hens supplemented with plant extracts. *Animals* 2021; 11(975): 1-13.
10. Gulati S., Madan, V. K., Singh S., Singh I. and Dusyant. Chemical and phytochemical composition of ashwagandha (*Withania somnifera* L.) roots. *Asian J. Chem.* 2017; 29(8):1683-1686.
11. Holder D.P. and Bradferd M.V. Relationship of specific gravity of chicken eggs to number of cracked eggs and percent shells. *Poult. Sci.* 1979; 58: 250-51.
12. Ibrahim D., Ahmad, S. and Hussain S. Effect of supplementation *Withania somnifera* roots on some egg production and quality traits of heat stressed Japanese quails. *Scientific Papers Series D. Animal Science LIX.* 2016; 59: 200-205.

Commented [RA27]: Follow one style

Commented [RA28]: Should follow similar pattern with other references

13. John J. Therapeutic potential of *Withania somnifera*: A report on phyto-pharmacological properties. *Int. J. Pharm. Sci. Res.* 2014; 5:2131-2148.
14. Kumar S., Berwal R.S., Ramasawroop and Sihag S. Effects of dietary supplementation of ashwagandha (*Withania somnifera*) root powder on production performance, egg quality and blood biochemical constituents of laying hens. *Indian J. Anim. Nutr.* 2020;37(4):352-357.
15. Kumar S., Berwal R.S., Ramasawroop and Singh A. Effects of dietary supplementation of ashwagandha (*Withania somnifera*) root powder on external egg parameters and body weight changes in laying hens. *The Pharma Inn. J.* 2021; 10(9):475-478.
16. Nayal K., Kumar A., Kharvi S., Verma P., Saxena S., Yogesh, Joshi D. and Sharma A. Effect of dietary supplementation of arjun bark powder on production and egg quality parameters in uttar layers *Int. J. Adv. Biochem. Res.* 2024; 8(4):529-532.
17. Saki A.A., Aliarabi H., Siyari S.A.S., Salari J. and Hashemi M. Effect of a phyto-genic feed additive on performance, ovarian morphology, serum lipid parameters and egg sensory quality in laying hen. *Vet. Res. Forum* 2014; 5(4): 287-293.
18. Saraswati, T.R., Manalu, W., Ekastuti, D.R. and Kusumorini, N. The role of turmeric powder in lipid metabolism and its effect on quality of the first quail's egg. *J. Indones. Trop. Anim. Agric.* 2013; 38: 123-130.
19. Singh, P.K., Kumar, A. and Tiwari, D.P. Effects of dietary supplementation of black cumin, garlic and turmeric on the production performance and egg quality of white leghorn hens. *Anim. Nutri. Feed Tech.* 2019; 19: 361-370.
20. Snedecor, G.M. and Cochran, W.G. Statistical methods (8th ed) IOWA State University Press, Ames, IOWA, USA (1994).
21. Yalcin, S., Onbasular, E.E., Reisli, Z. and Yalcin, S. Effect of garlic powder on performance, egg traits and blood parameters of laying hens. *J. Sci. Food Agric.* 2006; 86:1336-1339.