

Original Research Article

Performance of seaweed extracts on yield and quality of black cumin (*Nigella sativa* L.) with organic input management

ABSTRACT

A field experiment was conducted during **rabi** season of the two consecutive years of 2020-21 and 2021-22 at the Horticultural Research Station, Mondouri, BCKV, Nadia, West Bengal with the objective to examine the effectiveness of seaweed extract on yield and quality of black cumin. The treatments **composed** of 9 different doses of seaweed extract granule soil application, liquid foliar spray and control. Among the 10 treatments, T_5 (e.g., SWE granules @4kg ha⁻¹) recorded **highest** average plant height of 29.80 cm, 59.45cm and 73.30 cm, at 60, 90 and 120 DAS and **highest** number of primary branches (7,10) and secondary branches (24,10) plant⁻¹ respectively at 120 DAS. Similarly, **highest** seed yield of 536 kg ha⁻¹ along with different yield attributing characters like **highest** mean fresh pod weight (1.40g), dry pod weight (0.36g) and pod diameter (0.91cm), seed yield plant⁻¹ (2.76g) and 1000 seed weight (2.45g) were recorded under the treatment T_5 . **Highest** B: C ratio (1.97) and highest net return (Rs.52816 ha⁻¹) were also recorded under T_5 during the same period of experimentation. From the above **experiment** it may be concluded that the soil application of **Sea Weed** Extract in granular formulation @ 4kg ha⁻¹ has a positive influence on growth, yield and quality of black cumin.

Keywords: B:C ratio, organic inputs, biostimulants, Black Cumin, **FYM**, SWE, yield, quality

Comment [Y1]: the rabi

Comment [Y2]: to examine

Comment [Y3]: were composed

Comment [Y4]: the highest

Comment [Y5]: the highest

Comment [Y6]: the highest

Comment [Y7]: the highest

Comment [Y8]: 0.36 g

Comment [Y9]: 2.45 g

Comment [Y10]: The highest

Comment [Y11]: experiment,

Comment [Y12]: Seaweed ?

Comment [Y13]: The abbreviation FYM appears only once in the text, in the list of keywords and is not explained. It is necessary to explain it on a non-advised basis because in agriculture it abbreviates the expression Farm Yard Manure, but it also abbreviates a non-academic expression but very used in the common language.

INTRODUCTION

Black cumin, scientifically known as *Nigella sativa* L. is a revered flowering plant native to Southwest Asia, celebrated for its rich historical, cultural, and medicinal significance. Commonly referred to as black seed or kalonji, it boasts a plethora of therapeutic benefits deeply rooted in traditional medicine systems such as Ayurveda, Unani, and traditional Middle Eastern medicine. Its small, crescent-shaped seeds, with their distinct aroma and flavour, are not only prized as a spice in global cuisines but also esteemed for their potent medicinal properties.

Comment [Y14]: Suggestion: manny

Biostimulants are indispensable in organic farming as they provide a holistic solution to the challenges faced by growers striving for sustainable agricultural practices. These natural compounds, sourced from a variety of organic materials like seaweed, microorganisms, and plant extracts, offer multifaceted benefits that extend from the roots of plants to the health of the soil. By improving nutrient uptake efficiency, biostimulants ensure that crops receive essential elements necessary for robust growth and development, ultimately leading to increased yields and improved crop quality. Moreover, they bolster plants' innate defense mechanisms, equipping them to withstand environmental stresses such as drought, extreme temperatures, and pest pressures. Seaweed extract falls under this category which holds significant importance in promoting plant growth and development due to its rich nutritional composition and bioactive compounds. Derived from various species of marine algae, seaweed extract is a natural source of macro and micronutrients, vitamins, amino acids, and growth-promoting hormones such as auxins, cytokinins, and gibberellins. These components play key roles in stimulating root development, enhancing nutrient uptake, and promoting overall plant vigor.

Comment [Y15]: various

Comment [Y16]: defence

Comment [Y17]: falls

Comment [Y18]: vigour

MATERIAL AND METHODS

A field experiment (*Rabi* seasons of 2020-21 and 2021-22) was carried out at the Horticultural Research Station, Mondouri and the laboratory of the Department of Plantation, Spices, Medicinal and Aromatic crops, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal. The experimental material comprised of 9 different doses of seaweed extract (SWE) as foliar spray and soil application as granules given on black Cumin at 30, 60 and 90 days after sowing (T₁-SWE foliar spray@1ml^l⁻¹, T₂-SWE foliar spray @2ml^l⁻¹, T₃-SWE foliar spray @3ml^l⁻¹, T₄- SWE granules @ 3kg ha⁻¹, T₅-SWE granules @4kg ha⁻¹, T₆-SWE granules @ 5kg ha⁻¹, T₇ - SWE foliar spray @1ml^l⁻¹+ SWE granules @3kg ha⁻¹, T₈-SWE foliar spray @2ml^l⁻¹+ SWE granules @ 4kg ha⁻¹, T₉ - foliar spray @ 3ml^l⁻¹ + SWE granules @ 5kg ha⁻¹ and T₁₀ - control) following randomized block design with 3 replications.

The soil of the experiment at the field was Gangetic Alluvial sandy clay loam texture, well-drained, good water holding capacity with moderate soil fertility status (Sand- 54.25%, Silt- 30.20% Clay-14.30%, pH-5.74, Organic carbon (%) - 0.85, N (kg/ha) – 207, P₂O₅(kg/ha)- 380.1, K₂O (kg/ha)- 526.6, S (mg/ha) - 60.18, Zn (mg/ha)- 1.66, Ca (mg/ha)- 949.55, B (mg/ha)-0.44). Well rotten Farm Yard Manure @ 10 t ha⁻¹ was applied at the time of field preparation as basal application and mixed well with the soil, no chemical fertilizers were applied. The entire experimental area was leveled and then divided into 30 plots of 1.8m × 2.5 m size. Black Cumin variety Rajendra Shyama seeds were collected from Rajendra Agricultural University, Bihar and treated with *Trichoderma viride* @ 4g kg⁻¹. Each year the seeds were sown during the first week of November at a spacing of 30 cm x 10 cm @ 6kg ha⁻¹. First irrigation was given immediately after sowing. Subsequent irrigations were given at an interval of 8-10 days depending upon the soil moisture. Necessary treatments were given as per schedule in the respective plots time to time. Randomly selected five tagged plants from

Comment [Y19]: There is no map (classical or GIS) that locates the experimental field and the involved laboratories on the territory of India.

Comment [Y20]: Data on the experimental field are missing:
- isolation measures of plots with different treatments;
- images of the experimental field and plots;
- comparative images of the harvest obtained on plots with different treatments.

Comment [Y21]: In size

Comment [Y22]: Data on the experimental field are missing:
- isolation measures of plots with different treatments;
- images of the experimental field and plots;

Comment [Y23]: from time

each plot were considered for recording the growth parameters like plant height(cm), number of primary and secondary branches and dry matter accumulation (g plant⁻¹). The following yield parameters like fresh and dry pod weight (g), pod diameter (cm), 1000 seed weight(g), plot yield(g) and projected yield (kg ha⁻¹) were also recorded. Harvesting was done during the first fortnight of March of each year based upon the climatic condition of that time. The quality parameters like essential oil content were done by hydro distillation using Clevenger apparatus. The prices of the inputs prevailed at the time of their use were taken into consideration to work out the cost of cultivation. Gross income was calculated by multiplying the seed yield with prevailing market price of seed. Fisher method of analysis of variance as given by Panse and Sukhatme (1967) was applied for analysis and interpretation of data. The level of significance used in 'F' was at P = 0.05 and critical difference (CD) values were worked out wherever 'F' test was significant. Correlations were performed using the Pearson's correlation coefficient (r).

Comment [Y24]: on

Comment [Y25]: conditions

Comment [Y26]: that prevailed

Comment [Y27]: by the prevailing

Comment [Y28]: the seed

Comment [Y29]: Fisher's

Comment [Y30]: The analysis

Comment [Y31]: the 'F'

Comment [Y32]: Pearson's

RESULTS AND DISCUSSION

Table.1: Performance of sea weed extract on plant height, no. of primary, secondary branches and dry matter of black Cumin (Mean of 2 years)

Comment [Y33]: Table 1

Comment [Y34]: seaweed

Comment [Y35]: number

Treatment	Plant height(cm)			Primary branch plant ⁻¹	Secondary branch plant ⁻¹	Dry matter plant ⁻¹ g
	Days after sowing					
	30	60	120			
T ₁	6.20	25.55	65.08	6.20	20.00	15.64
T ₂	5.95	26.35	67.82	6.70	21.30	15.99
T ₃	6.20	27.35	68.98	6.30	21.10	16.16
T ₄	6.65	27.95	69.65	6.30	20.20	16.44
T ₅	5.90	29.80	73.30	7.10	24.10	18.05
T ₆	6.55	27.85	70.15	6.70	21.80	15.75
T ₇	6.50	26.25	68.24	6.00	20.40	17.41
T ₈	6.20	25.90	69.50	6.60	21.30	16.38
T ₉	6.20	28.35	71.55	6.90	23.50	17.95
T ₁₀	5.40	25.10	64.02	5.70	18.70	15.19
SEm(±)	0.19	0.72	0.89	0.12	0.41	0.41
LSD (0.05)	0.57	2.14	2.61	0.36	1.20	1.21

(SWE=Sea weed extract, Liquid=L, Granules=G, T₁- SWE(L) foliar spray @ 1ml/l, T₂ - SWE(L) foliar spray @ 2ml/l, T₃ - SWE (L) foliar spray @ 3ml/l, T₄- SWE (G) @ 3kg/ha, T₅- SWE (G) @ 4kg/ha, T₆- SWE (G) @ 5kg/ha, T₇- SWE(L) foliar

spray @1ml/l+SWE (G) @ 3kg/ha, T₈ - SWE (L) foliar spray @2ml/l +SWE (G) @ 4kg/ha, T₉ - SWE (L) foliar spray @ 3ml/l +SWE (G) @ 5kg/ha and T₁₀– Control).

Growth Parameters

Data presented in Table-1 on plant height of black cumin as influenced by sea weed extract revealed that just before imposing treatments at 30DAS maximum plant height of 6.65 cm was recorded in the plot to be treated with T₄ (SWE granules @ 3kg ha⁻¹) and the lowest plant height of 5.40 cm was recorded in the control plot (T₁₀). At 60DAS, just before application of 2nd dose maximum plant height of 29.80 cm was recorded in the plot treated with T₅(SWE granules @4kgha⁻¹) followed by T₉ (SWE foliar spray @ 3ml l⁻¹+SWE granules @ 5kg ha⁻¹) which was 28.35 cm. At 120 DAS after imposing all the treatments as per schedule, highest plant height of 73.30cm was recorded in T₅ (SWE (G) @ 4kg ha⁻¹) followed by T₉ (71.55cm) and least in control plots (64.02cm). The number of primary and secondary branches plant⁻¹ at 120 DAS of black cumin varied significantly, the maximum number of primary branches plant⁻¹ (7.10) was observed in the plots treated with T₅ and it was on par with T₉ (6.90) while minimum of 5.70 was recorded under control. Maximum (24.10) number of secondary branches plant⁻¹ was recorded in T₅ followed by T₉ (23.50). Similarly, T₅ recorded highest dry matter accumulation of 18.05g plant⁻¹ at 120 DAS and it was *on par* (17.95g) with T₉ whereas control (T₁₀) recorded minimum of 15.19g dry matter plant⁻¹.

Comment [Y36]: seaweed

Comment [Y37]: 30 DAS

Comment [Y38]: 60 DAS

Comment [Y39]: the application

Comment [Y40]: the 2nd

Comment [Y41]: the highest

Comment [Y42]: 73.30 cm

Comment [Y43]: a minimum

Comment [Y44]: The maximum

Comment [Y45]: in plant⁻¹

Comment [Y46]: the highest

Comment [Y47]: the minimum or a minimum

Comment [Y48]: seaweed

Table 2: Performance of sea weed extract on fresh, dry pod weight and diameter of black Cumin pods (Mean of 2 years)

Treatment	Fresh pod weight(g)	Dry pod weight(g)	Pod diameter(cm)
T ₁	1.28	0.27	0.86
T ₂	1.29	0.29	0.89
T ₃	1.22	0.27	0.87
T ₄	1.22	0.26	0.87
T ₅	1.40	0.36	0.91
T ₆	1.27	0.31	0.90
T ₇	1.25	0.30	0.88
T ₈	1.34	0.28	0.89

T ₉	1.39	0.34	0.90
T ₁₀	1.20	0.25	0.85
Sem (±)	0.02	0.02	0.02
LSD (0.05)	0.06	0.06	0.06

(SWE=Sea weed extract, Liquid=L, Granules=G. T₁- SWE(L) foliar spray @ 1ml/l, T₂- SWE(L) foliar spray @ 2ml/l, T₃- SWE (L) foliar spray @ 3ml/l, T₄- SWE (G) @ 3kg/ha, T₅- SWE (G) @ 4kg/ha, T₆- SWE (G) @ 5kg/ha, T₇- SWE(L) foliar spray @ 1ml/l+SWE (G) @ 3kg/ha, T₈- SWE (L) foliar spray @2ml/l +SWE (G) @ 4kg/ha, T₉- SWE (L) foliar spray @ 3ml/l +SWE (G) @ 5kg/ha and T₁₀- Control).

Table 3: Performance of sea weed extract on seed yield plant⁻¹, plot⁻¹, 1000 seed weight,

Comment [Y49]: seaweed

projected yield ha⁻¹, increase in yield and essential oil content in black cumin (Mean of 2 years)

Treatment	Yield plant ⁻¹ (g)	Yield plot ⁻¹ (g)	1000seed weight(g)	Projected Yield (kg/ha ⁻¹)	Increase yield (%)	Essential Oil (%)
T ₁	2.14	198.50	1.94	352.59	7.20	0.90
T ₂	2.41	243.50	1.98	432.88	31.62	1.00
T ₃	2.48	252.00	2.09	448.00	36.21	1.00
T ₄	2.39	279.00	1.91	496.00	50.81	0.90
T ₅	2.76	301.50	2.45	536.00	62.97	0.90
T ₆	2.31	294.50	2.04	523.55	59.19	0.90
T ₇	2.26	213.50	2.17	379.55	15.40	0.90
T ₈	2.35	236.50	1.92	420.44	27.83	0.90
T ₉	2.51	299.00	2.36	531.55	61.62	1.00
T ₁₀	1.92	184.00	1.86	328.88	-	0.80
Sem (±)	0.18	13.44	0.08	23.96	-	0.07
LSD (0.05)	0.53	39.94	0.24	71.18	-	0.20

(SWE=Sea weed extract, Liquid=L, Granules=G. T₁- SWE(L) foliar spray @ 1ml/l, T₂- SWE(L) foliar spray @ 2ml/l, T₃- SWE (L) foliar spray @ 3ml/l, T₄- SWE (G) @ 3kg/ha, T₅- SWE (G) @ 4kg/ha, T₆- SWE (G) @ 5kg/ha, T₇- SWE(L) foliar spray @ 1ml/l+SWE (G) @ 3kg/ha, T₈- SWE (L) foliar spray @2ml/l +SWE (G) @ 4kg/ha, T₉- SWE (L) foliar spray @ 3ml/l +SWE (G) @ 5kg/ha and T₁₀- Control).

Yield Parameters

Perusal of data presented in Table-2 indicated that highest fresh pod weight of 1.40g was recorded in the treatment T₅ followed by T₉ (1.39g) which are *at par* with each other and minimum pod weight of 1.20g was recorded in control plot (T₁₀) only. Highest dry pod weight of 0.36g was recorded in the treatment T₅ followed by T₉ (0.34g) which are *at par* with each other and the least pod weight of 0.25g was recorded in control plot.

Comment [Y50]: Table 2

Comment [Y51]: the highest

Comment [Y52]: the minimum

Comment [Y53]: The highest

Comment [Y54]: 0.36 g

Comment [Y55]: 0.34 g

Comment [Y56]: lowest

Comment [Y57]: 0.25 g

Comment [Y58]: the control

Comment [Y59]: seaweed

Comment [Y60]: Table 3

The effect of sea weed extract on seed yield plant⁻¹ as presented in the Table-3 indicated that among the different treatments, T₅ recorded a significantly higher seed yield of

2.76g plant⁻¹ followed by T₉ (2.51g plant⁻¹) and the lowest seed yield of 1.92 g plant⁻¹ was recorded in T₁₀. Seed yield plot⁻¹ indicated that among the different treatments, T₅ recorded a maximum seed yield of 301.50 g plot⁻¹ followed by 299.00 g plot⁻¹ was recorded in T₉ while, lowest seed yield of 184.00 g plot⁻¹ was recorded in the T₁₀ and T₅ recorded a significantly highest seed weight of 2.45g (1000 seeds) followed by T₉ (2.36g) and lowest of 1.86g (1000 seeds) in the treatment T₁₀. Projected seed yield ha⁻¹ indicated that the highest projected seed yield of 536 kg ha⁻¹ was recorded in the treatment T₅ followed by T₉ (531.55kg ha⁻¹) while, the lowest projected seed yield of 328.88 kg ha⁻¹ was noted in the treatment T₁₀ and T₅ recorded the highest increase in yield of 62.97% followed by T₉ (61.62%). There were no significant differences in essential oil (%) content among the treatments. Among the treatments essential oil content ranges between 0.80-1.00% out of which the highest essential oil content 1.00% was recorded in the treatments T₂, T₃ and T₉ whereas in all the other treatments it was recorded 0.90 % except T₁₀ with 0.80%.

Comment [Y61]: 2.76 g

Comment [Y62]: while lowest

Comment [Y63]: in treatment

Comment [Y64]: while

Comment [Y65]: treatment

Comment [Y66]: 62.97 %

Comment [Y67]: 61.62 %

Comment [Y68]: treatments,

Comment [Y69]: at 0.90 %

Comment [Y70]: 0.80 %.

Comment [Y71]: seaweed

Comment [Y72]: the economics

Table 4: Performance of sea weed extract on economics of black cumin cultivation

Treatment	Cost of cultivation (Rs. ha ⁻¹)	Gross income (Rs. ha ⁻¹)	Net return (Rs. ha ⁻¹)	B:C ratio
T ₁	57523	70518	12995	1.23
T ₂	60372	86576	26204	1.43
T ₃	63222	89600	26378	1.42
T ₄	53994	99200	45206	1.84
T ₅	54384	107200	52816	1.97
T ₆	54804	104710	49906	1.91
T ₇	62748	75910	13162	1.21
T ₈	65988	84088	18100	1.27
T ₉	69258	106310	37052	1.53
T ₁₀	48768	65776	17008	1.35

(SWE=Sea weed extract, Liquid=L, Granules=G, T₁- SWE(L) foliar spray @ 1ml/l, T₂- SWE(L) foliar spray @ 2ml/l, T₃- SWE (L) foliar spray @ 3ml/l, T₄- SWE (G) @ 3kg/ha, T₅- SWE (G) @ 4kg/ha, T₆- SWE (G) @ 5kg/ha, T₇- SWE(L) foliar spray @ 1ml/l+SWE (G) @ 3kg/ha, T₈- SWE (L) foliar spray @ 2ml/l +SWE (G) @ 4kg/ha, T₉- SWE (L) foliar spray @ 3ml/l +SWE (G) @ 5kg/ha and T₁₀- Control).

Economics

Economics is an important and ultimate factor that decides the beneficial effect of imposing different **treatment** on the ultimate economic returns of the farmers. The data presented in Table 4 on the economics of black cumin cultivation indicated that T₅ recorded **highest** B: C ratio of 1.97 with a **highest** net return of Rs. 52816 ha⁻¹ followed by T₆ (1.91 BCR and Rs. 49,906 ha⁻¹).

Comment [Y73]: treatments

Comment [Y74]: the highest

Comment [Y75]: the highest




Figure-1. Correlation coefficient between traits of black cumin

Correlation Analysis between Yield and Agronomic Traits

Comment [Y76]: Between

The correlation analysis was performed between seed yield, plant height, number of primary and secondary branches, dry matter accumulation, dry pod weight, seed weight (1000) and **essential of content**. Yield showed a positive correlation with plant height and number of branches. It showed **negative** correlation with essential oil content. All the growth parameters showed **negative** correlation with essential oil content. Test weight had a negative correlation with yield.

Comment [Y77]: essential content

Comment [Y78]: a negative

Comment [Y79]: a negative

DISCUSSION

These results are in good agreement with previous studies in which **growth** and seedling vigor of mungbean and maize were enhanced by SWE treatment (Rengasamy *et al.*, 2015; Kannan *et al.*, 2016). The positive effect of SWE on cabbage may be due to the result

Comment [Y80]: the growth

of combined interactions between SWE and endogenous growth hormones (auxins and cytokinins) (Teixeira da Silva *et al.*, 2013). In **other way** we can say that organic molecules such as organic acids, methionine and even amino acids in SWE can increase nutrient absorption in plants by chelating the available nutrients, thereby increasing their absorbance (Papenfus *et al.*, 2013). Seaweed extracts can equilibrate growth as a result of auxin and gibberellin acid presence, which will increase **vitamins** and hormones **producing** in the treated plants (O' Dell, 2003). This **increase** Macronutrients like nitrogen, potassium, and phosphorous are very essential for **growth** and development of the plant (Attememe, 2009). It has been indicated that seaweed extract stimulates the mycorrhizal activities in the soil which plays a crucial role in **absorption** of water and nutrients, alleviating any shock. The seaweed extracts which promoted root and plant growth through uptake of nutrients may be due to the presence of plant growth regulators at low concentrations such as cytokinins and auxins, polyamines such as putrescine and spermine which had synergistic action with other plant growth regulators especially auxin (Almohammed *et al.*, 2014). The observation of **SWE treated** *Arachis hypogea* plants suggested that the growth and biochemical characteristics might be promoted by micro and macro elements and **growth promoting** hormones present in the SWE (Ganapathy-selvam and Sivakumar, 2014). Our findings coincide with those of earlier studies carried out on **soybean** (Rathore *et al.*, 2009) where there was an increase in vegetative growth by the application of SWE. Similar results were also observed in *Cajanus cajan* (L.) Mill sp. (Mohan *et al.*, 1994), *Vigna sinensis* L. (Sivasankari *et al.*, 2006), and *Abelmoschus esculentus* (Thirumaran *et al.*, 2009). Seaweed extracts contain cytokinins as well, which induce **the physiological** activities and increase the total chlorophyll in the plant. Also, this will positively reflect on the activity of photosynthesis and the synthesized materials, which will show on **shoots** characteristics. The yield of a particular crop may be due to the function of a gene, environment (climate and soil) and management practices

Comment [Y81]: another way,

Comment [Y82]: the vitamins

Comment [Y83]: produced

Comment [Y84]: O'Dell

Comment [Y85]: increases

Comment [Y86]: the growth

Comment [Y87]: the absorption

Comment [Y88]: SWE-treated

Comment [Y89]: growth-promoting

Comment [Y90]: Ganapathy-Selvam ?

Comment [Y91]: soybeans

Comment [Y92]: physiological

Comment [Y93]: shoot

adopted. Further, yield of the crop depends majorly on the accumulation of photo assimilates during the growth period and in what way they are partitioned between storage organs and vegetative parts of the plant. Photosynthetic ability, biomass accumulation, translocation efficiency and finally the economic yield and quality of the crops are highly influenced by organic matter, fertility and other soil characteristics which were improved by the sea weed extract. This superiority of yield attributes could have been because of improving the plant height, branches plant⁻¹ and dry weight by induction of biochemicals that transport from the vegetative part to the reproductive organs of the plant which ultimately reflects on yield components of the plant. Regarding the biochemical traits application of SWE showed a significant effect on the content of plant essential oil compared to the control. This result is in a good line with the findings of Golzadeh *et al.* (2011) on *Matricaria recutita* and Rafiee *et al.* (2013) on *Calendula officinalis* L. Ardebili *et al.* (2012) indicated that foliar application of SWE as a source of amino acids, at suitable concentrations, had positive effects on the content of secondary metabolites, antioxidants, and antioxidant activity. Naboulsi *et al.* (2018) indicated that secondary metabolites such as essential oils extracted from the plants, showed interesting biological activities against numerous harmful pests and microorganisms. In addition, some of the secondary metabolites have a beneficial effect as plant growth regulators which will help to increase growth and yield. Here the highest essential oil content (1.0%) was recorded in treatments T₂, T₃ and T₉ indicated that seaweed extract in different formulations either single or in combination may influence the essential oil content of black cumini by increasing the photosynthetic ability, biomass accumulation and translocation efficiency resulted in higher oil content. The increase in essential oil content due to the effect of SWE was also observed in sunflower by Osman and Salem (2011).

Comment [Y94]: the yield or a yield

Comment [Y95]: photoassimilates

Comment [Y96]: seaweed

Comment [Y97]: good

Comment [Y98]: indicating

Comment [Y99]: sunflowers

CONCLUSION

Comment [Y100]: The conclusions are few. If possible, they should be enriched.

It may be concluded that growth, yield and quality of black cumin can be enhanced by soil application of Sea Weed Extract in granular formulation@ 4 kg ha^{-1} .

Comment [Y101]: The conclusion is not certain. It can be completed with the necessary directions for further research for possible confirmations or increasing the probability of the capacity of the proposed solutions to improve cumin harvests in the respective areas.

Comment [Y102]: seaweed extract

Comment [Y103]: kg ha^{-1}

REFERENCES

Almohammedi, A. N., Almehemdi, A, F. and Al Ajeelee, R. K. 2014. Impact of Bat Guano *Otonycterishemp richii* and seaweed extract on some growth and yield traits of Baraka seed (*Nigella sativa*L.). *J. Biol. Agri. Healthcare*, **4**: 57-65.

Ardebili, Z. O., Moghadam, A. R. L., Ardebili, N. O. and Pashaie, A. R. 2012. The Induced Physiological Changes by Foliar Application of Amino Acids in *Aloe vera* L. *Plants. Plant Omics J.*, **5(3)**: 279 - 84.

Attememe, J. Y. A. 2009. The Effect of Humic Acid and Sea Weed Extracts on the Growth, Chemical Characteristics and Oil Characteristics of *Rosmarinus officinalis* L. The 6th Scientific Conference, Biology Dept., College of Education, University of Tikrit, *Plants Sci.*, PP. 1-17.

Bulgari, R., Franzoni, G. and Ferrante, A. 2019. Biostimulants application in horticultural crops under abiotic stress conditions. *Agronomy*, **9**:1-30.

Comment [Y104]: I did not find the citation in the text of the manuscript.

Ganapathy-selvam, G. and Sivakumar, K. 2014. Influence of Seaweed Extract as an Organic Fertilizer on the Growth and Yield of *Arachis hypogaea* L. and Their Elemental Composition Using SEM-Energy Dispersive Spectroscopic Analysis. *Asian. Pac. J. Reprod.*, **3(1)**:18-22.

Golzadeh, H., Mehrafarin, A., Naghdi Badi, H., Fazeli, F., Qaderi, A. and Zarinpanjeh, N. 2011. Effect of Biostimulators Compounds on Quantitative and Qualitative Yield of German Chamomile (*Matricaria recutita* L.). *Euro. J. Med. Plants*, **8(1)**: 195 - 208.

Huchhannavar, S., Yogesh, L.N. and Prashant, S. M. 2019. The black seed *Nigella sativa*: A wonder seed. *Int. J. Chem. Studies*, **7**:1320-24.

Comment [Y105]: I did not find the citation in the text of the manuscript.

Kannan, R. R., Rengasamy, Manoj,G., Kulkarni, G., Srinivasa, C. and Van Staden, J. 2016. Enhancing Growth, Phytochemical Constituents and Aphid Resistance Capacity in Cabbage with Foliar Application of Eckol: A Biologically Active Phenolic Molecule from Brown Seaweed. *New Biotech.*, **33**(2): 273-280.

Mohan, V. R., Venkataraman Kumar, R., Murugeswari, R. and Muthusamy, S. 1994. Effect of Crude and Commercial Seaweed Extracts on Seed Germination and Seedling Growth in *Cajanus cajan*. *L. J. Phycol. Soc.*, **33**: 47-51.

Naboulsi, I., Aboulmouhajir, A., Kouisni, L., Bekkaoui, F. and Yasri, A. 2018. Plants extracts and secondary metabolites, their extraction methods and use in agriculture for controlling crop stresses and improving productivity: A review, *Academia J. Med. Pl.*,**68**: 223-40.

Comment [Y106]: use in

O' Dell, C. 2003. Natural Plant Hormones Are Bio Stimulants Helping Plants Develop High Plant Antioxidant Activity for Multiple Benefits. *Virginia Vegetable Small Fruit and Specialty Crops*, **2**(6):1-3.

Osman, H. E. and Salem, O. M. A. 2011. Effect of seaweed extracts as foliar spray on sunflower yield and oil content. *Egyptian J. Phycol.*,**12**:57-70.

Panse, V. G. and Sukhatme, P. V. 1967. Statistical Methods for Agricultural Workers, ICAR, New Delhi, pp. 152-174.

Papenfus, H. B., Kulkarni, M. G., Stirk, W. A., Finnie, J. F. and Van Staden, J. 2013. Effect of a Commercial Seaweed Extract (Kelpak®) and Polyamines on Nutrient Deprived (N, P and K) Okra Seedlings. *Scientia Hortic.*, **151**: 142-146.

- Rafiee, H., Mehrafarin, A., Qaderi, A., Kalate Jari, S. and Naghdi Badi, H. 2013. Phytochemical, Agronomical and Morphological Responses of Pot Marigold (*Calendula officinalis* L.) to Foliar Application of Bio-stimulators (Bioactive amino Acid Compounds). *J. Med. Plants. Res.*, 12(47): 48 - 61.
- Rathore, S. S., Chaudhary, D. R., Boricha, G. N., Ghosh, A., Bhatt, B. P., Zodape, S. T. and Patolia J. S. 2009. Effect of Seaweed Extract on the Growth, Yield and Nutrient Uptake of Soybean (*Glycine max*) under Rainfed Conditions. *S. Afr. J. Bot.*, 75: 351-355.
- Rengasamy, K. R. R., Kulkarni, M. G., Stirk, W. A. and Van Staden, J. 2015. Improves Growth, Enzyme Activities, and Secondary Metabolite Content in Maize (*Zea mays* cv., Border King). *J. Plant. Growth. Regul.*, 34:410-6.
- Sivasankari, S.V., Venkatesalu, M., Anantharaj, A., and Chandrasekaran, M. 2006. Effect of Seaweed Extracts on the Growth and Biochemical Constituents of *Vigna siensis*. *Bioresour. Technol.* 97: 1745- 1751.
- Teixeira da Silva, J. A., Dobrańszki, J. and Ross, S. 2013. Phloroglucinol in Plant Tissue Culture. *In Vitro Cell Dev. Biol. Plant.*, 49:1-16.
- Thirumaran, G., Arumugam, M., Arumugam, R. and Anantharaman, P. 2009. Effect of Seaweed Liquid Fertilizer on Growth and Pigment Concentration of *Abelmoschus esculentus* (l) Medikus. *Am. Eurasian J. Agron.*, 2 (2): 57-66.