

***In vitro* development of two *Alternaria solani* strains, causal agent of Alternariose in tomato (*Lycopersicon esculentum*) under the influence of *Thevetia peruviana* seeds extracts**

Abstract

Alternaria solani is a fungus that causes yield losses of up to 80 % in tomato production in field. Synthetic fungicides are the most widely used for its control, but have harmful consequences. The objective of this work was to test *in vitro* the antifungal potential of *Thevetia peruviana* seed extracts against two *A. solani* isolates. Aqueous, methanol, ethyl acetate and acetone extracts, at concentrations 12.5, 25, 50 and 100 $\mu\text{L/mL}$ were used. Two synthetic fungicides Maneb (5.33 $\mu\text{g/mL}$) and Dimethomorph + Clorothalonil (3.75 $\mu\text{g/mL}$) and control (0 $\mu\text{L/mL}$) were also tested on two *A. solani* isolates (Mbal and Foun). The investigation was repeated three times. Phytochemical screening, mycelial growth, spore germination and minimum inhibitory concentrations (MIC50 and MIC90) were determined. The results showed that *T. peruviana* extracts are rich in many families of bioactive compounds such as alkaloids, phenolic compounds and sugars. All extracts tested show high inhibition of mycelial growth (100%) and spore germination (100%) of the two strains at highest concentration (100 $\mu\text{L/mL}$). Acetone extract at a concentration of 50 $\mu\text{L/mL}$ inhibited mycelial growth by 88.45 and 86.55% and spore germination by 88.33 and 80.33%, respectively for the Mbal and Foun isolates. The lowest MIC50 (16.63 $\mu\text{L/mL}$) and MIC90 (54.6 $\mu\text{L/mL}$) were obtained with the acetone extract on the Mbal isolate while the highest MIC50 (27.5 $\mu\text{L/mL}$) and MIC90 (61.7 $\mu\text{L/mL}$) were observed with ethyl acetate on the Foun isolate. These extracts can therefore be used in the biological control against Alternariose in tomato.

Keywords: *Thevetia peruviana*, extract, isolates, *Alternaria solani*, inhibition.

• INTRODUCTION

Tomato (*Lycopersicon esculentum* Mill.) is a plant belonging to the Solanaceae family, native to South America and cultivated in more than 177 countries [1]. Tomato is the most consumed vegetable in the world ahead of watermelon and cabbage [2]. It is

consumed as salad, puree, concentrate, condiment and sauce [3]; [4]. Tomato has a high nutritional and economic importance, and its fruit is composed of 95% water and 5 % dry matter[37-40].

World production in 2020 is estimated at about 186.8 million tons of fresh fruits on a cultivated area of 5,051.9 thousand hectares with an average yield of 37.1 tons per hectare [5], while total African production is estimated at about 17,175,228 tons. In 2020, Cameroon recorded a production of 1,246,658 tons on 101,350 ha, i.e. about 14 % of African production. This low tomato productivity is due not only to the influence of abiotic factors (temperature, humidity, water stress, etc.) but also to the strong pressure of biotic factors such as diseases and pests [6]. Among the devastating diseases of tomato fields, Alternariosis, a fungal disease caused by *Alternaria solani*, is one of the most important [7]. Alternariosis is manifested by severe and destructive blights, reduced germination capacity of seeds, defoliation of the plant, and deterioration of the products before and after harvest, resulting in yield losses of up to 80 % in field tomato production. [8]. Several strategies of combating this disease are used, including chemical control, which remains the most used method by producers because it is easy to use. However, these synthetic chemical products have an impact on the health of producers, consumers and the environment, and leads to the development of resistance by certain pathogens [2]. Considerable efforts are directed towards the search for effective alternative solutions that respect the environment and human health [9]. Numerous studies have demonstrated the importance of using plant extracts from *Azadirachta indica*, *Thevetia peruviana* rich in secondary metabolites (phenolic compounds, terpenoids and alkaloids) in the control of crop diseases [10].

Thevetia peruviana (yellow oleander) which belongs to the Apocynaceae family, is an ornamental shrub native to tropical America and is widely cultivated throughout the tropics including tropical Africa [11]. The roots, leaves, seeds and fruits of *T. peruviana* are sources of biological active compounds conferring bactericides [12], virucides [13], insecticides [14], [15] and fungicides [16-17]; [9], [18] potentials.

The present work aims to test *in vitro* the antifungal potential of *Thevetia peruviana* seed

extracts against two *Alternaria solani* isolates

- **MATERIAL AND METHODS**

- **Sample collection**

The leaves showing typical symptoms of the disease were collected in two agro-ecological zones of Cameroon, one in the bi-modal rain fall forest zone, more precisely in the locality of Mbamayo (Mbal) belonging to agro-ecological zone 5 with geographical coordinates (N5.2926, E10.3419). The other in the highland or humid savannah zone in the locality of Foubot (Foum) in agro-ecological zone 3 with geographical coordinates (N3.3109, E11.3010) in Cameroon. These leaves were placed in envelopes and then put in a cooler containing ice cubes and transported directly to the Biotechnology and Environment Laboratory, Plant Pathology and Protection Research Unit of the University of Yaounde 1.

Thevetia peruviana seeds were obtained from fruits collected under trees in Yaoundé and identified in the national herbarium.

The plant material consisted mainly of *T. peruviana* seeds while fungal material used was two *A. solani* isolates (Mbal and Foum) isolated from diseased organs in farmers' fields, in Mbalmayo and Foubot. Two synthetic fungicides including Plantineb 80 WP (Maneb) and Jumper D (Dimethomorph + Clorothalonil) were used as reference. Laboratory equipment was also used.

- **Obtaining of *Alternaria solani* strains**

Using the identification key of [19] symptomatic leaves were collected and transported directly to the laboratory of Biotechnology and Environment, University of Yaoundé I. These leaves were carefully washed with running tap water, cut into small fragments, immersed in 10 ml of alcohol at 5°C for 2 min and rinsed 3 times with sterile distilled water.

These leaf fragments were dried on sterile filter paper (Whatman N° 1) and inoculated into Petri dishes containing PDA (Potato Dextrose Agar) then sealed with film and incubated at 23 ± 2 °C until colonies appeared. After 5 days of incubation, the mycelia were removed

from the growth front of the fungus and transferred to new Petri dishes containing PDA. This last operation was repeated several times until pure strains were obtained, and their microscopic observations were performed following the protocol described by [20].

- **Obtaining of different extracts and their phytochemical screening**

The *T. peruviana* seeds were collected in Yaounde. The obtained kernels were dried in the laboratory at room temperature for two weeks and grinded using manual hand mill grinder. The aqueous extract was obtained by maceration of 100 g of paste in 200 mL of distilled water for 24 hours. The resulting mixture was filtered through a muslin cloth and used directly [16]. For organic extracts, 500 g of powder were macerated in 2 L of each organic solvent (ethyl acetate, acetone and methanol) for 72 hours. The mixture was filtered using filter paper and the filtrates were concentrated using rotavapor (Büchi R124) at a temperature of 60-70°C. The organic extracts obtained were stored in a refrigerator at 4 °C until their use.

Phytochemical screening of *T. peruviana* extracts was carried out to determine the composition of the major families of secondary metabolites contained in these plants using classical characterisation methods and specific dyes. Protocol used for phytochemical screening was achieved according to [21] ; [22].

- **Evaluation of the effect of the extracts on the mycelial growth of *A. solani* strains**
- **Preparation of the different concentrations tested**

A stock solution of 500 µL/mL was prepared by mixing 12 mL of extract organic (methanol, ethyl acetate, acetone) with 3 mL of 70°C alcohol added to 9 mL of distilled water. Increasing concentrations of (C=12.5; C2=25; C3=50 and C4=100 µL/mL) were prepared by taking 1.5; 3; 6 and 12 mL of each extract respectively and adding them to 58.5; 57; 54 and 48 ml of the PDA to give a final volume of 60 mL. The antibiotic-amended mixture is aseptically poured into 90 mm diameter Petri dishes at a rate of 10 mL per dish under the laminar flow hood. For the negative controls, 10 mL of medium without additive was poured directly into the Petri dishes. The media preparation was enriched with two synthetic fungicides (Plantineb 80 WP and Jumper D) at concentrations

of C5=5.33 µg/mL and C6=3.75 µg/mL.

- **Evaluation of mycelial growth**

The Explants about 6 mm in diameter of each *Alternaria solani* strains were collected and placed in the centre of each Petri dish containing PDA medium enriched with the different extracts. Each treatment was repeated three times and the whole set was incubated at a temperature of 23±1 °C under continuous light for one week. Measurements of mycelia growth were taken at approximately the same time each day from the second day until the control (C0=0) plates were completely filled with mycelium. The mycelia growth of both isolates was assessed by measuring the two perpendicular diameters plotted on the back of the Petri dish and calculated according to the formula used by [23].

$$D = \frac{D1 + D2}{2} - D0$$

Where D0 is the diameter of the explant; D1 and D2: culture diameters measured in the two perpendicular directions.

The inhibition percentage due to each treatment is assessed in relation to the mycelia growth in the control Petri dishes after 7 days according to the formula developed by [24].

$$IP (\%) = \frac{Dt (mmm) - Dx (mm)}{Dt (mmm)} \times 100$$

IP (%): Inhibition percentage; Dt: estimated growth diameter on control medium and Dx: estimated growth diameter in the presence of the tested extract or fungicide.

- **Determination of minimum inhibitory concentrations (MIC)**

The minimum inhibitory concentrations reducing mycelia growth by 50% and 90% were determined by comparing the values of the percentage inhibition (PI) with those of the natural logarithm of the corresponding concentrations [23].

The linear regression line of type $Y = ax + b$ from the function $IP = f(\ln Ci)$ was used

where Y = percentage inhibition of mycelia growth or spore germination, a = regression coefficient, b = constant and x = extract concentrations.

- **Evaluation of the effect of *T. peruviana* seed extracts on *Alternaria solani* spore germination**

From a 7-day old culture of the pathogen of both isolates, 20 µl solutions of a spore suspension calibrated at 4,105 spores/ml using a Malassez cell were deposited and spread with a micropipette on the slides. Each treatment was repeated 3 times. All the slides were kept in the dark for 24 hours and the parameters were taken on the basis of counting 100 germinated spores on three different zones of each slide under an ordinary microscope, i.e. a total of 300 spores per slide and 900 spores for each treatment. The calculation of the percentages of inhibition was done using the formula of [25] and [23].

$$IP = \frac{A - B}{A} \times 100$$

With IP: Inhibition percentage; A: number of spores germinated on the control; B: number of spores germinated in the presence of the test extract.

- **Evaluation of the fungicidal or fungistatic activity of *Thevetia peruviana* seed extracts**

At the end of each trial, the mycelia explants from the Petri dishes where growth was totally inhibited were removed and aseptically placed on the PDA culture medium containing no extract. After 7 days of incubation, depending on whether the fungus had resumed growth or not, the starting extract was classified as fungistatic or fungicide respectively [26].

- **Statistical analysis**

Data on mycelia growth of isolates and spore germination rate were subjected to an analysis of variance (ANOVA) using R software version 3.5.1. Tables and curves were drawn using Microsoft Excel.

- **RESULTS**

- **Characteristics of *Alternaria solani* isolates**

Macroscopic observation of the pure isolate obtained showed whitish colonies with a downy or cottony subaerial appearance with variations in mycelia growth and regular and irregular borders (Figure 1A). Microscopic observation showed a species with large spore conidia with a porri (Figure 1B) cross-section, with 1-2 filamentous spouts measuring between 12.8 and 121.6 mm; the length of the conidial body varying from 54.4 to 115.2 mm. These different forms identified refer to those of *Alternaria solani*, the causal agent of Alternariose in tomato.




Figure 1. Macroscopic and microscopic characteristics of pure isolates of *Alternaria solani* (A- Macroscopic aspect; B- Conidia seen by photonic microscopy at X 20 magnification).

- **Effect of *Thevetia peruviana* seed extracts on the *in vitro* radial growth of *Alternaria solani***
- **Phytochemical screening of *Thevetia peruviana* seed extracts**

The phytochemical screening carried out revealed the existence of several families of compounds such as: essential and saponified oils, coumarins, sterols, alkaloids, saponins, anthocyanins, steric glycosides and sugars. Coumarins and sterols were present in all extracts and phenols were absent in the aqueous extract, ethyl acetate and in trace

amounts in the methanol extract (Table 1).



Table 1. Family of compounds in the aqueous and organic extracts of *T. peruviana* seeds.

(-): absence of the product; (+): presence of the product; (+++): abundant presence of the product and T: presence in trace form.

Group of compounds	Results of the different extracts			
	Aqueous	Ethyl acetate	Acetone	Methanol
Essential oils	+	-	+	+
Saponifiable oils	+	+	+	-
coumarin	+	+	+	+
alkaloids	+	-	-	+
sterols	+++	+	+	+
Terpenoids	-	-	T	-
Flavonoids	-	-	-	-
Anthraquinones	+	-	-	-
Catechin tannins	-	-	-	+
Gallic tannins	-	-	-	-
Saponins	-+	-	+	+
Anthocyanins	-	+	-	+
Steric glycosides	+	T	+	+
Triterpenoid glycosides	-	-	-	T
Sugars	+++	-	T	T
phenols	-	-	-	T

- **Radial growth of individual isolates**

All extracts significantly reduced the radial growth of the different isolates. Mycelial growth decreased with increasing concentration of the different extracts until it was zero with the highest concentration (C4: 100 $\mu\text{L}/\text{mL}$ or $\mu\text{g}/\text{mL}$) as well as with the synthetic fungicides (C5 and C6) used based on Maneb and Dimethomorph + Clorothalonil (Figure 2).






Figure 2. Evolution of mycelia growth of *A. solani* isolates as a function of different extract concentrations on the 7th day after incubation (A- Aqueous; B- Methanol; C- Ethyl acetate and D- Acetone).

- **Inhibition percentage of the different extracts of *Thevetia peruviana* seeds**

The inhibition percentage of mycelia growth of *A. solani* varied with the increase of the different extracts. At 7 days post-inhibition (DPI), the acetone extract at concentration C3 (50 $\mu\text{L}/\text{mL}$) showed the highest inhibition percentages compared to the control, respectively 88.45 % for the Mbal strain and 86.55 % for the Foun isolate, followed by the aqueous extract (with 85.32 % for the Mbal isolate and 83.33 % for that of Foun isolate). On the other hand, at the highest concentration C4 (100 $\mu\text{L}/\text{mL}$), total inhibition of mycelia growth (100 %) was observed for the aqueous and organic extracts and the synthetic fungicides used (Figure 3).

* Means in the same column followed by the same letters are not significantly different according to the Tukey test at the 5% probability level;

Figure 3. Variation in the percentage of radial growth inhibition of isolates by the different extracts (A: Mbal; B: Foum) 7 JAI

- **Minimum inhibitory concentrations MIC50 and MIC90 of extracts**

The minimum inhibitory concentrations (MICs) for growth of *A. solani* isolates varied between extracts. The lowest minimum concentrations that effectively inhibited mycelia growth at 50% and 90% were recorded with the acetone extract where MIC50 and MIC90 were respectively 16.63 L/mL and 54.6 L/mL for the Mbal isolate. With the same extract, MIC50 and MIC90 were 18.63 L/mL and 55.6 μL/mL for the Foum isolate. The highest minimum concentrations were recorded with the ethyl acetate extract where MIC50 and MIC90 were respectively 26.63L/mL and 60 μL/mL for the Mbal isolate. With the same extract, MIC50 and MIC90 were 27.5 L/mL and 60.7 L/mL, respectively for the Foum isolate (Table 2).

Table 2. MIC50 and MIC90 values on mycelia growth of *Alternaria solani* according to aqueous and organic extracts and synthetic fungicides

Isolates	Extracts	MIC50	MIC90
Mbal	Aqueous (L/mL)	17.6	55.6
	Methanol (L/mL)	22.25	55.7
	Ethyl acetate (L/mL)	26.63	60
	Acetone (L/mL)	16.63	54.6
Foum	Aqueous (L/mL)	20.13	57
	Methanol ((L/mL)	23.5	60
	Ethyl acetate ((L/mL)	27.5	60.7
	Acetone (L/mL)	18.63	55.6

- **Correlation between the concentrations of the different extracts and the percentage of growth inhibition of *Alternaria solani***

The regression equations obtained from the tests with the different extracts showed increasing linear relationships with regression lines all with positive slopes. The correlation coefficient (r) varied from 0.97 to 0.99, i.e. $r > 0.8$ (Table 3). This shows a positive and

perfect correlation between the different concentrations tested and the percentage of radial growth inhibition. The latter is proportional to the different increasing concentrations tested.

Table 3. Correlation between inhibition percentage and the concentrations on the 2 isolates tested with the different extracts of *T. peruviana*

Isolates	Extracts	Correlation coefficient	Observations
Mbal	Aqueous	0.981	Highly correlated
	Methanol	0.996	Highly correlated
	Ethyl acetate	0.988	Highly correlated
	Acetone	0.953	Highly correlated
Foum	Aqueous	0.986	Highly correlated
	Methanol	0.997	Highly correlated
	Ethyl acetate	0.988	Highly correlated
	Acetone	0.973	Highly correlated

- **Effect of *Thevetia peruviana* seed extracts on spore germination of different isolates of *Alternaria solani***

The acetone extract had a very significant influence on the germination of spores of the two isolates compared to the other extracts from the first concentration with inhibition percentages of 56.67; 71.67; 88.33 and 100 % for the Mbal isolate, while 50.00; 68.33; 80.00 and 100 % for were recording the Foum isolate for C1, C2, C3 and C4 respectively. On the other hand, inhibition percentage of the ethyl acetate extract was lower compared to the other extracts for both isolates with 21.6; 41.67; 66.67 and 100 % for Mbal and 18.33; 35.00; 63.33 and 100 % for Foum respectively for the same concentrations (Figure 4).

* Means in the same column followed by the same letters do not differ significantly according to the Tukey test at the 5% probability level;

Figure 4. Inhibition percentage of *Alternaria solani* spore germination according to the different *T. peruviana* extracts (A: Mbal isolate and B: Foun isolate)

- **Minimum inhibitory concentrations MIC50 and MIC90 for germination of *Alternaria solani* isolates**

The minimum germination inhibitory concentrations of *A. solani* isolates varied between extracts. The minimum concentrations for spore germination at 50 % and 90 % were lowest with the acetone extracts where MIC50 and MIC90 were respectively 6.38 L/mL and 53.7 L/mL for the Mbal isolate. With the same extract MIC50 and MIC90 were 17.6 L/mL and 56.83 L/mL for the Foun isolate. In contrast, the highest minimum concentrations were recorded with the ethyl acetate extract where MIC50 and MIC90 were respectively 27.63 L/mL and 62.5 L/mL for the Mbal isolate. With the same extract MIC50 and MIC90 were 31.2 L/mL and 66.1 L/mL for the Foun isolate (Table 4).

Table 4. Minimum inhibitory concentrations of *Alternaria solani* spore germination in the presence of *Thevetia peruviana* extracts

Isolates	Extracts	MIC50	MIC90
Mbal	Aqueous (g/mL)	17.6	57.83
	Methanol (L/mL)	22.9	60.5
	Ethyl acetate (L/mL)	27.63	62.5
	Acetone (L/mL)	6.38	53.7
Foun	Aqueous (g/mL)	22.4	60.7
	Methanol (L/mL)	25.4	62.5
	Ethyl acetate (L/mL)	31.2	66.1
	Acetone (L/mL)	12.13	57.5

- **Correlation between different concentrations and germination percentages of *Alternaria solani* isolates**

The regression equations obtained with the different extracts from the germination test of the two isolates used showed increasing linear relationships with regression lines with positive slopes. Furthermore, the correlation coefficients r obtained were between 0.96 and 0.99. Thus, the inhibition percentage is proportional with the increase of the

concentrations of the tested extracts (Table 5).

Table 5. Correlation between the germination percentage and the concentrations of the different extracts tested with *Thevetia peruviana* extracts

Isolates	Extracts	Correlation coefficient	Observations
Mbal	Aqueous	0.980	Highly correlated
	Methanol	0.993	Highly correlated
	Ethyl acetate	0.986	Highly correlated
	Acetone	0,954	Highly correlated
Foum	Aqueous	0.961	Highly correlated
	Methanol	0.979	Highly correlated
	Ethyl acetate	0.973	Highly correlated
	Acetone	0.990	Highly correlated

- Fungicide or fungistatic activity of *Thevetia peruviana* seed extracts**

The *Alternaria solani* isolates tested were very sensitive to the different concentrations of extracts tested. For all the extracts, only concentration C4 (100 µl/ml) proved to be fungicide on both isolates in the same way as the synthetic fungicides used as reference; the other concentrations were fungistatic (Table 6).

Table 6. Fungicide or fungistatic activity of aqueous and organic extracts of *Thevetia peruviana* seeds

Extracts	Concentrations in (g/ml ou l/ml)	Activities
Aqueous	C1, C2 and C3	Fungistatic
	C4	Fungicide
Methanol	C1, C2 and C3	Fungistatic
	C4	Fungicide
Ethyl acetate	C1, C2 and C3	Fungistatic
	C4	Fungicide
Acetone	C1, C2 and C3	Fungistatic
	C4	Fungicide

- DISCUSSION**

Phytochemical screening revealed that aqueous and organic extracts of *T. peruviana* contain several families of bioactive compounds such as essential and saponifiable oils, coumarins, sterols, alkaloids, saponins, anthocyanins, steric glycosides and sugars. Plant

extracts of many plants with fungicide properties contain these major groups of bioactive phytochemicals as demonstrated by [9]; [27]; [28].

Spores germination was strongly inhibited by the different extracts with a total inhibition obtained with the highest concentration tested. This highly significant reduction in germination rate could be explained by the effect of secondary metabolites contained in the aqueous and organic extracts of *T. peruviana*. In addition, the present study showed the presence of sterols, saponins and essential oils which could have acted together or independently leading to an effective fungicide activity against the different isolates of *A. solani* as demonstrated by [29].

The different extracts tested significantly reduced the mycelia growth of *A. solani* compared to the control with an inhibition percentage of 100 % for the C4 concentration (100 L/mL) and (100 L/mL) in the same way as the synthetic fungicides used. This reduction was more effective with the acetone extract followed by the aqueous extract than with the methanol and ethyl acetate extracts. This would also be due to the presence of different secondary metabolites in these extracts. Indeed, [30] had previously reported that plants are abundant source of various bioactive compounds, many of which are secondary metabolites serving as signal chemicals and conferring resistance to many fungal plant pathogens. [18] previously reported that *Azadirachta indica* extracts had an effective inhibitory effect in the control of *Phakopsora pachyrhizi* responsible for Asian soybean rust. These results are in line with those of [31], who evaluated the effect of aqueous and organic extracts of *T. peruviana* on the *in vitro* development of *Phytophthora colocasiae* and found that they significantly reduced the growth of the fungi at the highest concentrations.

The percentages of inhibition of the plant extracts on the growth of the pathogen varied with increasing concentrations as well as the nature of the extraction solvents. At the highest concentration C4, all extracts showed total inhibition of *Alternaria solani* spore development and germination in the same way as the synthetic fungicides: Maneb and Dimethomorpe + Clorothalonil. This could be explained by the anti-root and antioxidant activity of the anthocyanins present in the said extracts as demonstrated by [32]. In these different tests, the inhibition was more pronounced with the acetone extract followed by the aqueous extract. Acetone showed a high inhibition percentage at the lowest

concentrations on both strains. The effectiveness of this extract on the inhibition of radial growth and spore germination of the two *A. solani* strains can be explained by the fact that it is a highly water miscible compound and by the polarity of this solvent compared to the other three as demonstrated by [33]. This is due to the fact that each compound acts differently on the fungi, i.e. one compound could have a very important action on the pathogen with the acetone extract. However, this result is contrary to that of [11] who in his work on the evaluation of the antifungal activities of *T. peruviana* against *Phytophthora colocasiae* showed that the reduction in growth was more pronounced with the ethyl acetate extracts. The same is true of the work of [10] who showed that it was the methanol extract that appeared to be more effective in inhibiting the mycelia growth of *Phytophthora infestans*.

With regard to MIC50 and MIC90 it was also the acetone extract that was more effective than the other extracts. This efficacy of the acetone extract was evidenced by the lowest MIC values obtained as demonstrated by [34-35] who argued that low MIC values obtained with an extract mark its proven efficacy. This result shows a more pronounced inhibition with the Mbal isolate compared to the Foun isolate. This would mean that the latter is less sensitive than the Mbal isolate and this difference is due to the membrane specificity of each isolate. The different antifungal tests carried out revealed that the C1 to C3 concentrations of the aqueous and organic extracts of *T. peruviana* were fungistatic compared to the highest C4 concentration (100 L/mL) which was fungicide. This result confirms that *T. peruviana* extracts would possess both fungistatic and fungicide properties and is in line with that of [36] who obtained fungistatic and fungicide activities with organic extracts (acetone, ethyl acetate, methanol and hexane) of *J. curcas* seeds against *Cercospora malayensis* causal agent of Cercosporiose and insect pests of okra.

CONCLUSION

The present study revealed the significant influence of the aqueous and organic extracts of *Thevetia peruviana*, on spore germination and mycelia development of *Alternaria solani*, and it was the acetone extract that was more effective followed by the aqueous extract compared to the other extracts (ethyl acetate and methanol). For both strains, the inhibition of both parameters was proportional to increasing concentrations of the different

extracts, with total inhibition obtained with the highest concentration (100 L/mL). At this concentration, all extracts were fungicide in the same way as the two synthetic fungicides used as reference. These extracts could therefore be used as a component of a more sustainable integrated control strategy against Alternariosis in tomato. This preliminary study provides a basis for future *in situ* (field) and *in vivo* (greenhouse) trials.

ACKNOWLEDGEMENTS

Our thanks to Professor MKOUNGA Pierre (MC) and his collaborators (Dr PAGNA MAWOUMA Issah Julio and Mr. LITCHANG Martial Lamer) of the laboratory of natural products and application of organic synthesis in the Department of Organic Chemistry of the University of Yaounde I who kindly put at our disposal the necessary equipment for the realization of a part of this work.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

- Anonymous. Annuaire de production: Rome, FAO. 2018;2.
- 2. Milet A. Isolation of microorganisms from the soil of arid regions and selection of isolates with antagonistic effect on the Alternaria blight agent. Thesis, University of the Mentouri Brothers Constantine 1, 2017;122.
- 3. Shankara N, Joep LJ, Marja G, Martin H, Barbara D. Tomato cultivation production,

processing and marketing. 2005;105.

4. Adegbola YP, Ahoyo Adjovi NR, Adekambi SA. Consumer Preferences for Fresh Tomatoes in Benin using a Conjoint Analysis. *J Int Food Agribus Mark* 2019;31:1-21. <https://doi.org/10.1080/08974438.2018.1469448>.

5. Anonymous. Production: Crops and livestock products. In: FAO. 2022 <https://www.fao.org/faostat/en/#data/QCL>, accessed on January 2023.

6. Koffi M.H, Yah M, Koffi BC, Amon MW. Effect of four substrate treatments on the appearance of tomato diseases [*Solanum Lycopersicum* L. Solanaceae)] in soilless cultivation, Ivory Coast. *European Scientific J.* 2021; 3-16.

7. Boutoumou H, Boumaza. Study of the activity of *Trichoderma* sp. Against Early Blight of tomatoes. Master's thesis, University of the Mentouri Constantine Brothers, 2016;78.

8. Blancard D. Early blight of tomato. *Ephytia Review*, INRAE, 2020.

9. Mboussi SB, Ambang Z, Ndogho A, Ngoh Dooh JP, Manga EFl. In vitro Antifungal Potential of Aqueous Seeds Extracts of *Azadirachta indica* and *Thevetia peruviana* against *Phytophthora megakarya* in Cameroon. *Jrn. Apl. L. Sc. Int.*, 2016;4(4): 1-12.

10. Dida LSL, Kuate T.N.W, Ngatsi Z.P, Heu A, Mboussi B.S, Kone N.A.N, Gbaporo G.C.F, Ambang Z. First report of different mating types of *Phytophthora infestans* isolates from potatoes in Cameroon and their control using *Thevetia peruviana* (Pers) K. Schum seeds extracts analyzed by GC-MS. *Indian Phytopathology.* 2024;1-9.

11. Essomé SC. Efficacy of aqueous and organic extracts of the seeds of *Thevetia peruviana* (Pers.) K. Schum in the fight against downy mildew of taro (*Colocasia esculenta* (L) Schott) caused by *Phytophthora colocasiae* Racid. Doctoral thesis Ph.D University of Yaoundé I, 2021;154.

12. Saxena VK, Jain SK. *Thevetia peruviana* kernel oil: a potential Bactericidal agent. *Fitoterapia*, 1990;61:348-349.

13. Tewtrakul S, Nakamura N, Hattori M, Fugiwara T, Suparita T. Flavone and flavonol glycosides from leaves of *Thevetia peruviana* and their HIV-1 reverse transcriptase and HIV-1 integrase inhibitory activities. *Cem. Pharm. Bull.* 2002;50:630-635.

14. Ambang Z, Ngoh Dooh JP, Djilé B. Effect of yellow laurel seeds (*Thevetia peruviana* (Pers) K. schum) on weevils (*Sitophilus zeamais* Motsch) stock pests. *Bios prod*, 2005;11:57-63.

15. Ngoh Dooh JP. Evaluation of the effect of Extracts of *Thevetia peruviana* (Pers.) K. Schum on three phytopathogenic fungi: *Phytophthora magakarya* Bras. And Grif., *Collectotrichum gloesporides* Penz f. sp. *Manihotis* HENN and *Mycosphaerella fijiensis* Morelet. Doctoral thesis Ph.D University of Yaoundé I, 2015;190.

16. Ambang Z, Ngoh Dooh JP, Essono G, et al (2010). Effect of *Thevetia peruviana*' Seeds Extract on in vitro Growth of Four Strains of *Phytophthora megakarya*. *Plant Omics* 2010;3:70-76.

17. Meena BR, Meena S, Chittora D, Sharma K. Antifungal efficacy of *Thevetia peruviana* leaf extract against *Alternaria solani* and characterization of novel inhibitory compounds by Gas Chromatography-Mass Spectrometry analysis. *Biochem Biophys Rep.* 2021;17;25:100914.
18. Ndogho PA, Asseng CC, Ndongo B, Makamte PED, Heu A, Mboussi SB, Ambang Z. Antifungal potential of Aqueous Extracts of Neem Seeds (*Azadiracta indica*) on the in-vitro development of Six Strains of *Phacopsora Pachyrhizi*, causal agent of Asian Rust of Soybean. *Journal of applied life science international*, 2017;15(4):1-9.
19. Nafisa S, Amna S, Javed I. Cultural, morphological, molecular comparison and pathogenicity of *Alternaria solani* causing early blight disease in tomato. *Mycopathology journal.* 2017;15(1): 7-11.
20. Bessadat N. Identification and characterization of *Alternaria* sp., responsible for the deterioration of market garden plants by enzymatic and molecular systems. LMD doctoral thesis. University of Oran es-Sena, 2014;213.
21. Edeoga HO, Okwu DE, Mbaebie BO. Phytochemical constituents of some Nigeria medicinal plants. Department of biological and chemical sciences, Michael Okpara University of agriculture. *African J. f Biotechnology.* 2005;4(7):685-688.
22. Sahuvinod K, Irlhaiya R, Shashi A, Gujar H. Phytochemical investigation and chromatographic evaluation of the ethanol extract of whole plant extract of *Dendrophthoe falcate* (L.F) Ettingsh. *International Journal of Pharmaceutical sciences and Research* 2010;1(1): 39-45.
23. Dohou N, Yamani K, Badoc A, Douira A. Antifungal activity of extracts of *Thymelaea lytroides* on three pathogenic fungi of Rice. *Bull. Oc. Pharm. Bordeaux.* 2004;143:31-38.
24. Opara EU, Wokocha RC. Efficacy of Some Plant extracts on the in-vitro and in-vivo control of *Xanthomonas campestris* Pv. *Vesicatoria.* *Agric. J.* 2008;3(3):163-170.
25. Widmer T, Laurent N. Plant extracts containing caffeie acid rosmarinic acid inhibitor zoospore germination of *Phytophthora* spp pathogenic to *Theobroma cacao*. *E.Jrn. Pit Phytol.* 2006;115:377-388.
26. Kishore N, Mishra AK, Chansouria JPN. Fungitoxicity of essential oils against Dermatophytes. 1993;5(6):211-215.
27. Dida Lontsi SL, Ambang Z, Djieto CL, Chedjou WBC, Heu A, Ndogho PA. Effect of aqueous extracts of *Thevetia peruviana* K. seeds on the control of late blight and pest insects of *Solanum tuberosum* L. in Cameroon. *Greener J. of Agric. Science* 2019;9(1):14-22.
28. Sanjukta R, Himanshu AP, Ahamedabad. Department of Botany, Bioinformatics and Management of the Impacts of Climate Change. *IJRAR International Journal of Research and Analytical Reviews.* 2019;935.
29. Mohammed Z. Phytochemical studies and biological activity of some plants medicines

of the northern and southwestern region of Algeria, doctoral thesis in biology. University of Aboubaker Belkaid Algeria. 2013;170.

30. Ramirez GXS, Jimenez GSN, Vicente B C, Lourdes GC. Plant metabolites in plant defense against pathogens in plant diseases; Current Threats and Management Trends 2019;1-19.

31. Essomé SC, Ngoh Dooh JP, Heu Alain, Ndogho PA, Ngatsi ZP, Chewachong G, Ambang Zachee. Evaluation of antifungal activities of *Thevetia peruviana* seed extracts against *Phytophthora colocasiae* (Oomycetes) causal agent of taro late blight (*Colocasia esculenta* (L.) Schott) J. of Applied Biosciences 2020;151:15584-15597.

32. Jin R, Bao H, Bai Y, Li X. Theoretical study on the antioxidant activity of alizarin, purpurin, and pseudopurpurin. In Interdisciplinary research and applications in bioinformatics, computational biology, and environmental sciences (IGI Global), 2011;130-140.

33. Bougandoura N, Bendimerad N. Antifungal Effect of Aqueous and Methanolic Extracts of *Satureja calamintha* ssp. (*Nepeta*) briq. Revue des Bio Ressources 2012;2:1-7.

34. Serghat S, Mouria A, Touhami AO, Douira A. In vivo effect of some fungicides on the development of *Pyricularia grisea* and *Helminthosporium oryzae*. Phytopathol. Mediterr. 2004;41:235-246.

35. Serghat S, Mouria A, Touhami AO, Douira A. Natural protection of plants in tropical zones. CTA. Agricultural. 2004;95 -99.

36. Koné SAN. Evaluation of the bioeffectiveness of *Jatropha curcas* seed extracts against *Cercospora malayensis*, causal agent of Cercosporiosis and insect pests of okra. Ph.D Doctoral Thesis University of Yaoundé I (Cameroon), 2019;115.

- Song W, Ma X, Tan H, Zhou J. Abscisic acid enhances resistance to *Alternaria solani* in tomato seedlings. Plant Physiology and Biochemistry. 2011 Jul 1;49 (7):693-700.
- Kumar S, Singh R, Kashyap PL, Srivastava AK. Rapid detection and quantification of *Alternaria solani* in tomato. Scientia horticultrae. 2013 Feb 28;151:184-9.
- Dole P. D, Gaikwad S. D, Digvijay C. Optimizing Tomato Health: Holistic Approach to Early Blight Disease Management. J. Adv. Biol. Biotechnol. [Internet]. 2024 May 17 [cited 2024 Jun. 13];27(6):492-7. Available from: <https://journaljabb.com/index.php/JABB/article/view/909>
- Dobariya KG, Pandya JR, Pandya RP, Joshi SH. Biological Control of Early Blight of Tomato (*Solanum lycopersicum* L.) by the Use of *Pseudomonas fluorescens* (Flugge and Migula) under Field Condition. Curr. J. Appl. Sci. Technol. [Internet]. 2023 Jun. 2 [cited 2024 Jun. 13];42(13):34-41. Available from: <https://journalcjast.com/index.php/CJAST/article/view/4115>