

**Effect of integrated nutrient management in
China aster cv. ArkaPoornima under mid hills of
Himalayas**

ABSTRACT

The present investigation conducted to investigate the effect of integrated nutrient management on vegetative parameters of China aster cv. ArkaPoomima under the mid hills of Himachal Pradesh. The experiment was laid out in simple randomized design (RBD) with eighteen treatments. The experiment was conducted at the Kakhali research farm of Dr. Khem Singh Gill Akal College of Agriculture, Eternal University, Baru Sahib, Himachal Pradesh during *kharif* season of 2023. For integrated nutrient management eighteen treatments were constituted with three levels of recommend dose of inorganic fertilizers (50, 75 and 100 %) and three biofertilizers viz., *Azotobacter*, PSB and KSB along with control. The observations for the vegetative parameters i.e., plant height (cm) at maturity, numbers of leaves, number of primary branches, number of secondary branches, leaf area per plant, leaf area index and plant spread were recorded and analyzed. The plant height (91.40 cm), No. of leaves (356.13), No. of primary branches (12.40) and secondary branches (33.70) were found maximum in treatment T₁₂ (RDF@75% + *Azotobacter* + PSB + KSB), while the lowest values were observed in T₀ (control). The highest leaf area (27.49 cm²) was found in T₈ (RDF@75% + *Azotobacter*), while the leaf area index (12.02) and plant spread (16.57 cm) were found in treatment T₁₁ (RDF@75% + *Azotobacter* + PSB). The lowest values for leaf area (17.77 cm²), leaf area index (5.48) and plant spread (11.82 cm) were found in treatments T₁₄ (RDF@100% + *Azotobacter*), T₁₆ (RDF@100% + KSB) and T₁₀ (RDF@75% + KSB), respectively. Therefore, it can be concluded that, application of T₁₂ (RDF@75% + *Azotobacter* + PSB + KSB) is best for the getting higher production in China aster followed by T₁₁ (RDF@75% + *Azotobacter* + PSB). This study will reduce the use of inorganic fertilizer without polluting the soil and environment.

Comment [H1]: was

Comment [H2]: Justify the rational of the study in a sentence before the methodology.

Comment [H3]: in

Comment [H4]: delete

Keywords: China aster, Integrated nutrient management, *Azotobacter*, Phosphorous solubilizing bacteria, Potassium solubilizing bacteria

1. INTRODUCTION

The China aster (*Callistephus chinensis* L) belongs to the Asteraceae family and is indigenous to China. It is a prominent annual flower crop cultivated worldwide for a variety of uses, including cut flowers, loose flowers, pot plants, and landscape bedding. The flower has grown in popularity in the floral business because to its small size, diverse colour palette, extended vase life, and transportability [1]. In India, it is mostly grown in Karnataka, Maharashtra, Andhra Pradesh, Tamil Nadu, and West Bengal. China aster cultivated in Indian fields could be impacted from toxic nutrient levels. The right fertilizer combinations are essential for growing robust plants with more shoots and leaves, improving flower quality, and lengthening the flowering season [2,3]. There hasn't been much research done on the effects of organic and inorganic fertilizers on China aster growth and production, though. Combining biofertilizers (*Azotobacter*, phosphorous solubilizing bacteria and potassium solubilizing bacteria) with balanced inorganic fertilizers and organic manures is crucial for achieving higher yields. Increased interest in organic farming can result from the use of organic materials such as green manures, which can enhance soil structure, nutrient exchange, and general soil health [4]. According to Shylaja et al. [5], FYM (farmyard manure) helps regulate soil pH and increases accessible NPK levels. Biofertilizers increase the quantity and quality of flowers while using less inorganic fertilizer. They improve soil qualities and provide a home for advantageous microbes, which promotes long-term soil fertility [6]. PSB raises the availability of phosphorus, increasing yield. Therefore, the prime objective of the present investigation is to investigate the effect of integrated nutrient management in China aster cv. ArkaPoomima using inorganic fertilizers and biofertilizers under mid hills of Himachal Pradesh.

Comment [H5]: of

Comment [H6]: the introduction is too brief, try and add other literature

2. MATERIAL AND METHODS

The present study was carried out in Kakhali research farm of Dr. Khem Singh Gill Akal College of Agriculture, Eternal University, Baru Sahib, Sirmour, Himachal Pradesh (India) during the *Kharif* season of 2023. The study geographical area lies on latitudes 30.7537° N and longitudes 77.2965° E of mid hills of Himachal Pradesh (India) in an altitude of 1067 m from above mean sea level. The research trail includes 18 treatments, combing different levels of recommended dose of fertilizers with biofertilizers laid out in randomized block design. The effect of different treatments of integrated nutrient management i.e., T₀ (Control), T₁ (RDF@50%), T₂ (RDF@50% + *Azotobacter*), T₃ (RDF@50% +PSB), T₄ (RDF@50% +KSB),

Comment [H7]: The coordinates did not cover the study area. For example latitude ?N to latitude ?N and so on

Comment [H8]: What informed your choice of using 18 treatments. Why not 10, 20 Or 30. Justify the choice of selection

T₅ (RDF@50% + Azotobacter+PSB), T₆ (RDF@50% + Azotobacter+PSB+KSB), T₇ (RDF@75%), T₈ (RDF@75% + Azotobacter), T₉ (RDF@75% +PSB), T₁₀ (RDF@75% +KSB), T₁₁ (RDF@75% + Azotobacter+PSB), T₁₂ (RDF@75% + Azotobacter + PSB + KSB), T₁₃ (RDF@100%), T₁₄ (RDF@100% + Azotobacter), T₁₅ (RDF@100% +PSB), T₁₆ (RDF@100% +KSB), T₁₇ (RDF@100% + Azotobacter+PSB) and T₁₈ (RDF@100% + Azotobacter + PSB + KSB) on 'ArkaPoornima' of China Aster (Table 1). The seeds of China aster cv. ArkaPoornima were sown in the nursery, and one month old seedlings of China aster were treated with the liquid consortium of *Azotobacter*, PSB (Phosphorous solubilizing bacteria) and KSB (Potassium solubilizing bacteria) separately for few minutes. After that the treated seedlings were planted in the experimental field in a spacing of 30 cm plant to plant and 30 cm row to row. The observations for the traits i.e. plant height (cm) at maturity, numbers of leaves, number of primary branches, number of secondary branches, leaf area per plant, leaf area index and plant spread were measured and recorded as per standard procedure. The plant parameters were recorded, by selecting the five random plants from each replication of a treatment and significant variations were recorded from the plant. The plant were undergone from the various inter culture operations and plant protection procedure required for the better plant production during the course of investigation. Further, the analysis of variance (ANOVA) was done through OPSTAT software [7].

Comment [H9]: grammar

Comment [H10]: state how you use the statistical technique mentioned to analyze the data generated in this study. It is ambiguous

3. RESULTS AND DISCUSSION

The observation recording for plant height (cm) was significantly affected by the treatments of various integrated nutrient management. The highest plant height (91.40 cm) of China Aster cv. ArkaPoornima was found in T₁₂ (RDF@75% + Azotobacter + PSB + KSB), which was closely followed by T₁₁ (RDF@75% + Azotobacter + PSB) and T₁₆ (RDF@100% + KSB) recording 90.07 cm and 87.00 cm, respectively. Whereas, the lowest plant height (74.13 cm) was observed in T₀ (Control) (Table 1). The increased plant height can be associated with PSB's critical contribution in the solubilization of insoluble P by the formation of organic acid. The appropriate supply of P appears to have encouraged root development, which leads to rapid vegetative growth of plants. [8] noticed a similar finding in tuberose. The data recorded for number of leaves per plant was significantly affected by the different treatments of integrated nutrient management. The maximum number of leaves per plant (356.13) of was found in T₁₂ (RDF@75% + Azotobacter + PSB + KSB), which was followed by T₁₁ (RDF@75% + Azotobacter + PSB) and T₁₆ (RDF @100% + KSB) recording 354.27 and 345.53 respectively (Table 1). Whereas, the minimum number of leaves per plant (276.07) was found in T₀ (Control). The increment in plant growth is can be due to reason that Biofertilizers produce a number of growth-promoting hormones (auxins, cytokinins, and gibberellins, etc.) while also improving the accessibility of nitrogen, phosphate, and potash to plants, leading to improved plant development. Several authors have observed similar results of increased plant height at harvest because of the combination application of biofertilizers with lower doses of NPK i.e. [9,10] in China Aster, [11] in annual chrysanthemum and [12] in chrysanthemum.

Comment [H11]: Recast

The observation regarding the number of primary branches per plant was showed significant variation for different treatments of integrated nutrient management. The highest number of primary branches per plant (12.40) of was recorded in T₁₂ (RDF@75% + Azotobacter + PSB + KSB), which was followed by T₁₁ (RDF@75% + Azotobacter + PSB) recorded as 11.77 number of primary branches per plant. While, the lowest number of primary branches per plant (9.13) was observed in T₀ (Control). The highest number of secondary branches per plant (33.70) of was found in T₁₂ (RDF@75% + Azotobacter + PSB + KSB), which was followed by T₁₁ (RDF@75% + Azotobacter + PSB) and T₁₆ (RDF @100% + KSB) recording 31.70 and 29.07, respectively. Whereas, the lowest number of secondary branches per plant (22.13) was observed in T₀ (Control) (Table 1). The increase in branches could be attributed to the formation of nitrogenous compounds such as proteins, amino acids, nucleic acids and other enzymes and coenzymes that were responsible for cell division and enlargement, as well as the role of phosphorous in structural components such as phospholipid and the translocation of food components. Additionally, Azotobacter's presence in the nitrogen fixation process and the release of growth-promoting chemicals such as gibberellins and IAA may contribute to an increase in the number of primary branches per plant. There have been reports of a similar result, with an increase in the number of primary branches following the immunization of African marigold with Azotobacter and PSB (Phosphorous solubilizing bacteria) by several authors [13,14] in annual chrysanthemum.

Comment [H12]: ???

Comment [H13]: Poor grammar. showed

Comment [H14]: Compare and contrast this finding/claim with other results in the literature

The perusal of data showed that, the maximum leaf area per plant (27.49 cm²) was observed in T₈ (RDF@75% + Azotobacter), which was followed by T₉ (RDF@75% + PSB) recording 27.13 cm². Whereas, the minimum leaf area per plant (17.77 cm²) was observed in treatment T₁₄ (RDF@100% + Azotobacter) (Table 1). These growth-promoting compounds may have resulted from increased cell division and elongation, which improved leaf explanation by increasing leaf length and width, thereby increasing leaf area. These results closely align also the conclusions of [15] and [11] in chrysanthemum and [16] in petunia [17] in China aster.

The highest leaf area index (12.02) of was found in T₁₁ (RDF@75% + Azotobacter + PSB), which was followed by T₁₅ (RDF@100% + PSB) and T₁₀ (RDF@75% + KSB) recording 10.40 and 10.15, respectively (Table1). Whereas, the lowest leaf area index (5.48) was found in T₁₆ (RDF@100% + KSB). Leaf area index is dependent to total leaf area and numbers of leaves, therefore treatment having the larger area and number of leaves will also have the higher leaf area index. These results closely align also the conclusions of [15] and [11] in chrysanthemum and [16] in petunia [17] in China aster.

The highest plant spread (16.57 cm) of was found in treatment T₁₁ (RDF@75% + Azotobacter + PSB), which was followed by T₁₄ (RDF@100% + Azotobacter) recording 16.35 cm. whereas, the lowest plant spread (11.82 cm) was observed in T₁₀ (RDF@75% + KSB) (Table 1). The increased plant spread could be related to the creation of colonies of the biofertilizers (Azotobacter and PSB) inoculates, as well as their growth-promoting actions, such as the generation of plant growth-promoting substances. A further potential reason for greater plant spread after biofertilizers inoculation is increased cell metabolism, which is induced by increased enzyme activity, chlorophyll content, and photosynthetic activities. The similar observations were also discovered by [18] in Annual chrysanthemum, [19] in petunia, [20] in marigold and [21] in bird of paradise. The use of NPK together with the inoculation of bio-fertilizer proved beneficial, presumably because it fixed atmospheric nitrogen in the soil and made phosphorus freely available to plants. The role and stimulating effects of non-symbiotic nitrogen-fixing bacteria (Azotobacter) in connection with N-fertilization have been thoroughly researched these nutrients are essential for increasing soil microbial activity. Similar results were reported [5, 22]

Comment [H15]: delete

Comment [H16]: why

Comment [H17]: is there studies related to this work. Please state their studies

Comment [H18]: Recast

Comment [H19]: delete

Table 1. Effect of inorganic and biofertilizers on vegetative plant parameters of China aster

4. CONCLUSION

From the findings of the current experiment, It can be inferred that, T₁₂ (RDF@75% + Azotobacter + PSB + KSB) was the best treatment for the plant height (cm), No. of leaves, No. of primary and secondary branches. Whereas, T₈ (RDF@75% + Azotobacter) is superior for leaf area (cm²) and T₁₁ (RDF@75% + Azotobacter + PSB) were best for the leaf area index and plant spread (cm). Therefore, it can be concluded that from the findings of the experiment, that plant treatment with T₁₂ (RDF@75% + Azotobacter + PSB + KSB) and T₁₁ (RDF@75% + Azotobacter + PSB) are best for the getting higher production in China aster under mid hills of Himalayas. Without causing pollution to the environment or soil, this study will decrease the need of inorganic fertilizers and directly increases the farmer's income.

Comment [H20]: The conclusion did not summarize all the key finding. Recommendation and area for further study should be highlighted

DATA AVAILABILITY STATEMENT

Data will be available in request

REFERENCES

1. Santhosh T, Anupama V, Sreelatha U, Minimol JS, Sankar M. Evaluation of China aster (*Callistephus chinensis* (L.) Nees.) genotypes in tropical plains of Kerala, J. Trop. Agric. 2020; 58(2).

Treatments	Plant Height at maturity (cm)	No. of Leaves Plant per plant	No. of Primary Branches Plant per plant	No. of Secondary Branches Plant per plant	Leaf Area Plant ⁻¹ (cm ²)	Leaf Area Index	Plant Spread (cm)
T ₀	74.13	276.07	9.13	22.13	23.38	8.53	13.31
T ₁	84.17	307.73	9.27	23.2	24.75	8.82	12.33
T ₂	86.47	312.53	9.57	24.43	25.53	8.93	12.83
T ₃	76	297.27	9.97	26.4	23.34	9.32	13.09
T ₄	78.8	304.67	11.27	23.5	21.4	8.4	13
T ₅	79.73	315.67	10.73	25	20.55	7.74	12.18
T ₆	83.73	284.9	10.53	23.13	21.65	8.03	13.92
T ₇	85	314.6	11.23	27	25.69	9.73	12.67
T ₈	83.27	313.6	10.67	23.23	27.49	9.7	13.45
T ₉	85.8	305.87	11.33	28.2	27.13	10.05	13.19
T ₁₀	76.93	322.8	10.7	26.5	24.55	10.15	11.82
T ₁₁	90.07	354.27	11.77	31.7	21.51	12.02	16.57
T ₁₂	91.4	356.13	12.4	33.7	19.81	7.11	12.28
T ₁₃	86	306.53	10.63	27.2	19.48	9.11	12.8
T ₁₄	86.6	296.6	11.37	25	17.77	7.67	16.35
T ₁₅	86.27	294.6	10.43	24.77	20.63	10.4	15.77
T ₁₆	87	345.53	11.47	29.07	22.47	5.48	13.75
T ₁₇	81.47	315.13	10.37	25.1	24.51	9.86	14.11
T ₁₈	85.73	286.67	11.4	25.33	24.13	9.71	12.94
C.D. at 5 %	9.5	5.23	1.61	3.73	1.72	0.76	1.79

2. Sultana S, Khan FN, Haque MA, Akhter S, Noor S. Effect of NPK on growth and flowering in tuberose. *J. Ornament. Hort.* 2006; 4(2): 111-113.

Comment [H21]: journal

3. Zhang WenJun ZW, Lu JianWei LJ, Jiang ZhiPing JZ, Wang YaYi WY. Effects of nitrogen, phosphorus and potassium fertilizer on the growth, blossom and N, P, K uptake in *Calendula officinalis* L. 2010; 28(4): 491-496.

4. Mitra M. Response of tuberose to integrated nutrient management in Int. Conf. Biodivers. Livelihood Clim. Change Himalaya Dep. Bot. Tribhuvan Univ. India. 2010.

5. Syamal MM, Dixit SK, Kumar S. Effect of bio-fertilizers on growth and yield in marigold. *J. Ornament. Hort.* 2006; 9:304-305.

Comment [H22]: Full meaning

6. Harris RF, Chesters G, Allen ON. Dynamics of soil aggregation. *Adv. Agron.* 1966; 18:107-169.

Comment [H23]: Antiquated

7. Sheoran OP, Tonk DS, Kaushik LS, Hasija RC, Pannu RS. Statistical software package for agricultural research workers. Recent advances in information theory, statistics and computer applications. Hooda DS, Hasija RC (Eds). Department of Mathematics Statistics, Chaudhary Charan Singh Haryana Agricultural University, Hisa, Haryana. 1998; 139-43.

8. Srivastava R, Preetham S, Chand S. Effect of organic manures and biofertilizers on vegetative, floral and post-harvest attributes in tuberose (*Polianthes tuberosa*) var. Shringar. *Asian J. Biol. Sci.* 2014; 3(1).

Comment [H24]: Full meaning

9. Chitra R, Patil V. Integrated nutrient management studies in china aster (*Callistephus chinensis* (L.) Ness). *Karnataka J. Agric. Sci.* 2007; 20(3):689-690.

10. Yaseen T, Khan F, Gani I, Abass M. Response of globe amaranth (*Gomphrenaglobosa* L.) to pinching and biofertilizer application. *Int. J. Hortic and Food Sci.* 2021; 3(1):1-5.

11. Airadevi A. Integrated nutrient management studies in garland chrysanthemum (*Chrysanthemum coronarium* L.). *Bioinfolet-A Quarterly J. Life Sci.* 2012;9(4):430-434.

12. Verma S, Angadi S, Patil V, Mokashi A, Mathad J, Mummigatti U. Growth, yield and quality of chrysanthemum (*Chrysanthemum morifolium* Ramat.) cv. Raja as influenced by integrated nutrient management. *Karnataka J. Agric. Sci.* 2012; 24(5).

13. Gupta N, Sadavarte K, Mahorkar V, Jadhao B, Dorak S. Effect of graded levels of nitrogen and bioinoculants on growth and yield of marigold (*Tagetes erecta*). 1999.

14. Panchal R, Parekh N, Parmar A, Patel H. Effect of biofertilizers and nitrogenous fertilizer on growth, flowering and yield of annual white chrysanthemum (*Chrysanthemum coronarium* L.) under middle Gujarat agroclimatic conditions. 2010.

Comment [H25]: Volume, number and pages

15. Verma S. Integrated nutrient management studies in chrysanthemum *Chrysanthemum morifolium* Ramat. cv. Raja. M. Sc. Thesis Unpub. Submitted to UAS, Dharwad. 2010.

16. Hoda E, Mona S. Effect of bio and chemical fertilizers on growth and flowering of *Petunia hybrida* plants. 2014.

17. Mahadik M, Dalal S, Patil STU. Effect of bioinoculants with graded doses of NPK on growth of chrysanthemum and availability of micronutrients in soil. *IJCS.* 2017; 5(5):892-895.

Comment [H26]: Write in full

18. Meshram N, Badge S, Bhongle S, Khiratkar S. Effect of bio-inoculants with graded doses of NPK on flowering, yield attributes and economics of annual chrysanthemum. 2008.
19. Moghadam Z, SHOOR M. Effects of vermi-compost and two bacterial bio-fertilizers on some quality parameters of Petunia. Not. Sci.Biol. 2013;5(2), 226-231.
20. Kumar S, Singh J, Mohan B, Nathiram N, Rajbeer R. Influence of integrated nutrient management on growth, flowering and yield parameters of marigold (*Tagetes erecta* L.) cv. PUSA BASANTI GAINDA. 2013.
21. Yathindra H, Krishna M, Rajesh A, Harshavardhan M. Effect of Integrated Nutrient Management on Growth Parameters of Bird of Paradise (*Strelitzia reginae* (L.)). The Bioscan. 2016; 11(1):565-568.
22. Rolaniya MK, Khandelwal S, Koodi S, Sepat S, Choudhary A. Effect of NPK, biofertilizers and plant spacings on growth and yield of african marigold (*Tagetes erecta* Linn.). Chem. Rev. Lett. 2017; 6(21):54-58.

Comment [H27]: Volume, number and pages