

Original Research Article

Performance of Different Hybrids of Bitter Gourd (*Momordica charantia* L.) in terms of Growth, Yield and Quality in Prayagraj Agro-Climatic Condition

ABSTRACT

Bitter gourd botanically known as *Momordica charantia* (L.) is an important commercial cucurbitaceous vegetable belonging to the family Cucurbitaceae, with a diploid chromosome number, $2n=22$. It is variously known as balsam pear, bitter melon, bitter gourd, and African bitter gourd. Therefore, the present investigation was carried out at the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, Uttar Pradesh during the *Zaid* season 2023 with a view to check performance of different genotypes of bitter gourd under Prayagraj agro climatic conditions. The experiment was laid in Randomized block design with 12 hybrids with 3 replications. Genotypes comprised of H₁ (AVT-II 2020/BIGHYB-1); H₂ (AVT-II 2020/BIGHYB-2); H₃ (AVT-II 2020/BIGHYB-3); H₄ (AVT-II 2020/BIGHYB-4); H₅ (AVT-II 2020/BIGHYB-5); H₆ (AVT-II 2020/BIGHYB-6); H₇ (AVT-II 2020/BIGHYB-7); H₈ (AVT-II 2020/BIGHYB-8); H₉ (AVT-II 2020/BIGHYB-9); H₁₀ (Shriram Samridhi); H₁₁ (Alpine F₁ Hybrid) and H₁₂ (F₁ Hybrid Aman). From the present investigation, it is concluded that among twelve hybrids of Bitter gourd, hybrid AVT-II 2020/BIGHYB-6 performed best in terms of growth & yield. The vine length was recorded 3.45 m, fruit yield was 122.48 q/ha. In the economic analysis of the hybrid AVT-II 2020/BIGHYB-6.

Keywords: *Momordica charantia*, Hybrids, Growth, Fruit Yield and Quality.

Introduction

Bitter gourd botanically known as *Momordica charantia* (L.) is an important commercial cucurbitaceous vegetable belonging to the family Cucurbitaceae, with a diploid chromosome number, $2n=22$. It is variously known as balsam pear, bitter melon, bitter gourd, and African bitter gourd (Heiser, 1979). Bitter gourd is grown in different varieties in different countries. In India the main varieties are India long green, India

long white and Hybrid India baby whereas Japan is famous for Japan Green Spindle, China is for green lovers and Hong Kong for its Hong Kong Green. In Bangladesh mainly two varieties are grown i.e., Uchee (small) and Korolla (long) (Alam *et al.*, 2015). The bitter gourd plant is perennial with climbing and flowering vine grows up to 5 m and fruits are elongated with ridges on surface (Lee *et al.*, 2009). The young fruit is emerald green

Commented [SC1]: Put comma

Commented [SC5]: Put comma

Commented [SC6]: Put comma

Commented [SC7]: Full stop

Commented [SC8]: In contrast,

Commented [SC2]: Put comma

Commented [SC9]: Put comma

Commented [SC10]: Put comma

Commented [SC3]: No need to write word variously

Commented [SC11]: Put comma

Commented [SC12]: Put comma

Commented [SC4]: Put comma

Commented [SC13]: the

and later changes to orange-yellow when it is ripened (Kandangath *et al.*, 2015). The shape and size of bitter gourds vary due to varieties, climatic factors, regions but in general bitter gourds are 1.0 to 9.8 inch long and 1.0 to 5.9 inch wide with round, oval, oblong and club in shape and colour varies from dark green to white. In India the length of bitter gourd ranges from 2.4 to 3.9 inch and dark green in colour with ridges on its surface (Kumar *et al.*, 2016). The bitter gourd fruit gets mature after 45-80 days and harvesting is carried out after 60 days and continues up to 150 days from the planting (Islam *et al.*, 2011). The shelf life of fresh bitter gourd is only 4 days at natural conditions, and it can be stored up to 3-4 weeks in cold storage (0 to 7 °C) (Wang *et al.*, 2007). The bitter gourd is highly nutritious due to presence higher amount of protein, ascorbic acid, calcium, iron, and phosphorus (Dandawate *et al.*, 2016), also important source of glucosides, carbohydrate, charantin, steroidal, saponin, momordium, vitamins, protein and minerals. The protein content in bitter gourd is fractioned into albumin (49.3 per cent), globulin (29.3 per cent) and glutelin (3.1 per cent) (Horaxet *al.*, 2010). The seeds of the bitter gourd contain 35 to 40 per cent of oil and fatty acid i.e., 3.33 per cent MUFA (monosaturated fatty acid) and 36.71 per cent SFA (saturated fatty acids) (Liu *et al.*, 2010). Bitter gourd contains high amounts of Vitamin A, Vitamin C and vitamins B1, B2, B3 and B12 (Joseph

and Jini, 2013). According to NHB, 2022, area under bitter gourd production in India accounts to 96.85 million ha with production of 1332.55 million tonnes in year 2021-22. Madhya Pradesh ranks first in area and production of bitter gourd in year 2021-22 followed by Chhattisgarh and Tamil Nadu. The production of bitter gourd in Uttar Pradesh is 89.73 million tonnes for year 2021-22. It has many uses in ayurvedic medicines. Study was undertaken to estimate the performance of different hybrid varieties of Bitter gourd (*Momordica charantia*) under Prayagraj Agro-climatic conditions. The varieties were evaluated based on growth, yield and quality of varieties, shelf life and economics of different Bitter gourd hybrid varieties. Yield is a complex character and their interactions. For any effective selection programme, it would be desirable to consider the relative magnitude of association of various characters yield.

MATERIALS AND METHODS

The present investigation entitled was done to understand the plant growth, fruit yield and quality of fruit of different F₁ hybrids of bitter gourd. The investigation was carried out at Horticultural Research Farm (HRF), Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj during the winter season of 2022-23. The experiment was laid in Randomized block design with 12 Hybrids and 3 replications. Hybrids

Commented [SC14]: no need to write sentence it is

Commented [SC32]: For

Commented [SC15]: and

Commented [SC33]: No need to write this word

Commented [SC16]: Inches

Commented [SC17]: Inches

Commented [SC34]: No need to write this word

Commented [SC18]: Put comma

Commented [SC19]: Put comma

Commented [SC20]: Inches

Commented [SC35]: No need to write this word

Commented [SC21]: No need to write it

Commented [SC22]: matures

Commented [SC23]: put comma

Commented [SC24]: Put comma

Commented [SC25]: No need to write it

Commented [SC26]: After

Commented [SC27]: under

Commented [SC36]: Put comma

Commented [SC28]: for

Commented [SC29]: Put comma

Commented [SC30]: Put comma

Commented [SC37]: Out in a

Commented [SC31]: Put comma

comprised of H₁ (2020/BIGHYB-1), H₂ (2020/BIGHYB-2), H₃ (2020/BIGHYB-3), H₄ (2020/BIGHYB-4), H₅ (2020/BIGHYB-5), H₆ (2020/BIGHYB-6), H₇ (2020/BIGHYB-7), H₈ (2020/BIGHYB-8) H₉ (2020/BIGHYB-9), H₁₀ (Shriram Samridhi), H₁₁ (Alpine F₁ Hybrid) and H₁₂ (F₁ Hybrid Aman). Observations were recorded at different stages of growth for parameters like vine length, days to flower emergence, fruit length, fruit diameter and yield per vine and quality parameters like TSS and vitamin C content. The data were statistically analysed by the method suggested by Fisher and Yates, 1936. The experimental site is levelled land with sandy loam soil of uniform fertility status with low clay and high sand percentage.

Soil samples were collected randomly from depth of 0-30 cm and the soil was analysed for pH found to be slight neutral (6.9), organic carbon was 0.36%, available nitrogen was 212.56 kg ha⁻¹, available phosphorus was 14.59 kg ha⁻¹, and available potassium was 225.10 kg ha⁻¹. The preparation of the experimental field involved several steps to ensure optimal conditions for cultivation. Initially, a Tractor drawn disc plough was used to plough the field. Following this primary ploughing, two cross harrowing sessions were conducted, and the field was then planked. To achieve a uniform surface, a leveller was employed to thoroughly level the field before proceeding with the experimental

layout. This sequence of activities was undertaken to create an environment conducive to the study's objectives and to promote consistent and reliable results. To maintain a weed-free field, regular and shallow cultivation was performed frequently. This process aimed to eliminate weeds, enhance soil aeration, and support healthy root development.

Furthermore, two to three hoeing sessions and earthing up were conducted to meticulously control weed growth and maintain the crop's weed-free status. Around FYM 40 t/ha as basal was applied to field and 35 kg of N/ha at 30 days after sowing. NPK are required for hybrid bitter gourd is 150:75:75 kg NPK/ha; accordingly urea, DAP, MOP was applied in field. Light irrigation was provided at critical stages of crop growth, such as just after transplanting, pre flowering, fruit formation.

RESULTS AND DISCUSSION

Growth Parameters

The data pertaining to Vine length and number of branches per vine significantly varied among different Hybrids. The hybrid AVT-II 2020/BIGHYB-6 had the longest vines overall, measuring 3.45m, while AVT-II 2020/BIGHYB-9 had the second longest, measuring 3.36 m. In AVT-II 2020/BIGHYB-8, shortest vine length of 2.51 m was noted. Hybrid AVT-II 2020/BIGHYB-6 of bitter gourd likely exhibits a longer vine length compared to other hybrids, due to its specific

Commented [SC43]: A leveller was employed to thoroughly level the field before proceeding with the experimental layout to achieve a uniform surface

Commented [SC44]: earthing-up

Commented [SC45]: put comma

Commented [SC46]: put comma

Commented [SC47]: was applied

Commented [SC48]: no need to write this word

Commented [SC38]: a

Commented [SC49]: crop growth stages

Commented [SC39]: Comma

Commented [SC50]: pre-flowering, and

Commented [SC40]: and

Commented [SC41]: Slightly

Commented [SC51]: the sentence written unclear

Commented [SC52]: no need to write this word

Commented [SC42]: Cross- harrowing

Commented [SC53]: than

genetic makeup and environmental interactions. Genetic factors may include traits favouring vine elongation, such as genes related to internode elongation, cell expansion, or hormone regulation. These genetic traits could contribute to increased cell division and elongation, resulting in longer vines.

Moreover, AVT-II 2020/BIGHYB-6 might possess alleles promoting vine growth under varying environmental conditions, ensuring consistent elongation throughout the growing season. Environmental factors like temperature, light intensity, and soil fertility can also influence vine length by affecting plant hormone levels and physiological processes. Therefore, the combination of favourable genetic attributes and environmental conditions likely contributes to the longer vine length observed in hybrid AVT-II 2020/BIGHYB-6, potentially leading to increased yield and productivity in bitter gourd cultivation. Research on bitter gourd (Kumari *et al.*, 2019; Kumar and Topno, 2022) reported similar results.

Earliness parameters

With 46.20 days, the hybrid AVT-II 2020/BIGHYB-1 was earlier most for male flowering overall; AVT-II 2020/BIGHYB-5 and Shriram Samridhi came second most early in number of days to first male flowering, with 51.80 days. The maximum days to first male flowering (53.67 days) was observed in AVT-II

2020/BIGHYB-8. With 54.67 days, the hybrid AVT-II 2020/BIGHYB-5 was the earlier for fruit harvest overall; AVT-II 2020/BIGHYB-7 came second most early in the number of days to the first fruit harvest, with 57.00 days. The maximum days to first female flower appearance

(53.83 days) was observed in AVT-II 2020/BIGHYB-8. With 54.67 days, the hybrid AVT-II 2020/BIGHYB-5 was earlier most for fruit harvest overall; AVT-II 2020/BIGHYB-7 came second most early in number of days to first fruit harvest, with 57.00 days. The maximum days to first fruit harvest (61.67 days) was observed in AVT-II 2020/BIGHYB-3.

Hybrid AVT-II 2020/BIGHYB-5 of bitter gourd likely exhibits early flowering and maturity compared to other hybrids due to a combination of genetic attributes and environmental influences. Genetic factors within AVT-II 2020/BIGHYB-5 may include alleles that accelerate the onset of male flower development, such as genes involved in floral initiation and hormone regulation. These genetic traits may trigger the expression of male flower buds at an earlier stage of plant development.

Additionally, environmental cues like temperature, light duration, and soil moisture can impact flowering and maturity time by modulating hormone levels and gene expression patterns. Consequently, the genetic predisposition of AVT-II 2020/BIGHYB-5, in conjunction with favourable environmental conditions, likely

Commented [SC54]: no need to write this word

Commented [SC58]: no need to write this word

Commented [SC59]: Environmental

Commented [SC55]: no need to write this word

Commented [SC56]: no need to write this word

Commented [SC57]: for

promotes early flowering and maturity, facilitating timely pollination and fruit set. This trait could confer advantages in bitter gourd cultivation, such as extended fruiting periods and improved yield potential. Similar findings were reported in studies on bitter gourd by **Kumar and Topno, 2022; Triveniet al., 2022.**

Yield parameters

The current study discovered that hybrid differences for the number of fruits per plant were statistically significant. The hybrid AVT-II 2020/BIGHYB-2 had maximum number of fruits per plant (28.07 fruits), followed by F₁ hybrid Aman (26.07 fruits). In AVT-II 2020/BIGHYB-8, the minimum number of fruits per plant (22.73 fruits) was noted. The hybrid Shriram Samridhi exhibited the longest fruit (13.57 cm), at par with AVT-II 2020/BIGHYB-6 (12.90 cm). The minimum fruit length (7.33 cm) was observed in AVT-II 2020/BIGHYB-1. The hybrid AVT-II 2020/BIGHYB-5 exhibited the maximum fruit diameter (40.67 mm), at par with AVT-II 2020/BIGHYB-3 (40.00mm). The minimum fruit diameter (34.00 mm) was observed in Alpine F₁ hybrid.

Hybrid AVT-II 2020/BIGHYB-1 of bitter gourd likely exhibits longer fruits than other hybrids due to specific genetic traits and environmental influences. Genetic factors within AVT-II 2020/BIGHYB-1 may include alleles that regulate fruit development processes such as cell elongation, fruit

expansion, and hormone signalling pathways. These genetic characteristics result in fruits with extended lengths during growth and maturation stages. The significance of longer fruits in hybrid AVT-II 2020/BIGHYB-1 lies in its potential to enhance marketability and consumer preference. Longer fruits typically offer more edible portions, providing better value for consumers and potentially commanding higher prices in the market. Additionally, longer fruits may contribute to increased yield per plant, thereby improving overall productivity and profitability in bitter gourd cultivation. Therefore, the genetic predisposition of hybrid AVT-II 2020/BIGHYB-1 for longer fruits presents a desirable trait with economic benefits for growers and consumers alike. The findings were earlier reported in studies on bitter gourd by **Ramyaet al., 2020; Kumar and Topno, 2022; Triveni et al., 2022.**

The hybrid AVT-II 2020/BIGHYB-6 exhibited the maximum fruit weight (53.13 grams) at par with AVT-II 2020/BIGHYB-2 having 47.60 grams. The minimum fruit weight (35.60 grams) was observed in AVT-II 2020/BIGHYB-3. The hybrid AVT-II 2020/BIGHYB-6 displayed the highest fruit yield per plant (1.38 kg/plant), comparable to AVT-II 2020/BIGHYB-2 with 1.34 kg/plant. In AVT-II 2020/BIGHYB-3, the lowest fruit yield per plant (0.82 kg/plant) was recorded.

Hybrid AVT-II 2020/BIGHYB-6 of bitter gourd likely achieves maximum fruit yield

per plant compared to other hybrids due to its advantageous genetic traits and favourable environmental interactions. Genetic factors within AVT-II 2020/BIGHYB-6 may include alleles associated with high fruit set, vigorous growth, and efficient resource utilization. These genetic characteristics contribute to the production of a greater number of fruits per plant. The significance of maximum fruit yield per plant in hybrid AVT-II 2020/BIGHYB-6 lies in its potential to increase profitability and meet market demand. Higher fruit yield translates to greater marketable produce per unit area, enhancing overall productivity and economic returns for growers. Additionally, increased yield per plant can help meet consumer demand, maintain market competitiveness, and ensure food security. Therefore, the genetic predisposition of hybrid AVT-II 2020/BIGHYB-6 for maximum fruit yield per plant presents a desirable trait with significant implications for commercial bitter gourd cultivation and sustainable agriculture. The findings were earlier reported in studies on bitter gourd by **Ramya et al., 2020; Kumar and Topno, 2022.**

Quality parameters

The highest TSS (4.30 °Brix) was displayed by the hybrid AVT-II 2020/BIGHYB-9, which was comparable to Shriram Samridhi with 4.27 °Brix. In AVT-II 2020/BIGHYB-8, the lowest TSS (2.93 °Brix) was recorded. The highest ascorbic acid content (1.67mg) was displayed

by the hybrid AVT-II 2020/BIGHYB-1, which was comparable to AVT-II 2020/BIGHYB-6 and F₁ hybrid Aman with 1.53mg. In AVT-II 2020/BIGHYB-3, the lowest ascorbic acid content (0.93mg) was recorded. The findings were earlier reported in studies on bitter gourd by **Poornima et al., 2022; Triveni et al., 2022.**

CONCLUSIONS

From the present investigation, it is concluded that among twelve hybrids of Bitter gourd, hybrid AVT-II 2020/BIGHYB-6 performed best in terms of growth & yield. The vine length was recorded 3.45 m, fruit yield was 122.48 q/ha. In terms of quality, hybrid AVT-II 2020/BIGHYB-5 was found to be best for purpose of chips. In the economic analysis of the hybrid AVT-II 2020/BIGHYB-6.

Table 1 Performance of different Hybrids of bitter gourd for various vine length, flowering and yield parameters studied

Hybrid Notation	Hybrid details	Vine length (cm)	Days to first male flower appearance	Days to first female flower appearance	Days to first fruit harvest	No of fruits per plant	Fruit length (cm)
H ₁	AVT-II 2020/BIGHYB-1	2.82	46.20	53.67	59.67	24.07	7.33
H ₂	AVT-II 2020/BIGHYB-2	3.17	53.20	53.67	60.33	28.07	10.80
H ₃	AVT-II 2020/BIGHYB-3	2.65	53.27	55.30	61.67	23.07	11.73
H ₄	AVT-II 2020/BIGHYB-4	-	-	-	-	-	-
H ₅	AVT-II 2020/BIGHYB-5	3.01	51.80	50.80	56.67	25.87	11.87
H ₆	AVT-II 2020/BIGHYB-6	3.45	52.90	54.60	60.33	25.93	12.90
H ₇	AVT-II 2020/BIGHYB-7	2.94	52.37	51.60	57.00	23.07	11.57
H ₈	AVT-II 2020/BIGHYB-8	2.51	53.67	53.83	61.33	22.73	12.87
H ₉	AVT-II 2020/BIGHYB-9	3.36	52.90	52.90	57.67	25.47	11.70
H ₁₀	Shriram Samridhi (c)	3.14	51.80	52.97	57.33	25.30	13.57
H ₁₁	Alpine F ₁ Hybrid (c)	3.02	52.90	53.80	58.00	25.23	12.80
H ₁₂	F ₁ hybrid Aman (c)	3.19	52.57	52.77	58.33	26.07	10.93
'F' Test		S	S	S	S	S	S
SE d (±)		0.22	1.19	1.15	1.27	0.80	0.83
CD. at 5%		0.45	2.49	2.40	2.65	1.66	1.73
CV (%)		8.76	2.00	2.64	2.64	3.91	8.74

Table 2 Performance of different Hybrids of bitter gourd for various yield and quality parameters studied

Hybrid Notation	Hybrid details	Fruit diameter (mm)	Fruit weight (grams)	Fruit yield per plant (kg/plant)	TSS [°Brix]	Ascorbic acid content (mg)
H ₁	AVT-II 2020/BIGHYB-1	37.67	37.57	0.90	3.37	1.67
H ₂	AVT-II 2020/BIGHYB-2	39.33	47.60	1.34	3.07	1.37
H ₃	AVT-II 2020/BIGHYB-3	40.00	35.60	0.82	3.27	0.93
H ₄	AVT-II 2020/BIGHYB-4	-	-	-	-	-
H ₅	AVT-II 2020/BIGHYB-5	40.67	47.13	1.22	3.50	1.27
H ₆	AVT-II 2020/BIGHYB-6	39.33	53.13	1.38	3.90	1.53
H ₇	AVT-II 2020/BIGHYB-7	39.67	38.03	0.88	4.17	1.50
H ₈	AVT-II 2020/BIGHYB-8	35.83	36.83	0.84	2.93	1.27
H ₉	AVT-II 2020/BIGHYB-9	39.33	46.13	1.17	4.30	1.07
H ₁₀	Shriram Samridhi	36.00	43.93	1.11	4.27	1.37
H ₁₁	Alpine F ₁ Hybrid	34.00	41.13	1.04	3.03	1.10
H ₁₂	F ₁ hybrid Aman	35.33	37.47	0.98	4.13	1.53
'F' Test		S	S	S	S	S
SE d (±)		2.04	1.92	0.07	0.15	0.14
CD. at 5%		4.26	4.01	0.15	0.32	0.29
CV (%)		6.60	5.57	8.32	5.13	12.87

Commented [SC60]: Shriram Samridhi ©

Commented [SC61]: Alpine F₁ Hybrid ©

Commented [SC62]: F₁ hybrid Aman (c)

References:

- Alam, M. A., Uddin, R., Subhan, N., Rahman, M. M., Jain, P. and Reza, H. M. (2015). Beneficial Role of Bitter Melon Supplementation in Obesity and Related Complications in Metabolic Syndrome. *Journal of Lipid*. 2: 1-18.
- Choudhary, B. (2013). Vegetables-Cucurbits, Cucumber nutritional quality. *National Book Trust India. Reprint edition*. 142 pp.
- Dandawate, P. R., Subramaniam, D., Padhye, S. B. and Anant, S. (2016). Bitter melon: a panacea for inflammation and cancer. *Chinese Journal of Natural Medicines*. 14(2): 81-100.
- Fisher, R. A. and Yates, F. (1963). Statistical Tables for Biological, Agricultural and Medical Research. *Oliver and Boyd, London*: 143 p.
- Heiser, C. B. (1979). The gourd books. University of Oklahoma Press, Norman, UK, pp 236.
- Horax, R., Hettiarachchy, N. and Chener, P. (2010). Extraction, Quantification, and Antioxidant Activities of Phenolics from Pericarp and Seeds of Bitter Melons (*Momordica charantia* Harvested at Three Maturity Stages (Immature, Mature, and Ripe). *Journal of Agricultural and Food Chemistry*. 58: 4428-4433.
- Islam, S., Jalaluddin, M. and Hettiarachchy, N. S. (2011). Bioactive compounds of bitter melon genotypes (*Momordica charantia* L.) in relation to their physiological functions. *Functional Foods in Health and Disease*. 2: 61-74.
- Joseph, B.; Jini, D. (2013). "Antidiabetic effects of *Momordica charantia* (bitter melon) and its medicinal potency". *Asian Pacific Journal of Tropical Disease*. 3(2): 93-102.
- Kandangath, R. A., Garlapati, P. K., and Nallamuthu, I. (2015). Nutritional, pharmacological, and medicinal properties of *Momordica charantia*. *International Journal of Nutrition and Food Sciences* 4: 75-83.
- Kumar, A. M., Naik, K. M., Pathare, J., Balfour, D. and Kotecha, P. M. (2016). Studies on osmo-air drying of bitter gourd chips: physico-chemical composition. *International Journal of Advanced Scientific Technical Research*. 3: 175-196.
- Kumar, P. R. and Topno, S. E. (2022). Performance of bitter gourd (*Momordica charantia* L.) hybrids

- under Prayagraj Agro-climatic condition. *The Pharma Innovation Journal*. 11(5): 2181-2184.
- Kumari, A., Shree, S., Kumar, R., Haque, M., Kishor, C., & Singh, V.K. (2019). Study of *Per se* Performance of Parents and Hybrids for Yield and Quality of Bitter gourd (*Momordica charantia* L.). *International Journal of Current Microbiology and Applied Sciences*. 8(7): 1781-1789.
- Lee, S. Y., Eom, S. H., Kim, Y. K., Park, N. I. and Park, S. U. (2009). Cucurbitane-type triterpenoids in *Momordica charantia* Linn. *Journal of Medicinal Plants Research*. 3(13):1264-1269.
- Liu, X. R., Deng, Z. Y., Fan, Y. W., Li, J. and Liu, Z. H. (2010). Mineral elements analysis of *Momordica charantia* seeds by ICP-AES and fatty acid profile identification of seed oil by GCMS. *Guang Pu Xue Yu Guang Pu Fen Xi*. 30(8): 2265-8.
- McKay, J. W. (1930). Chromosome numbers in the Cucurbitaceae. *Botanical Gazette*. 89:416-417.
- NHB, (2021). nhb.gov.in/statistics/2020-21. Area and Production of Horticulture Crops- All India. Visited on 08/12/2022.
- Poornima, V., Rajasree, V., Thangamani, C., ManikandaBoopathi, N., & Manoranjitham, S. K. (2022). Evaluation of Interspecific Hybrids for Yield and Quality Characteristics in Bitter Gourd (*Momordica charantia* L.). *International Journal of Plant & Soil Science*. 34(22): 655-662.
- Ramya, B., Kerketta, A. and Topno, S. E. (2020). Evaluation of Different Hybrids for Growth and Yield Attributes of Bitter Gourd (*Momordica charantia* L.) in Prayagraj Region. *International Journal of Current Microbiology and Applied Sciences*. 9(12): 1008-1012.
- Wang, L., Li, Q., Cao, J., Cai, T. and Jiang, W. (2007). Keeping quality of fresh-cut bitter gourd (*Momordica charantia*.) at low temperature of storage. *Journal of Food Processing and Preservation*. 31: 571-582.

UNDER PEER REVIEW

