

**Antibiotic resistance profile and safety assessment  
of *Lactobacillus acidophilus* and *Lactobacillus  
plantarum* isolated from milk**

**ABSTRACT**

**Aims:** The present study was undertaken to evaluate the safety of probiotics viz., *Lactobacillus acidophilus* and *Lactobacillus plantarum* through antibiotic susceptibility tests.

**Study design:** The study was conducted on two reference *Lactobacillus* strains viz., *Lactobacillus acidophilus* (NCDC 13) and *Lactobacillus plantarum* (NCDC 20) procured from the National Collection of Dairy Cultures (NCDC), ICAR-NDRI, Karnal, India.

**Place and Duration of Study:** Department of Veterinary Public Health and Epidemiology, College of Veterinary and Animal Sciences, G.B. Pant University of Agriculture & Technology, Pantnagar, Uttarakhand, between March - April 2021.

**Methodology:** The strains of *L. acidophilus* and *L. plantarum* were subjected to antibiotic susceptibility tests using the disc diffusion method against 26 antibiotics.

**Results:** The isolates were found to exhibit multiple resistance against some of the most commonly used antibiotics. The isolates showed a high level of resistance toward ampicillin, amoxicillin, cefotaxime, nalidixic acid, streptomycin, kanamycin and nitrofurantoin. The isolates showed low levels of resistance toward cephalothin, amikacin, erythromycin and azithromycin. They were susceptible to ciprofloxacin, penicillin G, cloxacillin, ofloxacin, norfloxacin, levofloxacin, moxifloxacin, sparfloxacin, enrofloxacin, gemifloxacin, chloramphenicol, gentamicin, co-trimoxazole and oxytetracycline.

**Conclusion:** The current study concluded that antibiotic resistance is prevalent among *L. acidophilus* and *L. plantarum*, which is a major concern for food safety. Hence, antibiotic susceptibility tests should be considered as an essential measure for the assessment of the safety of probiotics. Furthermore, studies to evaluate the presence of antibiotic-resistant genes in commercially available probiotics should be conducted.

*Keywords: Lactobacilli; safety; probiotics; antibiotic resistance.*

## **1. INTRODUCTION**

Lactic acid bacteria (LAB) are Gram-positive bacteria that can produce lactic acid as an end product of carbohydrate fermentation. They are widely used in food production and preservation (Lahtinen *et al.*, 2011). Of all the genera of LAB, *Lactobacillus* is the most economically important microbiota which is harbored mainly in the gut of man and animals (Tannock, 2005). They are beneficial for human health because of their antimicrobial and immunomodulatory activity. They can also help to restore the healthy gut microflora. Keeping in view of their benefits, they are included in our diet as probiotics, food preservatives or starters (Salminen *et al.*, 1998; Rozman *et al.*, 2020). There have been numerous studies on the commercial production and use of *Lactobacillus*-based probiotics because of their 'generally recognized as safe' (GRAS) status (García-Fruitós, 2012).

Antibiotics are an important therapeutic tool in tackling many infectious diseases of bacterial origin (Aminov, 2017). However, misuse and exploitation of antibiotics by human beings in animal husbandry as well as medical fields has caused the emergence of antibiotic-resistant bacteria (EFSA/ECDC, 2018), which has become a global cause of health concern (World Health Organization, 2014). The bacteria may become resistant in two ways i.e. naturally to escape the action of antibiotics or genetically horizontally resistant gene transfer through transposons or plasmids (Belletti *et al.*, 2009). Furthermore, there has been a rise in the frequency of documented cases of antibiotic-resistant LAB strains over the past decade. To assess the risk, it is important to evaluate if commonly used lactobacillus species isolates display phenotypic resistance to antibiotics and if they do, to determine the extent of resistance and identify the responsible genes. Standardized methods have been developed to determine levels of antibiotic resistance and, as they allow comparisons of results between laboratories, they are recommended by several international agencies including the Clinical and Laboratory Standards Institute (CLSI), International Organization for Standardization (ISO), European Food Safety Authority (EFSA) and the International Dairy Federation (IDF).

Since, probiotics are used in doses unlimited, ensuring their safety is of utmost importance, while there are as such no legislation/acts/rules for microorganisms that are deliberately added to our food as probiotics. It is therefore recommended that these products follow similar requirements as feed additives as precautionary measures (EFSA, 2007). Several reviews (Saarela *et al.*, 2000) on LAB have recommended safety criteria for probiotics such as the absence of ARGs which are responsible for resistance to clinically important antibiotics (EFSA-FEEDAP, 2018). Although LABs are

regarded as safe, they can act as a reservoir for ARGs, which can persist in our food chain (Guo *et al.*, 2017; Duranti *et al.*, 2017) and can be responsible for the transfer of ARGs via horizontal gene transfer (Liu *et al.*, 2012). Some studies have reported the transfer of ARGs between various lactobacilli (Nawaz *et al.*, 2011). *In vitro* and *in vivo* studies have also been conducted to evaluate the transfer of ARGs from lactic acid bacteria to other pathogenic bacteria (Kazimierczak *et al.*, 2006; Jacobsen *et al.*, 2007). Thus, the present study was conducted to characterize the phenotypic antibiotic resistance profile in isolates of *Lactobacillus acidophilus* and *Lactobacillus plantarum* using the disc diffusion method. It provides an insight into the safety of probiotics available in terms of antibiotic resistance.

## 2. MATERIAL AND METHODS

### Bacterial strains and propagation

Two reference *Lactobacillus* strains viz., *Lactobacillus acidophilus* (NCDC 13) and *Lactobacillus plantarum* (NCDC 20) were procured from the National Collection of Dairy Cultures (NCDC), ICAR-NDRI, Karnal, India. *Lactobacilli* were maintained and propagated in *Lactobacillus de Man Rogosa Sharpe* (MRS) broth (Hi-media Laboratories Pvt. Ltd.). It was prepared according to the instructions of the manufacturers. The final pH of the media was maintained at  $6.5 \pm 0.2$ . Strains were incubated at 37 °C for 48 h in anaerobic atmospheric conditions. They were further sub-cultured thrice before the experiments. It was followed by plating on MRS agar under the same above-mentioned conditions. The bacterial cultures were preserved in glycerol stocks at -80° C.

### Antibiotic discs

Twenty-six antibiotics that are used most commonly from different classes were purchased from Hi-media Laboratories Pvt. Ltd. Mumbai, India, and tested for procured probiotics. The details of the name of the drug, concentration, antibiotic group, and mode of action are described in Table 1.

**Table 1. Antibiotics used for antibiotic-resistant profile and their mode of action**

| S. No. | Name of drug | Concentration<br>(mcg) | Group antibiotics | Mode of action                 |
|--------|--------------|------------------------|-------------------|--------------------------------|
| 1.     | Ampicillin   | 10                     | β-Lactams         | Cell wall synthesis inhibition |
| 2.     | Amoxycillin  | 30                     |                   |                                |
| 3.     | Cloxacillin  | 30                     |                   |                                |

---

|     |                            |                 |                                    |   |
|-----|----------------------------|-----------------|------------------------------------|---|
| 4.  | Ampicillin/<br>cloxacillin | 10              |                                    |   |
| 5.  | Penicillin G               | 10 <sup>a</sup> |                                    |   |
| 6.  | Cephalothin                | 30              | First generation<br>Cephalosporins |   |
| 7.  | Cefotaxime                 | 10              | Third generation<br>Cephalosporins |   |
| 8.  | Ciprofloxacin              | 5               | Quinolones                         | DNA replication and transcription<br>inhibition |
| 9.  | Ofloxacin                  | 5               |                                    |   |
| 10. | Nalidixic acid             | 30              |                                    |   |
| 11. | Norfloxacin                | 10              | Fluoroquinolones                   |   |
| 12. | Levofloxacin               | 5               |                                    |   |
| 13. | Moxifloxacin               | 5               |                                    |   |
| 14. | Sparfloxacin               | 5               |                                    |   |
| 15. | Enrofloxacin               | 10              |                                    |   |
| 16. | Gemifloxacin               | 5               |                                    |   |
| 17. | Gentamicin                 | 10              | Aminoglycosides                    | Protein synthesis inhibition                    |
| 18. | Streptomycin               | 300             |                                    |   |
| 19. | Amikacin                   | 30              |                                    |   |
| 20. | Kanamycin                  | 30              |                                    |   |
| 21. | Chloramphenicol            | 30              | Other                              |   |
| 22. | Erythromycin               | 15              | Macrolides                         |   |
| 23. | Azithromycin               | 15              |                                    |   |

---

|     |                 |     |               |   |
|-----|-----------------|-----|---------------|---|
| 24. | Nitrofurantoin  | 300 | Other         |   |
| 25. | Oxytetracycline | 30  | Tetracyclines |   |
| 26. | Co-Trimoxazole  | 25  | Other         | Folic acid synthesis inhibitors or anti-metabolites |

<sup>a</sup> denotes concentration in units

### Antimicrobial susceptibility testing

The sensitivity or resistance of LAB to most commonly used antibiotics was evaluated by an antimicrobial susceptibility test. The standard disc diffusion assay was performed according to the Kirby–Bauer method (Bauer *et al.*, 1966). Isolates were cultured and grown overnight in MRS broth and 100 µl culture (0.5 McFarland equivalent to 10<sup>8</sup> cfu/ml) was spread on Mueller-Hinton agar plates with the help of an L spreader. Antimicrobial disks will be placed with the help of sterile forceps and incubated at 37°C for 24 h under anaerobic conditions. The zone of inhibition diameter was measured by zone reader (Hi Antibiotic zone scale, Hi-Media) and results were read according to the breakpoints recommended by Clinical and Laboratory Standard Institute standards for disc-diffusion assay (CLSI, 2016) (Table 2) as described by Sharma *et al.* (2017).

**Table 2. Susceptible and resistant strains were evaluated after being compared with known standard given by CLSI (2016)**

| Disc Diffusion Method | Diameter of zone of inhibition (mm) |
|-----------------------|-------------------------------------|
| Susceptible           | >20                                 |
| Intermediate          | 15–19                               |
| Resistant             | ≤14                                 |

### Statistical evaluation

The disc diffusion method was performed in triplicate and the average diameters calculated are presented as resistant (R), sensitive (S) or intermediate (I).

### 3. RESULTS AND DISCUSSION

Antimicrobial disc susceptibility tests were performed according to the procedures described by CLSI (CLSI, 2016). Comparative evaluation of the diameter of the zone of inhibition has been shown in graphical form (Fig. 1). The growths of both tested LAB strains i.e. *L. acidophilus* and *L.*

*plantarum* were homogenous over MRS. The results for the reference LAB strains have been documented in terms of resistant (R), susceptible (S) and intermediate (I) (Table 3).




Fig. 1. Comparative evaluation of Diameter of zone of inhibition of *L. acidophilus* and *L. plantarum*

Table 3. Susceptibility of LAB to commonly used antibiotics using the disc diffusion method

| S. No. | Antibiotics            | Inhibition zone diameter range (mm) |                                |
|--------|------------------------|-------------------------------------|--------------------------------|
|        |                        | <i>Lactobacillus acidophilus</i>    | <i>Lactobacillus plantarum</i> |
| 1      | Ampicillin             | R (11)                              | R (13)                         |
| 2      | Amoxycillin            | R (10)                              | R (8)                          |
| 3      | Cloxacillin            | S (37)                              | S (34)                         |
| 4      | Ampicillin/cloxacillin | S (20)                              | I (19)                         |
| 5      | Penicillin G           | S (21)                              | S (23)                         |
| 6      | Cephalothin            | I (19)                              | R (13)                         |
| 7      | Cefotaxime             | R (14)                              | R (13)                         |
| 8      | Ciprofloxacin          | S (35)                              | S (38)                         |
| 9      | Ofloxacin              | S (25)                              | S (23)                         |
| 10     | Nalidixic acid         | R (10)                              | R (10)                         |
| 11     | Norfloxacin            | S (24)                              | S (23)                         |
| 12     | Levofloxacin           | S (25)                              | S (33)                         |
| 13     | Moxifloxacin           | S (24)                              | S (25)                         |
| 14     | Sparfloxacin           | S (23)                              | S (23)                         |
| 15     | Enrofloxacin           | S (25)                              | S (25)                         |
| 16     | Gemifloxacin           | S (36)                              | S (33)                         |

|    |                 |        |        |
|----|-----------------|--------|--------|
| 17 | Gentamicin      | S (27) | S (24) |
| 18 | Streptomycin    | R (13) | R (14) |
| 19 | Amikacin        | S (23) | I (15) |
| 20 | Kanamycin       | R (13) | R (13) |
| 21 | Chloramphenicol | S (39) | S (39) |
| 22 | Erythromycin    | I (17) | I (18) |
| 23 | Azithromycin    | I (17) | S (25) |
| 24 | Nitrofurantoin  | R (10) | R (8)  |
| 25 | Oxytetracycline | S (20) | S (28) |
| 26 | Co-Trimoxazole  | S (38) | S (34) |

In our study, phenotypic resistance to ampicillin, amoxicillin, cefotaxime, nalidixic acid, streptomycin, kanamycin and nitrofurantoin was exhibited by *L. acidophilus*. While *L. plantarum* showed phenotypic resistance to cephalothin in addition to the above-mentioned antibiotics. There were no significant differences observed in the antibiotic resistance profile of *L. acidophilus* and *L. plantarum*. A low level of resistance was exhibited by *L. acidophilus* toward cephalothin, erythromycin and azithromycin. Whereas, *L. plantarum* showed a low level of resistance toward ampicillin/cloxacillin, amikacin and erythromycin. High susceptibility was exhibited by both isolates toward cell wall synthesis inhibitors ( $\beta$ -Lactams- cloxacillin and penicillin G), DNA replication and transcription inhibitors (Quinolones- ciprofloxacin and ofloxacin; Fluoroquinolones) and protein synthesis inhibitors (chloramphenicol, oxytetracycline and co-trimoxazole). There was no significant difference in the antibiotic susceptibility pattern of *L. acidophilus* and *L. plantarum*.

Lactobacilli are generally considered susceptible to the cell wall synthesis inhibitors ( $\beta$ -Lactams) (Guo *et al.*, 2017; Gueimonde *et al.*, 2013) and more resistant towards cephalosporins (Karapetkov *et al.*, 2011) which corroborated with our study. In our study, *L. acidophilus* displayed susceptibility toward  $\beta$ -lactam antibiotics except for ampicillin and amoxicillin for *L. acidophilus* and *L. plantarum* which was contrary to the findings of Klare *et al.* (2007) and Nawaz *et al.* (2011). In addition, *L. plantarum* showed intermediate susceptibility towards ampicillin/cloxacillin while *L. acidophilus* was susceptible to it. Resistance towards cephalosporins in our study for both strains was reported which was similar to the study conducted by Karapetkov *et al.* (2011). Lactobacilli were reported to have intrinsic resistance to aminoglycosides (Abriouel *et al.*, 2015; Sharma *et al.*, 2017; Štšepetova *et al.*, 2017; Zhou *et al.*, 2005) which was similar to our study except for gentamicin

and amikacin. In a study conducted by Pell *et al.* (2021), they also observed the susceptibility of *L. plantarum* towards gentamicin. Lactobacilli are considered to have intrinsic resistance to aminoglycosides because of the absence of cytochrome-mediated electron transport, which is responsible for antibiotic uptake (Charteris *et al.*, 2001). Resistance to other antibiotics varies greatly among lactobacilli. The genetic determinants often present in mobile elements like transposons and plasmids have the potential to transfer antibiotic resistance genes primarily through conjugation mechanisms (Aquilanti *et al.*, 2007). Another mechanism of resistance against antimicrobials is known to be present in certain lactobacilli which are mediated by multidrug resistance (MDR) transporters (Gueimonde *et al.*, 2009).

#### **4. CONCLUSION**

Foods and food supplements having naturally occurring or intentionally added bacteria, such as probiotics can serve as a potential reservoir for antibiotic resistance genes. Hence, the safety of probiotics requires the assessment of resistance /resistance genes carried by them against the clinically significant antimicrobials. Future research exploring their presence and transferability will certainly be a step towards safety in true terms. As the probiotics in our study possessed resistance levels exceeding the limit recommended by CLSI, it is suggested that the proposed limit should be re-examined. Regulatory guidelines/legislation for the assessment of the safety of lactobacilli for their approval as starter cultures or probiotics should be made. It will also facilitate screening of probiotics from a safety point of view. Further studies can be done to evaluate the transferability of genes responsible for resistance to most commonly used antibiotics.

## REFERENCES

1. Abriouel H, Casado Muñoz MDC, Lerma LL, Montoro PB, Bockelmann W, Pichner R, et al. New insights in antibiotic resistance of *Lactobacillus* species from fermented foods. *Food Res Int.* 2015;78:465–81.
2. Aminov R. History of antimicrobial drug discovery: Major classes and health impact. *Biochem Pharmacol.* 2017;133:4-19.
3. Ammor MS, Flòrez AB, Mayo B. Antibiotic resistance in non-enterococcal lactic acid bacteria and bifidobacteria. *Food Microbiol.* 2007;24:559-70.
4. Aquilanti L, Garofalo C, Osimani A, Silvestri G, Vignaroli C, Clementi F. Isolation and molecular characterization of antibiotic-resistant lactic acid bacteria from poultry and swine meat products. *J Food Prot.* 2007;70:557–65.
5. Bauer AW, Kirby WMM, Sherris JC, Turk, M. 1966. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol.* 1966;45:493–96.
6. Belletti N, Gatti M, Bottari B, Neviani E, Tabanelli G, Gardini F. Antibiotic resistance of lactobacilli isolated from two Italian hard cheeses. *J Food Prot.* 2009;72:2162–69.
7. Charteris WP, Kelly PM, Morelli L, Collins JK. Gradient diffusion antibiotic susceptibility testing of potentially probiotic lactobacilli. *J Food Prot.* 2001;64:2007-14.
8. CLSI. Wayne PA. Performance standards for antimicrobial susceptibility testing: 26th informational supplement. M100-S26. 2016; pp 1-25.
9. Duranti S, Lugli GA, Mancabelli L, Turrone F, Milani C, Mangifesta M, et al. 2017. Prevalence of antibiotic resistance genes among human gut-derived bifidobacteria. *Appl Environ Microbiol.* 2017;83:e02894-16.
10. EFSA. Opinion of the Scientific Committee on a request from EFSA on the introduction of a Qualified Presumption of Safety (QPS) approach for assessment of selected microorganisms referred to EFSA. *EFSA J.* 2007;587:1-16
11. EFSA/ECDC. The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2016. *EFSA J.* 2018;16:5182.
12. EFSA-FEEDAP. Guidance on the characterization of microorganisms used as feed additives or as production organisms. *EFSA J.* 2018;16:5206.
13. Gueimonde M, Flórez AB, de los Reyes-Gavilán CG, Margolles A. Intrinsic resistance in lactic acid bacteria and bifidobacteria: the role of multidrug resistance transporters. *Int J Probiotics Prebiotics.* 2009;4:181–86.

14. Gueimonde M, Sánchez B, de Los Reyes-Gavilán CG, Margolles, A. Antibiotic resistance in probiotic bacteria. *Front Microbiol.* 2013;4:202.
15. Guo H, Pan L, Li L, Lu J, Kwok L, Menghe B, et al. Characterization of antibiotic resistance genes from *Lactobacillus* isolated from traditional dairy products. *J Food Sci.* 2017;82:724–30.
16. Jacobsen L, Wilcks A, Hammer K, Huys G, Gevers D, Andersen SR. Horizontal transfer of *tet(M)* and *erm(B)* resistance plasmids from food strains of *Lactobacillus plantarum* to *Enterococcus faecalis*JH2-2 in the gastrointestinal tract of gnotobiotic rats. *FEMS Microbiol Ecol.* 2007;59:158–166.
17. Karapetkov N, Georgieva R, Rumyan N, Karaivanova E. Antibiotic susceptibility of different lactic acid bacteria strains. *Benef Microbes.* 2011;2(4):335-39.
18. Kazimierczak KA, Flint HJ, Scott KP. Comparative analysis of sequences flanking *tet(W)* resistance genes in multiple species of gut bacteria. *Antimicrob Agents Chemother.* 2006;50:2632–39.
19. Klare I, Konstabel C, Werner G, Huys G, Vankerckhoven V, Kahlmeter G, et al. Antimicrobial susceptibilities of *Lactobacillus*, *Pediococcus* and *Lactococcus* human isolates and cultures intended for probiotic or nutritional use. *J Antimicrob Chemother.* 2007;59:900–12.
20. Liu L, Chen X, Skogerbø G, Zhang P, Chen R, He S. et al. The human microbiome: a hot spot of microbial horizontal gene transfer. *Genomics.* 2012;100: 265–70.
21. Nawaz M, Wang J, Zhou A, Ma C, Wu X, Moore JE, et al. Characterization and transfer of antibiotic resistance in lactic acid bacteria from fermented food products. *Curr Microbiol.* 2011;62:1081–89.
22. Pell LG, Horne RG, Huntley S, Rahman H, Kar S, Islam MS, et al. Antimicrobial susceptibilities and comparative whole genome analysis of two isolates of the probiotic bacterium *Lactiplantibacillus plantarum*, strain ATCC 202195. *Sci Rep.* 2021;11(1):1-18.
23. Rozman V, Lorbeg PM, Accetto T, Matijašić BB. Characterization of antimicrobial resistance in lactobacilli and bifidobacteria used as probiotics or starter cultures based on integration of phenotypic and in silico data. *Int J Food Microbiol.* 2020;314:108388.
24. Saarela M, Mogensen G, Fondén R, Mättö J, Mattila-Sandholm T. Probiotic bacteria: safety, functional and technological properties. Review article. *J Biotech.* 2000;84:197–215.

25. Salminen S, von Wright A, Morelli L, Marteau P, Brassart D, de Vos WM, et al. Demonstration of safety of probiotics—a review. *Int J Food Microbiol.* 1998;44:93–106
26. Sharma C, Gulati S, Thakur N, Singh BP, Gupta S, Kaur, S., et al. Antibiotic sensitivity pattern of indigenous lactobacilli isolated from curd and human milk samples. *3 Biotech.* 2017;7:1-14. DOI: <https://doi.org/10.1007/s13205-017-0682-0>.
27. Štšepetova J, Taelma H, Smidt I, Hütt P, Lapp E, Aotäht E, et al. Assessment of phenotypic and genotypic antibiotic susceptibility of vaginal *Lactobacillus* sp. *J Appl Microbiol.* 2017;123:524–34
28. World Health Organization. 2014. Antimicrobial Resistance. Global Report on Surveillance. World Health Organization, Geneva. 2014. Accessed 7 August 2021. Available: <http://www.who.int/iris/handle/>
29. Zhou JS, Pillidge CJ, Gopal PK, Gill HS. Antibiotic susceptibility profiles of new probiotic *Lactobacillus* and *Bifidobacterium* strains. *Int J Food Microbiol.* 2005;98:211-17.