

# Impact of plant growth regulators on growth, fruit yield and quality of strawberry (*Fragaria × ananassa* Duch.) cv. Winter dawn

## Abstract

The present study aimed to investigate the impact of plant growth regulators on the growth, yield, and quality of strawberry (*Fragaria × ananassa* Duch) cv. Winter Dawn. A total of ten treatments were examined, including control (Water Spray), individual treatments of NAA (Naphthalene Acetic Acid) and GA<sub>3</sub> (Gibberellic Acid) and BA (6-benzyladenine). The experiment was conducted under controlled conditions, and various parameters related to vegetative growth, quality, and yield were evaluated.

The results demonstrated that treatment T<sub>9</sub>, combination BA (6-benzyladenine) 75 ppm exhibited superior vegetative growth characteristics. Plants treated with T<sub>9</sub> displayed increased plant height, number of leaves, plant spread, petiole length, chlorophyll content, and early onset of flowering and first fruiting. Moreover, T<sub>9</sub> also demonstrated remarkable performance in terms of fruit production, with a higher number of fruits, flowers, and excellent fruit set, and TSS, Acidity (%), and Ascorbic acid.

These findings highlight the beneficial impact of plant growth regulators on the growth, yield, and quality of strawberry plants. The combined application of BA (6-benzyladenine) 75 ppm (T<sub>9</sub>) resulted in superior vegetative growth. These results suggest that the incorporation of plant growth regulators in strawberry cultivation practices can be a promising approach to optimize both yield and fruit quality. Further studies are warranted to understand the underlying mechanisms and optimize the application dosage and timing of these bio inputs for sustainable strawberry production.

*Keywords: Naphthalene acetic acid, Gibberellic acid, BA(6-Benzyladenine), Winter dawn.*

## **Introduction**

“Strawberry is one of the most delicious, attractive, nutritious and refreshing soft fruits of the world. The cultivated strawberry (*Fragaria × ananassa*) is a hybrid of two native American sp; *F. chiloensis* and *F. virginiana*. Strawberries are good source of natural anti-oxidant. Owing to its medicinal properties (anticarcinogenic, antidiabetic and antioxidant), strawberry is gaining popularity among all age group consumers. Strawberries are good source of natural anti-oxidant. In India, it is mainly grown in Maharashtra and in hills of Himachal Pradesh, J&K and Uttarakhand, Uttar Pradesh and Haryana”. [25] “Strawberry usually grows in temperate zones, and it is classified as a short-day plant based on its behaviour and eco system. Strawberry is a temperate plant that can be grown in both farmlands and mountains, but the fruit results are better on the hills. For best progress and growth, the strawberry needed a day temperature of 22°C and a night temperature of 7°C to 13°C. Frost and winter injuries decreased yield. Strawberry might cultivate in particular land, extending from thick clay to

light sand and gravel. Strawberry plants, on the other hand, grow on sandy loam soil with a pH of 5.5 to 6.5. It is a heavy conveyor crop that produces more”(Yadav *et al.*, 2018). “The main objective of the strawberry growers is to produce a fruit with appealing appearance (size, color and shape) not necessary accompanied by the same appealing tasteful characteristics. In order for the farmers to achieve such fruit growth enhancement, they often use plant growth regulating compounds. Plant growth regulators (PGRS) have proven their role in augmenting yield and quality in many fruits. Some of plant bioregulators are synthesized endogenously, but occasionally they are needed to get supplemented exogenously for additional stimulus for short duration crop like strawberry, which require quick response for increased growth, fruit set and yield. Use of plant bioregulators plays an important role in vegetative growth, flowering, yield and quality. Use of GA3 in strawberry has been reported in early flowering, increased duration of flowering, harvesting and yield. It increases yield and quality of fruits, helps in cell elongation and cell enlargement, increases vegetative growth and minimizes time of maturity and increases fruit set. Application of NAA increases fruit size and delays ripening and increases anthocyanin accumulation in strawberry fruits. It also increases duration of flowering, improves yield and quality of fruits. BA, as a plant growth regulator enhances the size and shape of fruits, lateral bud break and lateral shoot growth, leading to improved branching in fruit trees and flowering. BA increases fruit size and delay chlorophyll breakdown and fruit ageing. BA also decreases loss in firmness, delay ethylene production, decreases respiration rate and induces mechanical resistance which reduces senescence rate after harvest”. [25] “In India, strawberry fruit crop is still grown in open areas using a paddy straw mulching strategy by poor or marginal farmers, and it accounts for a large part of the national annual strawberry cultivable land. According to scientific findings, the strawberry plant responded positively to the application of a growth regulator” (Sharma and Sharma, 2004). “Because of their suitability for treatment at a lower cost, naphthalene acetic acid (NAA) and gibberellins (GA3) have been widely investigated in present agricultural systems. In many fruits, the role of these plant growth regulators has been explored”(Bist *et al.*, 2018). Premature flowering, enhanced flowering time, collecting, and yield have all been observed with the use of GA3 in strawberries. It boosts fruit output and quality, promotes cell elongation and expansion, helps to improve vegetative growth, reduces the period to maturity, and boosts fruit set (Sharma and Singh, 2009). In strawberry fruits, NAA promotes growth parameters, slows ripening, and enhances anthocyanin accumulation. It furthermore enhances the blooming period and improve fruit output and quality (Mir *et al.*, 2004).

## Materials and Methods

The experiment was conducted during 2023-2024 at experimental field of department of horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj to experiment the impact of plant growth regulators on growth, yield, and quality of strawberry. The practical was carryout in a randomized block design with seven treatments consisted of control (no application of plant growth regulators), GA3 (25, 50, 75 ppm), NAA (25, 50, 75 ppm), and BA (25, 50, 75ppm). All the treatment were replicated thrice. All the runners are equal and vigour transplanted during morning hours at a 30cm x 30cm. Daily watering was done for first week with drip irrigation and gap filling was done after first week of transplanted. After the transplant, the plant growth regulators were sprayed at 30, 60, and 90 days. In each replication, the observation was noted

in three randomly selected plants. According to A.O.A.C (1990), the quality of the fruits was tested during harvest using a standard method and procedure, and the results were statistically analysed. Randomized Block Design was used to set up the experiment, and ten treatments were reproduced three times. The ten treatments consist of (Water spray) T0, NAA (Naphthalene Acetic Acid) 25ppm T1, NAA (Naphthalene Acetic Acid) 50ppm T2, NAA (Naphthalene Acetic Acid) 75ppm T3, GA3 (Gibberellic Acid) 25ppm T4, GA3 (Gibberellic Acid) 50ppm T5, GA3 (Gibberellic Acid) 75ppm T6, BA(6-benzyladenine) 25ppm T7, BA(6-benzyladenine) 50ppm T8, BA(6-benzyladenine) 75ppm T9. All the doses of NAA, GA3, BA combination was applied at the time of planting and during flowering initiation and observations were recorded on plant height (cm), number of leaves per plant, plant spread (cm), petiole length (cm), Days to first flowering, number of flowers per plant, number of fruits per plant, diameters of fruits, fruits length, fruit weight, fruit set, TSS, acidity, ascorbic acid. The statistical procedure for agricultural research states that an analysis of variance will be performed on the data's mean values. Randomized Block Design. A method and algorithms were used to compute different statistical parameters. The Analysis of Variance (ANNOVA) method was used to compare the means of the attributes.

### List 1. Treatment Details

Treatment	Treatment Details
T0	Control
T1	NAA (Naphthalene Acetic Acid) 25ppm
T2	NAA (Naphthalene Acetic Acid) 50ppm
T3	NAA (Naphthalene Acetic Acid) 75ppm
T4	GA3 (Gibberellic Acid) 25ppm
T5	GA3 (Gibberellic Acid) 50ppm
T6	GA3 (Gibberellic Acid) 75ppm
T7	BA (6-benzyladenine) 25ppm
T8	BA (6-benzyladenine) 50ppm
T9	BA (6-benzyladenine) 75ppm

### Abbreviations

C.D. = Critical difference

F = Fertilizer

F test S = F test significant

S = Spacing

SE (d) = Standard error of difference

### Vegetative characters

#### . Plant Height (cm)

The height of plants was measured by using a measuring scale from crown level of plants to the apex of primary leave at was recorded at 30, 60 and 90 DAP and results were expressed in cm.

### **. Plant Spread (cm)**

The spread of the tagged plants was recorded at 30, 60 and 90 DAP in east-west and north-south direction separately with the help of a meter scale and the average for each direction was calculated.

### **. Petiole length (cm)**

The petiole is the stalk of the entire leaf, but for the operation of this key this feature is applied also to the leaflet stalk of compound leaves. It is measured at 30, 60 and 90. Their average was calculated and subjected to statistical analysis.

### **. Numbers of leaves per plant**

Total number of leaves was counted from tagged plants in each replication at 30, 60 and 90 DAP and expressed as average number of leaves per plant.

### **Floral characters**

#### **. Days to first flowering**

It was recorded after planting when 5-6 plants in each replication started to flower. The average number of days from planting date was calculated to make the observation.

#### **. No of flowers/plants**

The total number of flowers per plant was recorded on the five tagged plants. The number of flowers was counted from first flower initiation after planting till last harvesting and the value was expressed as number of flowers per plant.

### **Fruit and yield attributes**

#### **.Number of fruits per plant**

It was recorded after planting when 5-6 plants in each replication started to fruiting. The average number of days from planting date was calculated to make the observation.

#### **. Number of fruits/plants**

The number of fruits per plant was recorded on the same three tagged plants on which fruit set was studied. The number of fruits reaching harvestable maturity was counted at each harvest and the value was expressed as number of fruits per plant.

#### **Length – diameter of the fruit (cm)**

The length diameter ratio is the ratio of the flighted length of the fruit to its outside diameter of the fruit. The ratio calculation is calculated by dividing the flighted length of the fruit by its nominal diameter.

#### **FruitWeight(g)**

Selected fruit were harvested from each replication to measure the fruit weight. The weight was measured on electronic balance and average berry weight was calculated and expressed in grams (g).

#### **Fruit length (mm)**

Fruit length of 10 fruits from each treatment was taken with the help of digital vernier caliper. The average length was then worked out to record the data.

### . Fruit set (%)

For each stem, calculate the percent fruit set as follows: divide the number of fruits by the number of blossoms, then multiply by 100.

### Quality parameters

#### . Total soluble solid (°Brix).

Total soluble solids (TSS) were recorded with the help of digital refractometer. Fully ripe fruits of each treatment were taken and few (2-3) drops of juice from 5 fruits were taken separately and dropped over the prism of the refractometer. The value as observed was averaged to record the TSS (°Brix).

#### . Ascorbic acid (mg/100g of fresh fruits)

Ascorbic acid content was estimated by grinding 5-gram fruit pulp with 30 per cent metaphosphoric acid as buffer. The extract was filtered with muslin cloth and appropriate volume was made. A suitable aliquot was titrated against 2,6-dichlorophenol dye solution till light pink colour appeared. The result was calculated with help of following formula and expressed as milligram ascorbic acid per 100 gram of fruit pulp. (AOAC 1975)

#### . Acidity (%)

Titrate acidity (% malic acid) was measured by using a standard procedure of Hortwitz (1980) with a slight modification. For this, a known weight of the fruit sample was crushed and taken in a 100ml volumetric flask and the volume was made up by adding distilled water. Add filtration, 10 ml of the filtrate was taken in a separate conical flask and titrated against 0.1 N sodium hydroxide (NaOH) using phenolphthalein dye as an indicator. The end point was determined by the appearance of a faint pink colour. Titrate acidity was calculated by using the formula given below:

Titrate acidity (%)

$$= \frac{\text{Titre volume} \times \text{Normality of alkali} \times \text{volume made up} \times \text{Equivalent weight of acid} \times 100}{\text{Volume of aliquot sample} \times \text{weight of sample} \times 1000}$$

### Result and Discussion

Plant height, petiole length, number of leaves and plant spread all showed in the data (Table 1). Result on different PGR (NAA, GA3, BA) combination indicated that T9{BA(6-benzyladenine) 75ppm} recorded maximum plant height (cm) of 12.56cm (30DAT), 13.22cm (60 DAT), 14.16cm (90 DAT) whereas minimum plant height (cm) of 5.95cm (30DAT), 7.54cm (60 DAT), 9.12cm (90 DAT) was recorded in T0 (Control), recorded maximum number of leaves per plant of 4.80 (30DAT), 12.93 (60DAT), 16.33 (90DAT) whereas minimum number of leaves per plant of 2.67 (30DAT), 7.56 (60DAT), 10.54 (90DAT) was recorded in T0 (Control). recorded maximum plant spread of 16.26cm (30DAT), 21.52cm (60DAT), 32.54cm (90DAT) whereas minimum plant spread of 6.54cm (30DAT), 10.86cm (60DAT), 18.56cm (90DAT) was recorded

inT0(Control). recorded maximum petiole length (cm) of 8.50cm (30DAT), 10.00cm (60DAT), 12.04cm (90DAT) whereas minimum petiole length of 3.31cm (30DAT), 4.72cm (60DAT), 5.98cm (90DAT) was recorded in T0(Control). “When no of runners found best in treatment T9. The number of fruits, flowers, and fruit productivity all significantly increased after BA application, according to their observations of plant height, leaf count, leaf area, and fruit productivity. The evidence of, who also discovered that the application of BA greatly improves plant height, leaf area, fruit setting percentage, and the number of runners is consistent with the study's findings as well”. (Qureshi *et al.*, 2013)

**Table 1: Impact of plant growth regulators on Plant Height (cm), Number of leaves per plant, plant spread (cm), petiole length (cm) at (30,60,90 DAT)**

Notation	Plant height (cm)			Number of leaves			Plant spread (cm)			Petiole length (cm)		
	30 Days	60 Days	90 Days	30 Days	60 Days	90 Days	30 Days	60 Days	90 Days	30 Days	60 Days	90 Days
T0	5.95	7.54	9.12	2.67	7.56	10.54	6.54	10.86	18.56	3.31	4.72	5.98
T1	8.09	9.92	11.83	3.80	10.00	12.60	11.88	18.28	28.86	4.36	6.57	8.79
T2	9.06	9.20	10.99	3.20	11.40	13.40	12.28	17.54	28.84	4.77	6.74	8.89
T3	9.30	10.19	11.22	3.86	10.33	12.33	13.06	17.82	29.12	4.29	5.61	7.77
T4	10.31	10.20	12.04	3.66	11.20	13.93	14.15	19.13	30.20	4.80	6.20	8.36
T5	9.27	10.41	12.31	3.77	10.53	13.20	14.03	18.45	29.00	5.16	6.49	7.66
T6	7.43	11.40	11.41	3.73	10.80	14.00	11.94	17.14	27.63	4.57	5.98	8.02
T7	8.40	9.25	13.05	3.93	11.06	13.80	13.12	17.96	28.98	6.17	7.37	8.39
T8	7.60	11.75	13.43	4.00	11.53	14.13	11.80	19.31	26.69	7.52	9.38	11.54
T9	12.56	13.22	14.16	4.80	12.93	16.33	16.26	21.52	32.54	8.50	10.00	12.04
Sed	0.23	0.20	0.20	0.24	0.45	0.62	0.33	0.47	0.833	0.38	0.37	0.37
CD	0.69	0.61	0.59	0.50	0.95	1.31	0.71	1.01	1.76	0.81	0.79	0.79
F-Test	S	S	S	S	S	S	S	S	S	S	S	S

Days to first flowering, number of flowers per plant, number of fruits per plant, diameters of fruit, length of fruit, fruit weight, fruit set (%) all showed in the data (Table 2). Result on different PGR (NNA, GA3, BA) combination indicated that T9{BA(6-benzyladenine) 75ppm} recorded maximum days to first flowering 62.91 whereas minimum days to first flowering 48.66 recorded in T0 (control), and recorded maximum number of flowers per plant of 1.60 (60DAT), 4.96 (70DAT), 13.45 (80DAT) whereas minimum number of flowers per plant of 0.87 (60DAT), 2.83 (70DAT), 4.76 (80DAT) was recorded in T0(Control). recorded maximum number of fruits per plant of 1.26 (75DAT), 4.47 (85DAT), 7.13 (95DAT) whereas minimum number of fruits per plant of 0.54 (75DAT), 2.30 (85DAT), 4.32 (95DAT) was recorded in T0(Control), and recorded maximum diameters of fruit (cm) 3.98 cm whereas minimum diameters of fruits 2.97 cm recorded in T0 (control), and recorded maximum length of fruit 60.66 mm whereas minimum length of fruits 37.10 mm recorded in T0 (control), and recorded maximum fruit weight 31.55 g whereas minimum fruit weight 11.13 g recorded in T0 (control). “In foliar spray with BA, additional biomass may be able to generate extra metabolites during photosynthesis, which eventually sank into the producing fruits and generated berry with the most weight. In strawberries, the use of BA has been observed to boost berry weight” (Sharma and Singh, 2009). “A higher number of

marketable fruits were produced as a result of the exogenous application of BA, which also indirectly affected the benzylidene metabolism and increased the fruit yield. It has also been previously documented that applying BA increases fruit yield in strawberry”(**Rathod *et al.*, 2021**).

**Table2:Impact of plant growth regulators on Days of first flowering, Number of flowers per plant at 60 days,70 days,80 days, Number of fruits per plant at 75 days,85 days,95 days, Diameters of fruit (cm), Length of fruit (mm), Fruits weight (g), Fruit set (%).**

Notation	Days to first flowering	Number of flower/plants			Number of fruits/plants			Diameters of fruit	Length of fruit	Fruit weight	Fruit set
	Days	60 Days	70 Days	80 Days	75 Days	85 Days	95 Days	(cm)	(mm)	(g)	(%)
<b>T0</b>	62.91	0.87	2.83	4.76	0.54	2.30	4.32	2.97	37.10	11.13	63.80
<b>T1</b>	58.08	1.13	4.33	6.80	0.86	3.93	5.87	3.20	38.60	16.55	63.83
<b>T2</b>	56.58	1.20	4.20	6.46	1.06	3.73	5.93	3.23	38.80	17.63	66.33
<b>T3</b>	55.25	1.46	4.53	6.93	1.00	3.80	6.13	3.30	40.80	19.22	77.93
<b>T4</b>	55.16	1.26	4.80	7.13	0.73	4.13	6.60	3.26	41.50	19.79	80.86
<b>T5</b>	54.83	1.33	4.39	9.85	0.66	3.66	6.73	3.39	51.50	20.48	75
<b>T6</b>	55.25	1.40	4.66	10.45	1.13	4.21	6.20	3.53	52.33	20.73	81.66
<b>T7</b>	52.72	1.35	4.85	11.32	0.98	3.76	5.45	3.76	51.83	24.39	76.66
<b>T8</b>	50.50	1.54	4.79	11.67	1.20	4.36	6.80	3.81	54.50	28.41	82
<b>T9</b>	48.66	1.60	4.96	13.45	1.26	4.47	7.13	3.98	60.66	31.55	83.10
<b>Sed</b>	0.61	0.22	0.35	1.87	0.22	0.36	0.41	0.30	1.12	0.38	5.04
<b>CD</b>	0.82	---	1.46	3.35	----	1.34	1.57	0.91	2.37	1.14	10.6
<b>F-Test</b>	S	NS	S	S	NS	S	S	S	S	S	S

Number of fruits per plant, TSS (Total Solid Soluble), Acidity (%), Ascorbic acid(**Table 3**). Result on different PGR (NNA, GA3, BA) combination indicated that T9{BA(6-benzyladenine) 75ppm} recorded maximum Number of fruits per plant16.66 whereas minimum number of fruits per plant8.66 recorded in T0 (control), and recordedmaximumTSS (Total Solid Soluble)of10.17%whereas minimum TSS (Total Solid Soluble)of5.33%was recorded inT0(Control), and recordedmaximumacidity of 1.33% whereas minimum 0.60%was recorded inT0(Control), and recorded maximum ascorbic acid % (mg/100g) 58.66% whereas minimum ascorbic acid % (mg/100g) 51.30%recorded in T0 (control),When minimum result show in control T0. BA (6-benzyledenin) was applied to fruits, which massively improved total soluble solids and decreased titratable acidity. With foliar treatment of 75 ppm BA, the highest total soluble solid was recorded. However, in the current investigation, the control group had the lowest levels of total soluble solids. These results support those of **Prasad *et al.* (2013)**, who demonstrated that BA concentrations responded favourably to a strawberry quality measure.

**Table 3:Impact of plant growth regulators on Number of fruits per plant, TSS, Acidity (%), Ascorbic acid % (mg/100g).**

	Number of fruits per plant	TSS (Total solid soluble)	Acidity	Ascorbic acid
Notation	No.	(%)	(%)	%(mm/100g)
<b>T0</b>	8.66	5.33	0.6	51.30
<b>T1</b>	11	8.33	0.6	53.83
<b>T2</b>	11.33	8.33	0.83	53.83
<b>T3</b>	13	8.50	0.90	54.50
<b>T4</b>	14	7.67	0.86	54.83
<b>T5</b>	11	7.83	1.03	56.16
<b>T6</b>	12	8.50	0.86	56.83
<b>T7</b>	12.66	9.50	0.80	56
<b>T8</b>	14	9.50	1.06	57.16
<b>T9</b>	16.66	10.17	1.33	58.66
<b>Sed</b>	0.90	0.63	0.19	0.56
<b>CD</b>	1.89	1.32	NA	1.19
<b>F-Test</b>	S	S	S	S

## Conclusion

From the results obtained during the present investigation with different treatment of NAA, GA3 and {BA6-benzyladenine) 75ppm} on vegetative growth, flowering, yield and quality of strawberry (fragaria x ananassa duch) cv. (winter down), it is concluded that plants treated with Treatment T9 {BA6-benzyladenine) 75ppm} significantly increased the height of plant days to first flower, Days to first fruiting, fruit set, number of fruits per plant and fruit yield per plant. So far as the yield, the response of strawberry plants treated with T9 {BA6-benzyladenine) 75ppm} plants produced higher yield. TSS of T9 {BA6-benzyladenine) 75ppm} was higher as compared to other treatments. Total Acidity treated with {BA6-benzyladenine) 75ppm} were higher than the other treatments. Ascorbic acid titratable was recorded in plants treated with T9 {BA6-benzyladenine) 75ppm}. Treatment T9 recorded as the best treatment of yield and quality of strawberry. The B: C ratio was highest in the treatment T<sub>9</sub>{BA6-benzyladenine) 75ppm}with (4.45).

## Disclaimer (Artificial intelligence)

### Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

### Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the

generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

## REFERENCES

1. **Archana, B. and Patil, V. S., (2013)**, Studies on ethyl methane sulphonate (EMS) induced mutations for enhancing variability of gladiolus varieties (*Gladiolus hybridus* Hort.) in M1V2 generation. *Karnataka Journal of Agriculture Sciences*, 26 (3): 403-407.
2. **Anitha, K., Jegadeeswari, V. and Selvaraj, N., (2017)**, Studies on genetic variability, heritability and genetic advance in lisianthus (*Eustoma grandiflorum L.*). *Agriculture Update*, 12(2):
3. **Balamurugan, Rengasamy, P. and Arumugam, T. (2002)**. Variability studies in gladiolus. *Journal of Ornamental Horticulture*, 5(1): 5 38-39
4. **Balaram, M. V. and Janakiram, T. (2009)**. Correlation and path coefficient analysis in gladiolus. *Journal of ornamental horticulture*, 12(1): 22-29.
5. **Bhatia, R. and Grewal, H. P. S. (2009)**. Genetic variability and heritability studies in gladiolus. *Indian Agriculturist*, 53(2): 1-6.
6. **V.K. Tripathi, Sanjeev, K. and Gupta, A.K.** Influence of NAA and vermicompost on growth, flowering, yield and quality of strawberry cv. Chandler Indian J. Hort., 72(2): 201-205(2015).
7. **Wange, S.S., Patil, M.T. and Singh, B.R. 1998.** Cultivar x biofertilizer interaction study in strawberry. *Recent Hort.* 4: 43-49.
8. **Yadav, S.K., Prasad, R. and Khokhar, U.U. 2010.** Optimization of integrated nutrient supply system for strawberry cv. Chandler in H.P. (India). *Scientia Hort.* 124(1):62-6.
9. **Zargar MY, Baba ZA, Sofi PA.** Effect of N, P and biofertilizers on yield and physico-chemical attributes of strawberry. *Agro Thesis*. 2008; 6(1):3-8.
10. **Singh, M., Gogoi, N., Singh, P., Singh, H., & Srivastava, A.K. (2015).** Phosphate-solubilizing bacteria enhance the growth and yield but not the phosphorus uptake of wheat grown in phosphorus-deficient soils. *Biology and Fertility of Soils*, 51(6), 723-731.
11. **Staudt, G. 1962.** Taxonomic studies in the genus *Fragaria* typification of *Fragaria* species known at the time of Linnaeus. *Canad. J. Bot.* 40(6): 869-886.
12. **Tripathi V.K., Kumar. N.; Shukla, H.S. and Mishra, A.N. 2010.** Influence of NAA, Azospirillum and GA3 on growth, yield and quality of strawberry cv. Chandler. In: National Symposium on Conservation Horticulture held at Dehradun during March, 21-23, 2010. pp-198-199.
13. **Elkoca, E., Kantar, F. & Sahin, F. (2008).** Influence of Nitrogen Fixing & Phosphorus Solubilizing Bacteria on the Nodulation, Plant Growth, and Yield of Chickpea. *J. Of P. Nutr.* 31: 157-171.
14. **El-Zeiny, O.A.H., El-Behairy, U.A. and Zaky, M.H. (2001),** Influence of bio-fertilizer on growth, yield and fruit quality of tomato grown under plastic house. *J. Agricult. Sci. - Mansoura University*, 26 (3): 1749-1763.
15. **Gajbhiye, R.P., Sharma, R. R. and Tewari R.N. 2003.** Effect of bio-fertilizers on growth and yield parameters of tomato. *Indian J. Hort.* 60 (4):368-371.
16. **Gogoi D, Kotoky U, Hazarika S.** Effect of biofertilizers on productivity and soil characteristics in banana. *Indian J Hort.* 2004;61(4):354-356.

17. Gore NS, Sreenivasa MN. Influence of liquid organic manures on growth, nutrient content and yield of tomato (*Lycopersicon esculentum* Mill.) in the sterilized soil. *Karnataka Journal of Agricultural Sciences*. 2011; 24:153-156.
  18. Hammam, M.S. 2003. Effect of bio-fertilizers on growth and fruiting of Cavendish and Williams banana. *Egyptian J. Hort.* 30 (1-2): 67-81.
  19. Iqbal U, Wali VK, Kher R, Jamwal M. Effect of FYM, urea and NAA on growth, yield and quality of strawberry cv. Chandler. *Notulae Not. Bot. Hort. Agrobot. Cluj*. 2009;37(1):139-143.
  20. Joshi PS. Effect of growing media, bioregulators and nutrients on growth, yield and quality of strawberry cv. Chandler. Ph.D. Thesis. Dr YSP Parmar University of Horticulture and Forestry, Solan, HP, 2003, 130.
  21. Kumar, S., Kundu, M., Das, A., Rakshit, R., Siddiqui, Md, W. and Rani, R. (2019), Substitution of mineral fertilizers with biofertilizer: an alternate to improve the growth, yield and functional biochemical properties of strawberry (*Fragaria × ananassa* Duch.) cv. Camarosa.
- 
- 22 Singh S, Bahadur V, Mishra S. Effect of Different Spacing and NPK Combination on Plant Growth, Fruit Yield and Fruit Quality of Strawberry (*Fragaria ananassa* Duch.) cv. Winter Dawn. *J. Adv. Biol. Biotechnol.* [Internet]. 2024 May 16 [cited 2024 Jun. 24];27(6):415-27. Available from: <https://journaljabb.com/index.php/JABB/article/view/900>
- 
- 23 Khatoon F, Kundu M, Mir H, Nahakpam S. Efficacy of foliar feeding of brassinosteroid to improve growth, yield and fruit quality of strawberry (*Fragaria × ananassa* Duch.) grown under subtropical plain. *Communications in Soil Science and Plant Analysis*. 2021 Apr 28;52(8):803-14.
- 
- 24 Paikra S. INFLUENCE OF NAA AND GA3 ON GROWTH, FLOWERING, YIELD AND QUALITY OF STRAWBERRY (*Fragaria x ananassa* Duch.) CV. SABRINA UNDER NET TUNNEL (Doctoral dissertation, Indira Gandhi Krishi Vishwavidyalaya, Raipur).
- 
25. Yadav I, Singh J, Meena B, Singh P, Meena S, Neware S, Patidar DK. Strawberry yield and yield attributes after application of plant growth regulators and micronutrients on cv. Winter Dawn. *Chemical Science Review and Letters*. 2017;6(21):589-94.