

Effect of Biochar Application on Biochemical and Physiological Processes in Wheat (*Triticum aestivum* L.) Under Water Stress Condition

Abstract

A field experiment was conducted at Udaipur (Rajasthan) during rabi season of 2021-22 to evaluate the effect of water stress and biochar application on biochemical and physiological processes in wheat. The experiment consisted of four levels of water stress as main-plot and four levels of biochar as sub-plot treatments conducted in split plot design (SPD) replicated thrice. Results revealed that water stress at various growth stages and biochar application had effect on proline, chlorophyll and relative water content (RWC). Water stress at grain filling stages resulted in higher chlorophyll and relative water content as compared to no water stress. Further, water stress at tillering stages resulted in higher proline over no water stress. Application of biochar had no effect on proline and chlorophyll contents. Further, application of Biochar @ 4 t/ha in significant relative water content.

Key words: Biochar, water stress, wheat, physiological and biochemical

Introduction

Wheat (*Triticum aestivum* L.) is the world's major cereal crop known as "king of Cereals". It is locally known as Gehun, belongs to the Poaceae family. It contains about 8-15% protein, 1-1.5% fat, 2-2.5% fibre, and 62-71% carbohydrate and supplies 73% of the calories of the average diet. It is consumed in the form of chapatti and its straw is used for feeding the animals (Heyne, 1987). In India wheat is grown in 30.47 M ha area with production of 106.84 MT and productivity 3507 kg ha⁻¹, while in Rajasthan it is cultivated in 2.58 M ha area with production and productivity of 9.48 M T and 3673 kg ha⁻¹, respectively (DAC&FW, 2021-22). Currently water scarcity has become the leading menace to curtail crop productivity around the globe (Hussain *et al.*, 2008). Water stress is major harmful factor in arid and semi-arid regions worldwide that limits the area under cultivation and yield of crops. Drought is observed in irrigated areas due to insufficient supply of water and canal closure. Recently, biochar is getting importance worldwide to improve the water holding capacity and physiochemical properties of soil [14-17].

Biochar is a carbon-rich co-product resulting from pyrolyzing biomass. When applied to the soil it resists decomposition, effectively sequestering the applied carbon and mitigating anthropogenic CO₂ emissions. Zhang *et al.* (2012) reported that application of biochar to calcareous and infertile dry croplands poor in soil organic carbon enhanced crop productivity and reduced GHGs emissions. Other benefits of biochar application to soil included enhanced

plant productivity and reduced nutrient leaching. Biochar soil amendment can affect leaf N status and photosynthesis, but the effect varied with soil type (Xu *et al.*, 2015). Biochar is preferred due to its unique properties of low density (providing additional void age and aeration in the soil), significant adsorption and cation exchange capacity, and the ability to promote living microbiology in the soil.

Materials and Methods

The field experiment was conducted at Rajasthan College of Agriculture, MPUAT, Udaipur during rabi season of 2021-22. Treatments comprised of sixteen treatment combinations consisting of four water stress *viz.*, No water stress (W₁), water stress at tillering (W₂), water stress at flowering (W₃) and water stress at grain filling stage (W₄) as main plot and four biochar levels (0, 2, 3 and 4 t/ha) as sub plot were tested in SPD with three replications. In water stress treatments, water stress was imposed at tillering, flowering and grain filling stage of the crop as per treatments, whereas in no water stress six irrigations were provided at critical growth stages of wheat. Biochar was prepared at the Department of Renewable Energy Engineering, College of Technology and Engineering, Udaipur from the twigs of trees, weeds and agricultural waste. Biochar was applied as per treatments and mixed well in soil prior to sowing of the crop. Recommended dose of fertilizers *i.e.*, 90 kg N, 60 kg P₂O₅ and 40 kg K₂O / ha were applied through commercial fertilizers *viz.*, urea, DAP and MOP. Estimation of proline, chlorophyll and relative water content (RWC) in leaves.

Chart 1: Physiological/Biochemical process

S.No.	Physiological/Biochemical process	References
1.	Proline	Bates <i>et al.</i> (1973)
2.	Chlorophyll	SPAD
3.	Relative water content (RWC)	Barrs and Weatherley (1992)

Data were analysed statistically for analysis of variance (ANOVA). Treatments were compared by computing the 'F-test'. The significant differences between treatments were compared by critical difference at 5% level of probability (Panse and Sukhatme, 1989).

Results and discussion

Proline content at 10 days after water stress

A perusal of the data (Table 1) revealed that water stress caused significant variation

in proline contents in leaves. At tillering 10 days after water stress, significantly higher proline content was recorded under water stress ($8.59\mu\text{g g}^{-1}$) over control ($7.33\mu\text{g g}^{-1}$), water stress at flowering ($7.48\mu\text{g g}^{-1}$) and grain filling ($7.41\mu\text{g g}^{-1}$). Whereas, significantly higher proline content was recorded at 10 days after water stress at flowering under treatment water stress at flowering over control ($8.48\mu\text{g g}^{-1}$), water stress at tillering ($8.82\mu\text{g g}^{-1}$) and water stress at grain filling ($8.62\mu\text{g g}^{-1}$). At 10 days after water stress at grain filling, water stress at grain filling treatment ($9.39\mu\text{g g}^{-1}$) recorded significantly higher proline content over rest of the treatments. These results are conformity with the result obtain by Hafez *et al.*, (2021) and Chowdhury *et al.*, (2024). Application of biochar failed to register any significant variation in proline content at all the three stages of proline estimation.

Chlorophyll content at 10 days after water stress

Water stress treatments brought about significant variation in chlorophyll content (SPAD) at 10 days after water stress at tillering and flowering stages Whereas, water stress treatments didn't alter SPAD reading for chlorophyll at 10 days after grain filling stage of crop over control. At 10 days after water stress at tillering and flowering, significantly higher SPAD reading was recorded for chlorophyll under water stress treatments over no water stress. All the water stress treatments were at par with each other in respect of SPAD reading for chlorophyll content in leaves at all the three stages. Application of biochar didn't significantly influence chlorophyll content in leaves at 10 days after water stress at tillering, flowering and grain filling over control (Table 1).

Relative water content at 10 days after water stress

At 10 days after water stress at tillering, relative water content (RWC) in leaves significantly reduced under treatment water stress at tillering over control, water stress at flowering and grain filling. Whereas, at 10 days after water stress at flowering, significant reduction in relative water content was recorded in treatment water stress flowering over no water stress, tillering and grain filling stage. Significant decrease in RWC was recorded under water stress at grain filling at 10 days after water stress at grain filling stage over rest of the treatments. Under the moisture stress condition in soil, less water is available to plant for absorption which reduced both turgidity of cell and relative water content. The relative water content and transpiration intensity of leaves decreased. Similar results were recorded by Gupta and Gupta (2011), Nezhedahmedi *et al.* (2013) and Meena (2015) in wheat crop. Application of biochar at all three stages significantly increased RWC in leaves at 10 days

after water stress at tillering, flowering and grain filling stages over control. Further, application of biochar 2 t ha⁻¹, 3 t ha⁻¹ and 4 t ha⁻¹ were at par with each other in respect of relative water content at all the three stages (Table 1). The data showed that under no water stress, the addition of biochar at 2 t ha⁻¹, 3 t ha⁻¹, and 4 t ha⁻¹ significantly increased relative water content (RWC) at 10 days after flowering compared to no water stress. However, only biochar at 3 t ha⁻¹ was superior to the control. Biochar at 2 t ha⁻¹ was superior under water stress at tillering. Under water stress at flowering, biochar at 2 t ha⁻¹ was significantly superior to the control. Biochar at 4 t ha⁻¹ also significantly increased RWC at 10 days after flowering compared to biochar at 3 t ha⁻¹ under water stress at grain filling.

Conclusion

The study concludes that water stress significantly increased proline content in wheat leaves at different growth stages, while biochar application did not alter proline content. Additionally, water stress reduced chlorophyll content and relative water content (RWC), whereas biochar application mitigated these effects, enhancing RWC uniformly across different application rates.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

1. Barrs, H.D. and Weatherly, P.E. (1992) A re-examination of the relative turgidity techniques for estimating water deficit in leaves. *Australian Journal of Biological Science* **15**: 413-428.
2. Bates, L.S., Waldren, R.P. and Teare, I.D. (1973) Rapid determination of free proline for water stress studies. *Plant and Soil* **39**(1): 205-207.
3. Chowdhury, M.S.N., Sani, M.N.H., Siddique, A.B., Hossain, M.S. and Yong, J.W.H. (2024) Synergistic effects of biochar and potassium co-application on growth, physiological attributes, and antioxidant defense mechanisms of wheat under water deficit conditions. *Plant Stress* **12**:100452.
4. DAC & FW. (2022) Agricultural statistics at a glance. Website: <http://eands.dacnet.nic.in> retrieved on dated **02-10-2022**.
5. Gupta, S. and Gupta, N.K. (2011) Field efficacy of exogenously applied putrescine in wheat (*Triticum aestivum* L.) under water-stress conditions. *Indian Journal of Agricultural Sciences* **81**(6): 516-9.
6. Hafez, E.M., Omara, A.E.D., Alhumaydhi, F.A. and El-Esawi, M.A. (2021) Minimizing hazard impacts of soil salinity and water stress on wheat plants by soil application of vermicompost and biochar. *Physiologia Plantarum* **172**(2): 587-602.
7. Heyne, E. G. (1987). *Wheat and wheat improvement* (Vol. 2). American Society of Agronomy, Crop Science Society of America, Soil Science Society of America.
8. Hussain, M., Malik, M.A., Farooq, M., Ashraf, M.Y. and Cheema, M.A. (2008) Improving drought tolerance by exogenous application of glycine betaine and salicylic acid in sunflower. *Journal of Agronomy and Crop Science* **194**: 193- 199.
9. Meena, Pinky. (2015) Effect of putrescine on growth and productivity of wheat (*Triticum aestivum* L.) under water stress condition. M.Sc. Thesis, Maharana Pratap University of Agriculture and Technology, Udaipur.
10. Nezhadahmadi, A., Prodhan, Z.H. and Faruq, G. (2013) Drought tolerance in wheat. *The Scientific World Journal* **2013**: 610721.
11. Panse, V.G. and Sukhatme, P.V. (1989) Statistical methods for agricultural workers. Indian Council of Agricultural Research, New Delhi.
12. Xu, C., Hosseini-Bai, S. and Hao, Y. (2015) Effect of biochar amendment on yield and photosynthesis of peanut on two types of soils. *Environmental Science and Pollution Research* **22**: 6112-6125.

13. Zhang, A., Liu, Y., Pan, G., Hussain, Q., Li, L., Zheng, J. and Zhang, X. (2012) Effect of biochar amendment on maize yield and greenhouse gas emissions from a soil organic carbon poor calcareous loamy soil from central china plain. *Plant and Soil* **351** (1):263-275.
14. Afrad, Md. Safiul Islam, G. K. M. Mustafizur Rahman, Mohammad Saiful Alam, Md. Zulfiker Ali, and Aliyu Akilu Barau. 2021. "Effects of Organic and Inorganic Fertilizers on Growth and Yield of Different Crops at Charlands in Bangladesh". *Asian Journal of Advances in Agricultural Research* **17** (3):27-40. <https://doi.org/10.9734/ajaar/2021/v17i330198>.
15. Hassan , Reem H. I., Abbas S. M. T., and Ismail A. Y. 2023. "Influence of Biochar and Irrigation Levels on Productivity of Marjoram (*Origanum Majorana* L.) under Sandy Soil Conditions". *Asian Journal of Agricultural and Horticultural Research* **10** (4):538-58. <https://doi.org/10.9734/ajahr/2023/v10i4293>.
16. Oliveira FR, Patel AK, Jaisi DP, Adhikari S, Lu H, Khanal SK. Environmental application of biochar: Current status and perspectives. *Bioresource technology*. 2017 Dec 1;246:110-22.
17. Atkinson CJ, Fitzgerald JD, Hipps NA. Potential mechanisms for achieving agricultural benefits from biochar application to temperate soils: a review. *Plant and soil*. 2010 Dec;337:1-8.

Table: 1 Effect of Water Stress and Biochar on Proline, Chlorophyll and Relative Water Content (RWC) in Leaves of Wheat

Treatments	Proline content ($\mu\text{g g}^{-1}$)			Chlorophyll content (SPAD)			RWC (%)		
	Water stress at tillering (after 10 days)	Water stress at flowering (after 10 days)	Water stress at grain filling (after 10 days)	Water stress at tillering (after 10 days)	Water stress at flowering (after 10 days)	Water stress at grain filling (after 10 days)	Water stress at tillering (after 10 days)	Water stress at flowering (after 10 days)	Water stress at grain filling (after 10 days)
Water stress									
W ₀	7.33	8.48	8.90	53.81	54.03	55.66	54.37	56.91	59.50
W ₁	8.59	8.82	8.96	56.16	56.35	54.90	51.76	56.12	58.66
W ₂	7.48	9.42	9.00	55.93	56.12	54.67	54.24	53.53	58.96
W ₃	7.41	8.62	9.39	56.22	56.42	54.96	54.62	56.93	55.87
SEm \pm	0.11	0.11	0.05	0.50	0.52	0.50	0.49	0.62	0.60
CD (P=0.05)	0.39	0.38	0.17	1.73	1.81	NS	1.68	2.13	2.07
Biochar application									
Control	7.83	8.96	9.08	55.49	55.69	55.07	52.53	53.96	56.70
2 t ha ⁻¹	7.75	8.88	9.08	55.68	55.88	55.18	53.89	56.24	57.84
3 t ha ⁻¹	7.69	8.82	9.02	55.46	55.66	54.96	54.11	56.46	59.09
4 t ha ⁻¹	7.54	8.67	9.05	55.49	55.69	54.99	54.46	56.84	59.35
SEm \pm	0.08	0.08	0.04	0.49	0.50	0.49	0.47	0.60	0.58
CD (P=0.05)	NS	NS	NS	NS	NS	NS	1.36	1.76	1.69
Interaction (W \times B)									
SEm \pm	0.16	0.16	0.07	0.99	0.99	0.97	0.94	1.21	1.16
CD (P=0.05)	NS	NS	NS	NS	NS	NS	NS	3.52	NS

*W₀: Control (no water stress) W₁: Water stress at tillering W₂: Water stress at flowering W₃: Water stress at grain filling