

## Original Research Article

### Eco- friendly management of root rot disease in Cowpea caused by *Macrophomina phaseolina*

#### ABSTRACT

The research was conducted on evaluation of biological agents and organic amendments against *Macrophomina phaseolina* of cowpea at The Department of Plant Pathology, S. D. Agricultural University Sardarkrushinagar, Gujarat during 2018-2019. Seven known antagonists were tested *in vitro* for their antagonism to *M. phaseolina* using dual culture method. *T. harzianum* (Sardarkrushinagar) and *T. viride* (Sardarkrushinagar) were potent antagonists of *M. phaseolina*. The six organic amendments at three different concentrations were tested against *M. phaseolina* by poisoned food technique *in vitro*. Significantly least mycelium growth of *M. phaseolina* was recorded in the extract of neem cake followed by mustard cake. Vermicompost and Farm yard manure were less effective against *M. phaseolina*.

**KEYWORDS:** *M. phaseolina*, antagonists, organic amendments, *in vitro*

#### INTRODUCTION

Cowpea [*Vigna unguiculata* (L.) Walp.] is locally known as chowli, lobiya, southern pea or black eye pea in India (Singh, 1998). Tropical Africa is a native of cowpea. It is grown extensively in West Africa, Brazil and India (Stele and Mehra, 1980). It has been referred to as “poor man’s meat” because of its high protein content (20 - 25%). It contains large amount of quality protein (23.4%), carbohydrate (60.3%), fat (1.8%) and sufficient amount of calcium (76.00 mg/100 g), iron (57 mg/100 g) and vitamins such as thiamine (0.92 mg/100 g), riboflavin (0.18 mg/100 g) and nicotinic acid (1.9 mg/100 g) (Chatterjee and Bhattacharya, 1986).

Cowpea is infected by many diseases caused by viruses, bacteria and fungi. The root rot caused by *Macrophomina phaseolina* (Tassi.) Goid. causes significant loss in yield. Concurrent heat and moisture stress favor development of charcoal or dry root rot disease makes cultivation of cowpea uneconomical (Singh *et al.*, 2012). Incidence of this disease ranging from 5 - 39 per cent (Ushamaliniet *al.*, 1997). *M. phaseolina* (Tassi.) Goid. is a soil borne plant pathogen with a very wide host range. *M. phaseolina* (Tassi.) Goid. Attacks a oilseeds, legumes and vegetable

crops. Its micro-sclerotia are formed in senescing shoot tissues, survive well in soil. The fungus poses great problem in managing the disease.

The hazardous effect of chemicals used in plant disease management has diverted plant pathologists to find alternative methods having little or no adverse effect on the environment. Nowadays the commercial formulations of some of the biological agents are already made available in the market for farmers use. However, inadequate information on the performance of the antagonists under varying condition is a major constraint in the large scale adoption of this technology. Management of plant pathogen through botanicals, organic amendments, antagonist micro-organism is the alternative way of managing the plant pathogens without use of chemicals. Its good for sustainable disease management.

## **MATERIAL AND METHODS**

### **Isolation, purification and identification of pathogen**

The diseased samples of cowpea showing typical root rot symptoms were collected from Pulses Research Station, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar as well as farmer's field. The infected diseased samples were brought to the laboratory for microscopic examination and tissue isolation. The culture obtained was purified by hyphal tip method/single spore isolation technique (Rangaswami, 1972). The obtained culture was maintained on PDA slants for further investigations. The pure culture was maintained by periodical transfer on PDA slants. The pure cultures of the isolates were maintained on PDA slants in refrigerator at  $5 \pm 2^\circ\text{C}$  temperature for further study.

Identification of the pathogen causing root rot of cowpea grown on PDA medium was examined visually as well as microscopically for cultural and morphological characters viz., mycelial growth, color and shape of sclerotia were recorded and compared with the available literature viz., 'Illustrated Genera of Imperfect Fungi' by Barnett and Hunter (1972) as well as description given by Ashby (1927) and Goidanich (1947). The cultural characters were recorded right from initiation of growth (24 hrs) up to 15 days. While morphological characters were measured under high power magnification from 7 days old culture of *Macrophomina* sp. The microphotograph of micro-sclerotia was also taken.

### **Efficacy of different biological control agents against pathogen *in vitro***

The effects of different antagonists (Table 1) were studied by dual culture technique against *M.phaseolina*. The seven antagonistic microorganisms employed for the *in vitro* antimicrobial

assay obtained from the Department of Plant Pathology, Chimanbhai Patel College of Agriculture, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar. The antagonistic fungal cultures were maintained on PDA culture medium and bacterial culture was maintained on nutrient agar media in 90 mm diameter sterilized Petri-plates.

**Table 1: List of bio-control agents tested against *Macrophomina phaseolina* in vitro**

Sr. No.	Treatment	Name of the bio-control agents and location of isolation
1	T <sub>1</sub>	<i>Trichoderma viride</i> (Navsari isolate)
2	T <sub>2</sub>	<i>Trichoderma harzianum</i> (Sardarkrushinagar isolate)
3	T <sub>3</sub>	<i>Trichoderma viride</i> (Junagadh isolate)
4	T <sub>4</sub>	<i>Trichoderma harzianum</i> (Junagadh isolate)
5	T <sub>5</sub>	<i>Trichoderma harzianum</i> (Anand isolate)
6	T <sub>6</sub>	<i>Pseudomonas fluorescens</i> (Sardarkrushinagar isolate)
7	T <sub>7</sub>	<i>Trichoderma viride</i> (Sardarkrushinagar isolate)
8	T <sub>8</sub>	Control (Test pathogen only)

The assay for antagonism was performed on PDA medium on Petri plates by the dual culture method. Mycelial disc (5 mm diameter) from seven days old actively growing culture of bio-agents and the test pathogen were cut aseptically from the periphery of the colony with the help of sterilized cork borer and placed on solidified PDA approximately 60 mm away from each other. To test for antagonistic bacteria, a 5 mm of mycelial disc of pathogen cultures was placed on the one side of a Petri plate containing PDA medium. A loopful of bacteria was then streaked 60 mm away from the disc of *Macrophomina* isolate on the same plate. Culture plates were incubated at 28 ± 2°C. The plates inoculated only with test pathogens served as control. The experiment was conducted with three repetitions of each treatment.

#### **Efficacy of different organic amendments against pathogen in vitro**

Effects of different organic amendments (Table 2) were studied at three different concentrations (5, 10 and 20 %) on the growth of *M. phaseolina* under *in vitro* condition. All the amendments as mentioned in Table 2 were crushed to make fine powder. Fifty gram powder of each amendment was taken into 250 ml flask and then 150 ml sterilized water added to the flask. All these flasks were plugged with cotton and allowed to decompose the material for 15 days. After 15 days, the material was strained with muslin cloth to obtain the extract. The strained liquid was autoclaved at 1.045 kg/cm<sup>2</sup> ( 15 psi) pressure and 121° C temperature for 20 minutes and considered as cent per cent (100 %) concentration (standard solution). The measured

quantity of standard solution of the organic amendments were incorporated separately in melted potato dextrose agar (PDA) medium in conical flasks aseptically at the time of pouring the medium to obtain desired concentration. The medium was shaken well to give uniform dispersal and then poured about 20 ml in each sterilized Petri plates. After solidification of the medium, the Petri plates were inoculated in the centre by placing seven days old mycelial discs (5 mm) of the pathogen and then incubated in incubator at  $28 \pm 2^\circ\text{C}$  temperature. A control was also maintained by growing the pathogen on organic amendment free medium.

**Table 2: List of different organic amendments tested against *M. phaseolina in vitro***

Sr. No.	Treatment	Organic amendment	Concentrations (%)		
			5	10	20
1	T <sub>1</sub>	Mustard cake	5	10	20
2	T <sub>2</sub>	Neem cake	5	10	20
3	T <sub>3</sub>	Castor cake	5	10	20
4	T <sub>4</sub>	Cotton cake	5	10	20
5	T <sub>5</sub>	Farm yard manure (FYM)	5	10	20
6	T <sub>6</sub>	Vermicompost	5	10	20
7	T <sub>7</sub>	Control	-	-	-

### Observations recorded

Observations on the radial growth (mm) were recorded from 24 h of incubation at  $28 \pm 2^\circ\text{C}$  till the complete growth of test pathogen achieved in control plates. The per cent growth inhibition (PGI) over control was calculated by using following formula given by Vincent (1947).

$$\text{PGI} = \frac{C - T}{C} \times 100$$

Where,

- PGI = Per cent growth inhibition,
- C = Colony in diameter in control (mm), and
- T = Colony in diameter in treatment (mm).

## RESULTS AND DISCUSSION


### Identification of pathogen

The studies on the morphological and cultural characters of isolated *Macrophomina* sp. showed its close identity with *Macrophomina phaseolina* (Tassi) Goid as described by Grover

and Sakhuja (1981). The results obtained by Agarwal (1993) and Tandel (2004) also similar with our present finding. Thus, the causal organism of root rot of cowpea was confirmed as *Macrophomina phaseolina* (Tassi) Goid.


### Cultural and morphological characters

The mycelium of the fungus was initially white, gradually turned brown to black in colour due to formation of numerous small black sclerotia. In a course of mycelial development, which grew fast on potato dextrose agar (PDA) covering the entire Petri plate (90 mm) surface within 5 days at  $28 \pm 2^\circ\text{C}$  temperature (Plate 1). The fungus produced initially white mycelial growth on PDA later changing to brown black centered due to formation of numerous small black sclerotia. The mycelium was hyaline to brown, branched, septate and dendroid (Plate 2A). The sclerotia formed in culture were black and hard (Plates 2 B ). The pycnidia were not produced in culture. The morphology of *M. phaseolina* causing dry root rot of green gram was studied by Agrawal (1993).



### Efficacy of different bio-agents against pathogen *in vitro*

The results are presented in Table 3 revealed that, all the antagonists tested against *M. phaseolina* were effective in checking the growth of *M. phaseolina* (Plate 3). Out of seven antagonists tested, least growth of the pathogen was recorded in *T. harzianum* (Sardarkrushinagar) (27.93 mm) which was followed by *T. viride* (Sardarkrushinagar) (31.81 mm), *T. viride* (Navsari) (34.18 mm), *T. harzianum*(Junagadh) (38.54 mm), *T. viride*(Junagadh) (40.87 mm), *T. harzianum* (Anand ) (42.31 mm) and *P. fluorescence* (43.49 mm).



Treatment	
T <sub>1</sub>	= <i>Trichodemaviride</i> (Navsari isolate)
T <sub>2</sub>	= <i>Trichodermaharzianum</i> (Sardarkrushinagar isolate)
T <sub>3</sub>	= <i>Trichodemaviride</i> (Junagadh isolate)
T <sub>4</sub>	= <i>Trichodermaharzianum</i> (Junagadh isolate)
T <sub>5</sub>	= <i>Trichodermaharzianum</i> (Anand isolate)
T <sub>6</sub>	= <i>Pseudomonas fluorescense</i> (Sardarkrushinagar isolate)
T <sub>7</sub>	= <i>Trichodemaviride</i> (Sardarkrushinagar isolate)
T <sub>8</sub>	= Control (Test pathogen only)

Plate 3 : Efficacy of different bio-control agents against *M. phaseolina* in vitro

The per cent growth inhibition was computed in Table 3 it revealed that *T. harzianum* (Sardarkrushinagar) (68.90 %), *T. viride* (Sardarkrushinagar) (64.6 %) and *T. viride* (Navsari) (62.34 %) significantly inhibited the growth of the pathogen. Whereas, *T. harzianum* (Junagadh) (56.76 %), *T. viride* (Junagadh) (54.53 %), *T. harzianum* (Anand) (52.62 %) and *P. fluorescense* (Sardarkrushinagar) (44.87 %) were comparatively found less effective.

It is evident from the study that among all the antagonists evaluated by dual culture method, *T. harzianum* (Sardarkrushinagar), *T. viride* (Sardarkrushinagar) and *T. viride* (Navsari) consistently showed strong antagonistic properties against *M. phaseolina* as compared to the other antagonists tested hence considered as potential antagonists.

Table 3: Efficacy of different bio-control agents against *M. phaseolina* in vitro

Tr. No.	Bio-agents	Colony diameter of pathogen (mm)	Growth inhibition over control (%)
T <sub>1</sub>	<i>Trichodemaviride</i> (Navsari isolate)	34.18	52.15 <sup>abc</sup> (62.34)
T <sub>2</sub>	<i>Trichoderma harzianum</i> (Sardarkrushinagar isolate)	27.93	56.11 <sup>a</sup> (68.90)
T <sub>3</sub>	<i>Trichodemaviride</i> (Junagadh isolate)	40.87	47.60 <sup>cd</sup> (54.53)
T <sub>4</sub>	<i>Trichodermaharzianum</i> (Junagadh isolate)	38.54	48.89 <sup>bcd</sup> (56.76)
T <sub>5</sub>	<i>Trichodermaharzianum</i> (Anand isolate)	42.31	46.52 <sup>de</sup> (52.65)
T <sub>6</sub>	<i>Pseudomonas fluorescense</i> (Sardarkrushinagar isolate)	43.49	42.06 <sup>e</sup> (44.87)
T <sub>7</sub>	<i>Trichodemaviride</i> (Sardarkrushinagar isolate)	31.81	53.49 <sup>ab</sup> (64.6)
T <sub>8</sub>	Control	90.00	4.05 <sup>f</sup> (0.49)
S.Em.±			0.647


C.D. at 5%	1.956
C.V. %	2.584

Figures in parentheses are retransformed values of arc sine transformation, Treatment means with the letter(s) in common are not significant by DNMRT at 5% level of significance.

These results are in harmony with earlier workers Thombre and Kohire (2018) reported *T. harzianum* most effective and recorded significantly highest mycelial growth inhibition (77.59 %) of the *M. phaseolina* over untreated control. Kumar *et al.* (2020) also reported that maximum growth inhibition of *M. phaseolina* causing dry root rot of mungbean was recorded in *T. harzianum* (73.33 %) followed by *T. viride* (64.44 %) *in vitro* conditions.

### Efficacy of different organic amendments against pathogen *in vitro*

The aqueous extracts of six different organic amendments were evaluated for their inhibitory effect to *M. phaseolina*. The results presented in Table 4 and (Plate 4) indicated that organic extracts produced significant inhibitory effect on the fungal growth.



Treatment                      Concentration  
T<sub>1</sub> = Mustard cake              C<sub>1</sub> = 5%

T <sub>2</sub>	=	Neem cake	C <sub>2</sub>	=	10%
T <sub>3</sub>	=	Castor cake	C <sub>3</sub>	=	20%
T <sub>4</sub>	=	Cotton cake			
T <sub>5</sub>	=	Farm yard manure			
T <sub>6</sub>	=	Vermicompost			
T <sub>7</sub>	=	Control			

**Plate 4: Efficacy of different organic amendments against *M. phaseolina* in vitro**

Maximum per cent growth inhibition of *M. phaseolina* was recorded in neem cake at the rate of 20 % (59.54 %) which was at par with mustard cake at the rate of 20 % (56.57%), cotton cake at the rate of 20 % (56.38 %) and neem cake at the rate of 10 % (56.0 %). Whereas, vermicompost at the rate of 5 % (33.08 %) and Farm yard manure at the rate of 5 % (33.96 %) were less effective in inhibiting the growth of the pathogen .

**Table 4: Efficacy of different organic amendments against *M. phaseolina* in vitro**

Tr. No.	Organic amendments	Growth inhibition (%)			Mean
		Concentration (%)			
		5	10	20	
T <sub>1</sub>	Mustard cake	43.14 <sup>cd</sup> (46.75)	45.90 <sup>b</sup> (51.57)	48.78 <sup>a</sup> (56.57)	45.94 <sup>b</sup> (51.63)
T <sub>2</sub>	Neem cake	43.46 <sup>cd</sup> (47.31)	48.45 <sup>a</sup> (56.00)	50.50 <sup>a</sup> (59.54)	47.47 <sup>a</sup> (54.30)
T <sub>3</sub>	Castor cake	35.88 <sup>e</sup> (34.35)	36.66 <sup>e</sup> (35.64)	43.78 <sup>cd</sup> (47.87)	38.77 <sup>d</sup> (39.21)
T <sub>4</sub>	Cotton cake	42.07 <sup>d</sup> (44.89)	45.16 <sup>bc</sup> (50.27)	48.67 <sup>a</sup> (56.38)	45.30 <sup>b</sup> (50.52)
T <sub>5</sub>	Farm yard manure	35.65 <sup>e</sup> (33.96)	41.95 <sup>d</sup> (44.68)	43.46 <sup>cd</sup> (47.31)	40.35 <sup>c</sup> (41.98)
T <sub>6</sub>	Vermicompost	33.27 <sup>f</sup> (33.08)	34.98 <sup>ef</sup> (36.41)	36.77 <sup>c</sup> (40.03)	35.01 <sup>e</sup> (36.47)
T <sub>7</sub>	Control	4.05 <sup>g</sup> (0.49)	4.05 <sup>g</sup> (0.49)	4.05 <sup>g</sup> (0.49)	4.05 <sup>f</sup> (0.49)
Mean		34.04 <sup>b</sup> (31.33)	38.01 <sup>a</sup> (37.92)	38.04 <sup>a</sup> (37.97)	-
		Organic amendments	Concentration		Organic amendments × Concentration
S.Em. ±		0.349	0.228		0.604

C.D. at 5 %	0.995	0.652	1.24
C.V. %	2.85		

Figures in parentheses are retransformed values of arc sine transformation, Treatment means with the letter(s) in common are not significant by DNMRT at 5% level of significance.

From this study, it is clear that neem cake was found effective in reducing the growth of *M. phaseolina* causing root rot in cowpea. The present investigation is quite similar to the work done by earlier workers Makwana (2010) who concluded that neem cake at the rate of 10 per cent with maximum (53.69 %) growth inhibition against *M. phaseolina*. Dhingani *et al.* (2013) recorded the least growth of *M. phaseolina* mycelium was found in extracts of neem cake (59.40%) followed by farm yard manure (42.56%). Meena *et al.* (2014) also concluded that maximum mycelial growth inhibition (52.40 %) was recorded in neem cake (*Azadirachta indica*) at the concentration of 20 per cent followed by 42.61 and 29.60 per cent with concentrations of 15 and 10 per cent, respectively.

### Conclusion

Microscopic examination and tissue isolation from stem and root of infected cowpea plant yielded culture of *M. phaseolina*. Seven known antagonists were tested *in vitro* for their antagonism with *Macrophomina phaseolina* by dual culture method. Among them, *Trichoderma harzianum* (Sardarkrushinagar) and *Trichoderma viride* (Sardarkrushinagar) were potent antagonists of *M. phaseolina*.

The six organic amendments at three different concentrations *viz.*, 5, 10 and 20 per cent were tested against *M. phaseolina* by poisoned food technique *in vitro*. All the amendments were significantly inhibitory to *M. phaseolina*. Maximum per cent growth inhibition of pathogen was recorded by the extract of neem cake (54.30%) followed by and mustard cake (51.63%).

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name,

version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

### References:

1. Agrawal, S.C. (1993). "Disease of green gram and black gram" international book distributors. pp. 1-2, 5, 28, 32-33, 159, 181, 269.
2. Ashby, S.F. (1927). *Macrophominaphaseolina* (Maubl.) Comb. Nov. the pycnidial stage of *Rhizoctoniabataticola* (Taub.) Butl. *Transactions of the British Mycological Society*. **12** : 141-147.
3. Barnett, H.L. and Hunter, B.B. (1972). *Innervated Genera of Imperfect Fungi*, Burgess Publishing Company, Minneapolis, Minn, U.S.A.
4. Chatterjee, B.N. and Bhattacharya, K.K. (1986). *Principle and practices of grain legume production*. Oxford and IBH Publishing Co. Ltd., New Delhi.
5. Dhingani, J.C., Solanky, K. C. and Kansara S.S (2013). Management of root rot diseases [*Macrophomina phaseolina* (Tassi.) G. Goid] of chickpea through botanicals and oil cakes. *The Bioscan An International Quarterly Journal Of Life Sciences*. **8**(3): 739-742.
6. Goidanich, G. (1947). Revisione del genera *Macrophomina* Petrak. Species tipica : *Macrophomina phaseolina* (Tassi.) G. Goid. n. comb. nov. *M. phaseoli* (Maubl.) Ashby. *Annali della sperimentazione agrarian Roma N.S.* **1** : 449-461.
7. Grover, R.K. and Sakhuja, P.K (1981). Some pathological studies on *Rhizoctonia bataticola* leaf blight of mungbean *Indian Phytopathology*. **34** (1) : 24-29
8. Kumar, M.; Ram Kumhar, D., Lal Godara, S., & Garg, S. (2020). Evaluation the efficacy of plant extracts and bioagents against *Macrophomina phaseolina* *in vitro*. *International Journal of Current Microbiology and Applied Sciences*, **9** (5), 3474-3481.
9. Makwana, J.R. (2010). Investigations on charcoal rot disease [*Macrophominaphaseolina* (Tassi.) Goid.] of Soybean [*Glycine max* (L.) Merrill.] Under South Gujarat conditions. **M.Sc. (Agri.)** Thesis submitted to Navsari Agricultural University, Navsari.
10. Meena, P.N.; Tripathi, A.N.; Gotyal, B.S. and Satpathy, S. (2014). Bio-efficacy of phytoextracts and oil cakes on *Macrophomina phaseolina* (Tassi) Goid. causing stem rot disease of jute, *Corchorus* spp. *Journal of Applied Natural Science*. **6** (2) : 530-533.
11. Rangaswami, G. (1972). *Diseases of crop plants in India*. Prentice Hall of India Pvt. Ltd., New Delhi. p. 520.
12. Singh, R.S. (1998). *Plant Diseases* (Seventh Edition). Oxford and IBH Publication, New Delhi. p. 686.
13. Singh, R.; Maurya, S. and Upadhyay, R.S. (2012). Antifungal potential of *Trichoderma* species against *Macrophominaphaseolina*. *Journal of Agricultural Technology*. **8** : 1925-1933.

14. Stele, W.M. and Mehra, K.L. (1980). Structure, evolution and adaptation to farming systems and environments in *Vigna*. In : Advances in legume science (Eds. R.J. Summerfield and A.H. Bunting), HMSO, London, pp. 393-404.
15. Tandel D. H. (2004). Epidemiology and management of leaf blight of green gram (*Phaseolus aureus* Roxb.) caused by *Macrophominaphaseolina*(Tassi). Goid. M.Sc. (Agri.) Thesis (Unpublished) submitted to Navsari Agricultural University, Navsari.
16. Tassi, A. (1901). Novae Micromycetum species Ser HI Bull Lab Orto Bot Univ Sura. **4** : 7-12.
17. Thombre, B.B. and Kohire, O.D. (2018). Integrated management of *Macrophomina* blight of mungbean (*Vigna radiata* L.) caused by *Macrophomina phaseolina* (Tassi.) Goid. *Indian Phytopathology*.**71** : 423-429.
18. Ushamalini, C.; Rajappan, K. and Gangadharan, K. (1997). Management of charcoal rot of cowpea using biocontrol agents and plant products. *Indian Phytopathology*.**50** (4) : 504-507.
19. Vincent, J.M. (1947). Distortion of fungal hyphae in presence of certain inhibitors *Nature*. **159** : 850-856.
- 20. Chauhan , Sreyansh Singh, Ajinkya A. Telgote, Gaurav Gunjan, Saket Mishra, Shashi Kant Ekka, Johnson Lakra, and Reena Kujur. 2024. "A Review on Effects of Biofertilizers and Bio Capsules on Vegetative Growth, Flowering and Fruiting Attributes of Strawberry". Asian Journal of Advances in Agricultural Research 24 (1):33-38. <https://doi.org/10.9734/ajaar/2024/v24i1485>.**
- 21. Ishak, Mariam H., Marzouk R. Abdel-Latif, Harby M. Abd Alla, and Mohamed M. N. Shaat. 2020. "Thyme (Thymus Vulgaris L.) Root Rot/Wilt in Minia Governorate, Egypt". Asian Journal of Agricultural and Horticultural Research 5 (4):44-57. <https://doi.org/10.9734/ajahr/2020/v5i430060>.**
- 22. <https://www.sciencedirect.com/science/article/abs/pii/S0167198706001681>**
- 23. <https://link.springer.com/article/10.1007/s10327-018-0792-5>**