

## Review Article

# Arbuscular Mycorrhizal Fungi Efficiency on Plant Growth and Nutrient Acquisition: A Comprehensive Review

### Abstract

Arbuscular Mycorrhizal Fungi (AMF) play a crucial role in enhancing plant growth and nutrient acquisition. The impacts of pressures on crop yield have been made worse by climate change and agricultural practices such as overuse of pesticides and fertilizers, which have also damaged the environment. AMF are one environmentally friendly management strategy that is desperately needed to increase agricultural output. More importantly, it's widely acknowledged that the AMF inoculation confers resistance in host plants to a range of stressful environments, such as heat, salt, drought, metals, and abnormally high or low temperatures. This review paper provides a comprehensive analysis of the current understanding of AMF efficiency on various aspects of plant physiology, including nutrient uptake and overall plant development. The paper synthesizes information from recent studies to present a nuanced perspective on the multifaceted interactions between AMF and plants. Furthermore, challenges and future directions in AMF research are discussed, highlighting the need for a more integrated and holistic approach to harness the full potential of these fungi for enhancing plant productivity and nutrient acquisition.

**Keywords:** Arbuscular Mycorrhizal Fungi (AMF), plant growth, nutrient acquisition, plant tolerance

### Introduction


The growing concern about food security has brought plant science to light as a developing field which has driven researchers to come up with fresh ideas for raising crop yields. Food shortage is caused by both biotic and abiotic factors that restrict global agricultural yield, such as salt, drought, floods, plant diseases, nutritional deficiencies, and toxicity. It's critical to appropriately regulate and make use of good microbial activities in order to reduce adverse environmental consequences and accomplish sustainable goals (Laishram *et al.*, 2023). Arbuscular Mycorrhizal Fungi (AMF) form symbiotic relationships with the majority of plant species, influencing plant

growth and nutrient uptake. These beneficial fungi form mutualistic symbiotic associations with the roots of most plant species, creating a vast network of interconnected hyphae that extends the root system and greatly increases the surface area available for nutrient absorption (Aryal & Xu, 2001). This symbiotic relationship benefits both the fungi and the plants (Bücking *et al.*, 2012). The fungi receive carbohydrates from the plants, while the plants receive increased access to nutrients such as phosphorus and nitrogen (Huey *et al.*, 2020). Furthermore, arbuscular mycorrhizal fungi have been shown to improve plant tolerance to various abiotic stresses, including drought, salinity, and heavy metal toxicity.

Many studies have focused on the efficiency of arbuscular mycorrhizal fungi in enhancing plant growth and nutrient uptake (Aerts, 2003). These studies have consistently demonstrated the positive effects of arbuscular mycorrhizal fungi on plant growth and nutrient acquisition (Begum *et al.*, 2019). In addition to promoting plant growth and nutrient uptake, arbuscular mycorrhizal fungi also play a role in protecting plants against pathogens and toxic stresses (Huey *et al.*, 2020). Research has demonstrated that arbuscular mycorrhizal fungi can act as bioprotectants, helping plants defend against pathogens and tolerate toxic stresses. For example, Copetta *et al.* (2007) found that arbuscular mycorrhizal fungi enhanced the growth and essential oil production in *Ocimum basilicum* plants. These findings highlight the potential of harnessing the symbiotic relationship between arbuscular mycorrhizal fungi and plants for sustainable agriculture and plant health management practices. The research on arbuscular mycorrhizal fungi efficiency on plant growth and nutrient uptake has demonstrated consistent positive effects. Arbuscular mycorrhizal fungi have been shown to improve plant tolerance to various abiotic stresses, including drought, salinity, and heavy metal toxicity (Mitra *et al.*, 2019). In the case of metal toxicity, several studies have indicated that arbuscular mycorrhizal fungi can improve plant tolerance and reduce the detrimental effects of heavy metals on plant growth. Furthermore, arbuscular mycorrhizal fungi have been found to enhance nutrient acquisition by plants (Chen *et al.*, 2017). They are particularly effective at increasing the availability of phosphorus to plants, which is often a limiting nutrient in many soils.

In the scientific literature, arbuscular mycorrhizal fungi are generally acknowledged for their effectiveness in promoting plant growth and nutrient absorption. Arbuscular mycorrhizal fungi have been demonstrated to improve soil structure and stability, increase organic matter decomposition, and enhance nutrient uptake in addition to their direct effects on plant growth and nutrient uptake (Begum *et al.*, 2019) (Figure 1). Arbuscular mycorrhizal fungi also have

indirect effects on ecosystem functioning and soil fertility. In light of the significance of AMF and the developments in research about its applications in agriculture, this review centers on the function of AMF in the control of plant growth and development with enhanced nutrient absorption in demanding conditions.



**Fig. 1: A schematic depiction of how mycorrhizal fungi control several ecological processes and stimulate plant growth and nutrient uptake in stress conditions**

### **Mycorrhiza evolution and types:**

Mycorrhizae are ancient, maybe even older than the terrestrialization of plants, according to fossil and genetic data. Research reveals that proto-mycorrhizal fungi were a critical feature enabling plant terrestrialization. Genetic evidence implies that all land plants share a single common ancestor (Harris et al., 202), who appears to have swiftly acquired mycorrhizal symbiosis (Puginier et al., 2022). An assemblage of fossil plants from the 400 million-year-old Rhynie chert have been sufficiently preserved to allow for the observation of arbuscular mycorrhizae in the stems of *Aglaophyton major*, providing a lower constraint on the possible development time of late mycorrhizal symbiosis (Remy et al., 1994) While the majority of

contemporary mycorrhizal groups, such as orchid and erchoid mycorrhizae, date to the Cretaceous era's angiosperm radiation, ectomycorrhizae evolved far later, during the Jurassic period (Miyachi et al., 2020). Genetic research suggests that legume-fixing bacterium symbiosis is a genetic outgrowth of mycorrhizal symbiosis (Provorov et al., 2016). The distribution of mycorrhizal fungi now seems to be a reflection of the rivalry and complexity in root architecture that arose during the Cenozoic Era, when angiosperm dominance was the norm and complex ecological interactions existed between species (Brundrett et al., 2018)

An ectomycorrhiza and an endomycorrhiza are two categories for mycorrhizas. Hyphae of endomycorrhizal fungus infiltrate the cell wall and invade the cell membrane, but ectomycorrhizal fungi do not penetrate individual cells inside the root. This distinguishes the two types of fungi (Harley & Smith, 1983; Allen, 1991). While arbutoid mycorrhizas fall under the category of ectoendomycorrhizas, endomycorrhizas comprise arbuscular, ericoid, and orchid mycorrhizas. One distinct class of mycorrhizas is monotropoid.

#### **Mycorrhizal mutualist mechanisms:**

Most plant species have a mutualistic interaction with mycorrhizal fungi in their roots. Both the plants and the portions of the roots that house the fungus are referred to be mycorrhizal in such a connection. While only a small percentage of mycorrhizal relationships between plant species and fungi have been studied thus far, 95% of plant families that have been studied are primarily mycorrhizal, either because the majority of their species have a beneficial association with mycorrhizae or because they are completely dependent on mycorrhizae. It is well known that in the Orchidaceae family, seeds will not even germinate if the proper mycorrhizae are not present (Trappe, 1987). Mycorrhizal fungi and plants may have a more complicated interaction than only mutualism, according to recent studies on ectomycorrhizal plants in boreal forests. When it was surprisingly discovered that mycorrhizal fungi were storing nitrogen from plant roots during periods of nitrogen shortage, this link was recognized. According to research, certain mycorrhizae disperse nutrients in response to the surrounding plants and other mycorrhizae. They go on to describe how this revised model may account for the fact that mycorrhizae do not mitigate plant nitrogen restriction and that as soil nitrogen supply decreases, plants may suddenly transition from a mixed strategy including both mycorrhizal and nonmycorrhizal roots to a solely mycorrhizal approach (Franklin et al., 2014). Furthermore, a greater degree of variation in the

intensity of mycorrhizal mutualisms may be explained by evolutionary and phylogenetic links than by ecological variables (Hoeksema et al., 2018). Give an explanation of the mycorrhizal formation process, emphasizing the important phases of fungal colonization in plant roots.

### **Impact of AMF on Plant Nutrient Acquisition**

#### **Plant growth enhancement:**

Beneficial rhizosphere bacteria have the ability to increase crops' quality as well as their nutritional condition. Over the past 20 years, several studies on AMF have demonstrated its numerous advantages for crop yield and soil health. As a result, it is generally accepted that AMF may eventually be used in place of inorganic fertilizers due to the fact that mycorrhizal treatment may significantly lower the amount of chemical fertilizer input, particularly phosphorus (Ortas, 2012). As naturally occurring root symbionts, AMF help host plants by supplying vital inorganic nutrients, which enhances growth and production in both stressed and unstressed environments. AMF's ability to function as a biofertilizer may improve plants' capacity to adapt to shifting environmental conditions. Arbuscular mycorrhizal fungi (AMF) may form symbiotic associations with the majority of terrestrial plants (Smith et al., 2008). Plant growth and yield, nutrient acquisition (Qiao et al., 2015; Ercoli et al., 2017), especially P and other poorly mobile nutrients (Siasou et al., 2009), enhanced resistance to biotic stresses (Fileccia et al., 2017) and abiotic stresses (salinity, drought, etc.) (Rillig & Mummey, 2006) and enhanced soil aggregation and carbon sequestration are all impacted by this type of symbiosis. AMF can also change the microbial community's structure and promote the growth of other microorganisms in the rhizosphere (Secilia & Bagyaraj, 1987; Andrade, 2004), which can affect a number of biological processes, including biological N fixation (Püschel et al., 2017).

The strawberry that was colonized by AMF, for instance, showed higher levels of secondary metabolites, which enhanced its antioxidant properties (Castellanos-Morales et al., 2010). According to Hart et al. (2015), AMF can improve the nutritional value of crops by influencing the synthesis of carotenoids and other volatile chemicals. AMF has positive benefits on tomato quality, as noted by Bona et al. (2017). Citrus fruit quality was improved in another study by Zeng et al. (2014), which found that *Glomus versiforme* boosted the amounts of sugars, organic acids, vitamin C, flavonoids, and minerals. Stronger anthocyanin, chlorophyll, carotenoids, tocopherols, total soluble phenolics, and different mineral elements accumulate when

mycorrhizal symbiosis is present (Baslam et al., 2011). AMF have proven to have a significant potential for increasing crop productivity since they have been used in large-scale field production of potatoes (Hijri, 2016), yam (Lu et al., 2015), and maize (Sabia et al., 2015). According to Sbrana et al. (2014) and Rouphael et al. (2015), AMF can also improve the manufacture of beneficial phytochemicals in edible plants, enabling them to function properly in the food production chain. According to Rouphael et al. (2015), maintaining the pH of the soil can help AMF mitigate abiotic stress and preserve its horticultural value. As explained below, AMF may also be quite helpful in enhancing plants' ability to withstand harsh conditions.

### **Nutrient cycling and uptake:**

Many studies have demonstrated that excessive land use may have a significant influence on biodiversity overall, which can then have an impact on how well an ecosystem functions (Balliu et al., 2015; Wagg et al., 2015). Transferring resources, such as organic carbon (C), in the form of lipids and sugars, is one important function of such symbiotic relationships (Jiang et al., 2017). It is well accepted that AMF colonization increases plants' absorption of nutrients. It is clear that inoculating with AMF may greatly raise the concentration of different macro- and micronutrients, which increases the synthesis of photosynthate and, in turn, the accumulation of biomass (Chen et al., 2017; Mitra et al., 2019). AMF can increase nearly all plants' absorption of inorganic nutrients, particularly phosphate (Nell et al., 2010). Additionally, AMF are highly successful in facilitating plants' uptake of nutrients from nutrient-deficient soils (Kayama and Yamanaka, 2014). AMF interaction has been shown to enhance the phyto-availability of micronutrients such as copper and zinc in addition to macronutrients (Smith and Read, 1997). AMF increases the host roots' capacity to absorb surface energy (Bisleski, 1973). Increased leaf area and higher levels of nitrogen, potassium, calcium, and phosphorus have been seen in experimental experiments conducted on tomato plants infected with AMF, indicating greater plant development (Balliu et al., 2015). AMF have a symbiotic relationship with roots in order to get vital nutrients from the host plant and then give back mineral nutrients including N, P, K, Ca, Zn, and S. As a result, AMF give the plants nutritional assistance even in unsuitable root cell environments.

AMF create fungal structures called arbuscules that aid in the interchange of inorganic minerals and carbon and phosphorus molecules, giving host plants a significant boost in vitality (Prasad et

al., 2017). As a result, they can greatly increase the concentration of phosphorus in both the root and shoot systems (Al-Hmoud and Al-Momany, 2017). Mycorrhizal association enhances phosphorus delivery to the infected roots of host plants in phosphorus-limited environments (Bucher, 2007). For instance, the AMF-colonized maize plants showed a significant improvement in Pi absorption rate (Garcés-Ruiz, 2017). Enhanced growth frequency of AMF inoculation is directly correlated with increased photosynthetic activity and other leaf functions. This is because enhanced growth frequency of AMF inoculation is directly correlated with the intake of N, P, and carbon, which travel towards roots and stimulate tuber development. It has been noted that under various irrigation regimes, AMF maintains P and N absorption, eventually aiding in plant growth at higher and lower P levels (Liu et al., 2018). For instance, in *Pelargonium graveolens* L. under drought stress, mycorrhizal symbiosis favorably raised the concentrations of N, P, and Fe (Amiri et al., 2017). Gomez-Bellot et al. (2015) observed that rapid fungal attachment led to increased P, Ca, and K levels in *Euonymus japonica* under salinity stress.

AMF is thought to increase the absorption of nearly all necessary nutrients while decreasing the absorption of Na and Cl, which stimulates growth (Evelin et al., 2012). Plant growth and development can be enhanced by the extra-radical mycelium (ERM) through efficient nutrient absorption improvement (Lehmann and Rillig, 2015). Even in places with an adequate amount of livestock and farm-yard manure (FYM), nitrogen (N), a well-known mineral fertilizer, is a major source of soil nourishment. Numerous researchers have documented how AMF aids in the absorption of soil nutrients, particularly N and P, which can successfully stimulate the development of host plants (Smith et al., 2011). N is a key element restricting development in certain crops and higher plants. AMF can absorb and transport N to neighboring plants or host plants, according to a number of studies (Turrini et al., 2018). Zhang et al. (2018) have shown that, especially under low fertilizer levels, AMF mediated enhanced shoot biomass allocation to panicles and grains through increased N and P redistribution to panicles.

According to Zhu et al. (2016), AMF inoculation enhances C and N accumulation as well as N assimilation at both ambient and increased CO<sub>2</sub> concentrations. According to Bati et al. (2015), AMF, for instance, has been shown to enhance the development, accumulation of micro- and macronutrients, and distribution of these nutrients in plantlets cultivated in environments with

elevated Mn levels in olive plants. The important dynamic characteristics that contribute to improving the advantageous effects of AMF colonization on overall plant performance are the improvement of plant nutrition and the preservation of the  $\text{Ca}^{2+}$  and  $\text{Na}^{+}$  ratio (Evelin et al., 2012). Mycorrhizal chickpea was shown to have improved growth and levels of protein, iron, and zinc (Pellegrino and Bedini, 2014). Furthermore, mycorrhizal symbiosis's contribution to different micronutrients in crops was demonstrated by two metaanalysis papers that surfaced a few years ago (Lehmann et al., 2014). According to Asrar et al. (2012), while *Antirrhinum majus* was experiencing a drought, the specific fungal relationship increased the amount of macronutrients such N, P, K, Ca, and Mg. AMF also demonstrated efficacy in limiting the elevated build-up of Na, Mn, Mg, and Fe in roots (Bati et al., 2015). According to a number of research carried out in recent years, AMF, including *Glomus mosseae* and *Rhizophagus irregularis*, demonstrated enhanced heavy metal translocation in the shoot (Zaefarian et al., 2013). Mycorrhizal hyphae assist plants in absorbing micronutrients like zinc and copper, which have restricted dispersion in soil.

### **Challenges and Future Directions:**

#### **Limitations of AMF Application:**

In modern times, pesticide usage is a crucial component of contemporary agriculture that depends on technology. Since most high-yielding crops are more prone to disease than their wild counterparts, there is a rising demand for them. High concentrations of key fertilizers, particularly nitrogen and phosphate; pesticides and fungicides; extensive tillage; and crop rotation with non-mycorrhizal crops inhibit AM activity, variety, and association. According to Mbutia et al. (2015), the variety, population, and root colonization of AM flora in agricultural fields are therefore changed and poorer than in nearby natural soil. Mycorrhizal symbiosis is observed to be inhibited by high P concentrations in plants brought on by high P fertilization in soil (Balzergue et al., 2013). Plants may absorb sufficient phosphorus without sacrificing carbohydrates when high P fertilizer is applied (García-Caparrós et al., 2021). In return for phosphate, AM fungus wants a carbon source from plants. Cooperation from both parties is necessary for AM to exist and function, and the plant itself selects the most efficient and compatible strains by sharing more resources with them (Kiers et al., 2011). Numerous nitrogen fertilizers have been shown in pot and field trials to reduce colonization (Getman-Pickering et

al., 2021). Plant growth and root development are increased by low to medium levels of AM colonization and sporulation. Applying nitrogen fertilizer at a higher rate inhibits AM colonization in plants (Lin et al., 2020). Greater than ideal potassium concentration inhibits AM signaling, reduces root exudation, and accumulates soluble carbohydrates in the cortex.

Despite the fact that these agrochemicals generally have detrimental impacts on the environment, soil, and human health, crop productivity is never compromised. To control leaf, seed, and soil-borne pathogens, fungicides, both systemic and non-systemic, are employed; however, the majority of them negatively impact AM spore germination, colonization, extraradical hyphal growth, sporulation (Buysens et al., 2015), and the efficiency of P uptake by phosphatase activity (Channabasava et al., 2015). By inhibiting the processes of nutrient transfer, especially in dry soil lacking in nutrients, both fungicides hinder the growth of crops that are dependent on AM.

The hyphal network in the soil is destroyed by soil disturbances and tillage, which can physically crush AM spores, reducing root colonization (Brito et al., 2012). The amount of soil that AM can remove is reduced when colonized root fragments and hyphal networks are disrupted. The current favorable circumstances for AMF species are also altered by the movement of the soil layer. The physical disruption of the extra radical mycelium network during the early colonization stage is directly linked to the direct effects of conventional tillage systems on AM activity in nutrient and water uptake, glomalin-related soil aggregate formation, and bioprotection against soil pathogens (Brito et al., 2012). Propagule density, activity, and AMF community are all negatively impacted by extended periods of fallow land and crop rotation including non-mycorrhizal crops. According to Higo et al. (2015), the fallow time and the addition of non-host species have an impact on the density and quantity of AMF propagules, which has a negative effect on the production of subsequent crops. According to Valerino et al. (2014), waterlogged soil in paddy agriculture prevents AM activity since AM cannot develop in wet soil. Considering that conventional agrochemical-based agriculture disrupts the symbiosis and effectiveness of AM, the advantages of AM remain untapped in this context.

## **Conclusion**

The main discussion surrounding AMF has been its ability to help soil organisms absorb nutrients. Nevertheless, recent research has demonstrated that plants injected with AMF can successfully fend off a variety of environmental cues, such as salinity, drought, nutrient stress,

alkali stress, cold stress, and extreme temperatures, helping to increase the yield of numerous crops and vegetables per hectare. For contemporary global agricultural systems to remain consistently sustainable, it is crucial to promote the use of AMFs. Without a doubt, using AMF for agricultural enhancement may cut down on the usage of pesticides and synthetic fertilizers considerably, supporting bio-healthy farming practices. Crop plants with AMF-mediated growth and productivity increase may be able to meet the world's growing population's consumption needs. Additionally, because of their widespread use, environmentally friendly technology should be strongly supported.

### **Future Research Directions:**

Future research should primarily concentrate on identifying the genes and gene products regulating the AMF-mediated regulation of growth and development in response to stressful stimuli. Future studies in this area should focus on identifying the key cellular and metabolic pathways under varying environmental conditions, as well as the host and AMF specific protein components governing symbiotic interaction. Crop production may be increased by comprehending the crosstalk that is activated to control plant performance and the AMF-induced modifications in the tolerance mechanisms. All things considered, AMF need to be thoroughly investigated to learn more about their function in nature for sustainable agricultural output. To effectively define methods for managing the soil/plant system to enhance this significant ecological process, a greater comprehension of the significance of each route involved in AMF-mediated nutrient cycling and transfer is necessary. Novel and inventive research methodologies will be needed for this.

### **References**

1. Aerts, R. (2003). The role of various types of mycorrhizal fungi in nutrient cycling and plant competition. In *Ecological studies* (pp. 117–133). [https://doi.org/10.1007/978-3-540-38364-2\\_5](https://doi.org/10.1007/978-3-540-38364-2_5).
2. Al-Hmoud, G., and Al-Momany, A. (2017). Effect of four mycorrhizal products on squash plant growth and its effect on physiological plant elements. *Adv. Crop. Sci. Tech.* 5, 260. doi: 10.4172/2329-8863.1000260.
3. Allen, Michael F. 1991. *The ecology of mycorrhizae*. Cambridge University Press, Cambridge.
4. Amiri, R., Ali, N., Nematollah, E., and Mohammad, R. S. (2017). Nutritional status, essential oil changes and water-use efficiency of rose geranium in response to arbuscular

- mycorrhizal fungi and water deficiency stress. *Symbiosis* 73, 15–25. doi: 10.1007/s13199-016-0466-z.
5. Andrade G. Role of functional groups of microorganisms on the rhizosphere microcosm dynamics In: Varma A, Abbott L, Werner D, Hampp R, editors. *Plant Surface Microbiology*. Berlin: Springer; 2004. pp. 51–69.
  6. Aryal, U. K., & Xu, H. (2001). Mycorrhizal associations and their manipulation for Long-Term Agricultural Stability and Productivity. *Journal of Crop Production*, 3(1), 285–302. [https://doi.org/10.1300/j144v03n01\\_24](https://doi.org/10.1300/j144v03n01_24).
  7. Asrar, A. A., Abdel-Fattah, G. M., and Elhindi, K. M. (2012). Improving growth, flower yield, and water relations of snapdragon *Antirrhinum majus* L. plants grown under well-watered and water-stress conditions using arbuscular mycorrhizal fungi. *Photosynthetica* 50, 305–316. doi: 10.1007/s11099-012-0024-8.
  8. Balliu, A., Sallaku, G., and Rewald, B. (2015). AMF Inoculation enhances growth and improves the nutrient uptake rates of transplanted, salt-stressed tomato seedlings. *Sustainability* 7, 15967–15981. doi: 10.3390/su71215799.
  9. Balzergue, C., Chabaud, M., Barker, D. G., Bécard, G., & Rochange, S. (2013). High phosphate reduces host ability to develop arbuscular mycorrhizal symbiosis without affecting root calcium spiking responses to the fungus. *Frontiers in Plant Science*, 4. <https://doi.org/10.3389/fpls.2013.00426>.
  10. Baslam, M., Garmendia, I., and Goicoechea, N. (2011). Arbuscular mycorrhizal fungi (AMF) improved growth and nutritional quality of greenhouse grown lettuce. *J. Agric. Food Chem.* 59, 5504–C5515. doi: 10.1021/jf200501c.
  11. Bati, C. B., Santilli, E., and Lombardo, L. (2015). Effect of arbuscular mycorrhizal fungi on growth and on micronutrient and macronutrient uptake and allocation in olive plantlets growing under high total Mn levels. *Mycorrhiza* 25 (2), 97–108. doi: 10.1007/s00572-014-0589-0.
  12. Begum, N., Qin, C., Ahanger, M. A., Raza, S., Khan, M., Ashraf, M., Ahmed, N., & Zhang, L. (2019). Role of arbuscular mycorrhizal fungi in plant Growth Regulation: Implications in abiotic stress Tolerance. *Frontiers in Plant Science*, 10. <https://doi.org/10.3389/fpls.2019.01068>.
  13. Bisleski, R. L. (1973). Phosphate pools, phosphate transport, and phosphate availability. *Annu. Rev. Plant Physiol.* 24, 225–252. doi: 10.1146/annurev. pp.24.060173.001301.
  14. Bona, E., Cantamessa, S., Massa, N., Manassero, P., Marsano, F., Copetta, A., et al. (2017). Arbuscular mycorrhizal fungi and plant growth-promoting pseudomonads improve yield, quality and nutritional value of tomato: a field study. *Mycorrhiza* 27, 1–C11. doi: 10.1007/s00572-016-0727-y.
  15. Brito, I., Goss, M. J., & De Carvalho, M. (2012). Effect of tillage and crop on arbuscular mycorrhiza colonization of winter wheat and triticale under Mediterranean conditions. *Soil Use and Management*, 28(2), 202–208. <https://doi.org/10.1111/j.1475-2743.2012.00404.x>.

16. Brundrett, Mark C.; Tedersoo, Leho (2018). "Evolutionary history of mycorrhizal symbioses and global host plant diversity". *New Phytologist*. **220** (4): 1108–1115. doi:10.1111/nph.14976. PMID 29355963.
17. Bucher, M. (2007). Functional biology of plant phosphate uptake at root and mycorrhizae interfaces. *New Phytol.* 173, 11–26. doi: 10.1111/j.1469-8137.2006.01935.x.
18. Bücking, H., Liepold, E., & Ambilwade, P. (2012). The role of the mycorrhizal symbiosis in nutrient uptake of plants and the regulatory mechanisms underlying these transport processes. In InTech eBooks. <https://doi.org/10.5772/52570>.
19. Buysens, C., De Boulois, H. D., & Declerck, S. (2014). Do fungicides used to control *Rhizoctonia solani* impact the non-target arbuscular mycorrhizal fungus *Rhizophagus irregularis*? *Mycorrhiza*, 25(4), 277–288. <https://doi.org/10.1007/s00572-014-0610-7>.
20. Castellanos-Morales, V., Villegas, J., Wendelin, S., Vierheiling, H., Eder, R., and Cardenas-Navarro, R. (2010). Root colonization by the arbuscular mycorrhizal fungus *Glomus intraradices* alters the quality of strawberry fruit (*Fragaria ananassa* Duch.) at different nitrogen levels. *J. Sci. Food Agric.* 90, 1774–1782. doi: 10.1002/jsfa.3998.
21. Channabasava, A., Lakshman, H. C., & Jorquera, M. A. (2015). Effect of fungicides on association of arbuscular mycorrhiza fungus *Rhizophagus fasciculatus* and growth of Proso millet (*Panicum miliaceum* L. *Journal of Soil Science and Plant Nutrition*, ahead, 0. <https://doi.org/10.4067/s0718-95162015005000004>.
22. Chen, S., Zhao, H., Zou, C., Li, Y., Chen, Y., Wang, Z., et al. (2017). Combined Inoculation with multiple arbuscular mycorrhizal fungi improves growth, nutrient uptake and photosynthesis in cucumber seedlings. *Front. Microbiol.* 8, 25–16. doi: 10.3389/fmicb.2017.02516.
23. Copetta, A., Lingua, G., Bardi, L., Masoero, G., & Berta, G. (2007). Influence of arbuscular mycorrhizal fungi on growth and essential oil composition in *Ocimum basilicum* var. Genovese. *Caryologia*, 60(1–2), 106–110. <https://doi.org/10.1080/00087114.2007.10589555>.
24. Ercoli L, Schüßler A, Arduini I, Pellegrino E. Strong increase of durum wheat iron and zinc content by field-inoculation with arbuscular mycorrhizal fungi at different soil nitrogen availabilities. *Plant Soil*. 2017;419(1–2): 153–167.
25. Evelin, H., Giri, B., and Kapoor, R. (2012). Contribution of *Glomus intraradices* inoculation to nutrient acquisition and mitigation of ionic imbalance in NaCl-stressed *Trigonella foenum-graecum*. *Mycorrhiza* 22, 203–217. doi: 10.1007/s00572-011-0392-0.
26. Fileccia V, Ruisi P, Ingrassia R, Giambalvo D, Frenda AS, Martinelli F. Arbuscular mycorrhizal symbiosis mitigates the negative effects of salinity on durum wheat. *PloS one*. 2017;12(9): e0184158 10.1371/journal.pone.0184158.
27. Franklin, O.; Näsholm, T.; Högberg, P.; Högberg, M. N. (2014). "Forests trapped in nitrogen limitation - an ecological market perspective on ectomycorrhizal symbiosis". *New Phytologist*. **203** (2): 657–666. doi:10.1111/nph.12840. PMC 4199275. PMID 24824576.

28. Garcés-Ruiz, M., Calonne-Salmon, M., Plouznikoff, K., Misson, C., Navarrete- Mier, M., Cranenbrouck, S., et al. (2017). Dynamics of short-term phosphorus uptake by intact mycorrhizal and non-mycorrhizal maize plants grown in a circulatory semi-hydroponic cultivation system. *Front. Plant Sci.* 8, 1471. doi: 10.3389/fpls.2017.01471.
29. García-Caparrós, P., Lao, M. T., Rangel, P. P., & Sánchez, E. (2021). Phosphorus and carbohydrate metabolism in green bean plants subjected to increasing phosphorus concentration in the nutrient solution. *Agronomy*, 11(2), 245. <https://doi.org/10.3390/agronomy11020245>.
30. Getman-Pickering, Z. L., Stack, G. M., & Thaler, J. S. (2021). Fertilizer quantity and type alter mycorrhizae-conferred growth and resistance to herbivores. *Journal of Applied Ecology*, 58(5), 931–940. <https://doi.org/10.1111/1365-2664.13833>.
31. Gomez-Bellot, M. J., Ortuño, M. F., Nortes, P. A., Vicente-Sánchez, J., Bañón, S., and Sánchez Blanco, M. J. (2015). Mycorrhizal euonymus plants and reclaimed water: biomass, water status and nutritional responses. *Sci. Hort.* 186, 61–69. doi: 10.1016/j.scienta.2015.02.022.
32. Harley, J. L.; Smith, S. E. 1983. *Mycorrhizal symbiosis* (1st ed.). Academic Press, London.
33. Harris, Brogan J.; Clark, James W.; Schrempf, Dominik; Szöllösi, Gergely J.; Donoghue, Philip C. J.; Hetherington, Alistair M.; Williams, Tom A. (2022-09-29). "Divergent evolutionary trajectories of bryophytes and tracheophytes from a complex common ancestor of land plants". *Nature Ecology & Evolution*. 6 (11): 1634–1643. doi:10.1038/s41559-022-01885-x. PMC 9630106. PMID 36175544.
34. Hart, M., Ehret, D. L., Krumbein, A., Leung, C., Murch, S., Turi, C., et al. (2015). Inoculation with arbuscular mycorrhizal fungi improves the nutritional value of tomatoes. *Mycorrhiza* 25, 359–376. doi: 10.1007/s00572-014-0617-0.
35. Higo, M., Isobe, K., Kondo, T., Yamaguchi, M., Takeyama, S., Drijber, R. A., & Torigoe, Y. (2014). Temporal variation of the molecular diversity of arbuscular mycorrhizal communities in three different winter cover crop rotational systems. *Biology and Fertility of Soils*, 51(1), 21–32. <https://doi.org/10.1007/s00374-014-0945-4>.
36. Hijri, M. (2016). Analysis of a large dataset from field mycorrhizal inoculation trials on potato showed highly significant increase in yield. *Mycorrhiza* 2, 209– 214. doi: 10.1007/s00572-015-0661-4.
37. Hoeksema, Jason D.; Bever, James D.; Chakraborty, Sounak; Chaudhary, V. Bala; Gardes, Monique; Gehring, Catherine A.; Hart, Miranda M.; Housworth, Elizabeth Ann; Kaonongbua, Wittaya; Klironomos, John N.; Lajeunesse, Marc J.; Meadow, James; Milligan, Brook G.; Piculell, Bridget J.; Pringle, Anne; Rúa, Megan A.; Umbanhowar, James; Viechtbauer, Wolfgang; Wang, Yen-Wen; Wilson, Gail W. T.; Zee, Peter C. (16 August 2018). "Evolutionary history of plant hosts and fungal symbionts predicts the strength of mycorrhizal mutualism". *Communications Biology*. 1 (1): 116. doi:10.1038/s42003-018-0120-9.

38. Huey, C. J., Gopinath, S. C. B., Uda, M. N. A., Zulhaimi, H. I., Jaafar, M. S., Kasim, F. H., & Yaakub, A. R. W. (2020). Mycorrhiza: a natural resource assists plant growth under varied soil conditions. *3 Biotech*, 10(5). <https://doi.org/10.1007/s13205-020-02188-3>.
39. Jiang, Y. N., Wang, W. X., Xie, Q. J., Liu, N., Liu, L. X., Wang, D. P., et al. (2017). Plants transfer lipids to sustain colonization by mutualistic mycorrhizal and parasitic fungi. *Science* 356, 1172–1175. doi: 10.1126/science.aam9970.
40. Kayama, M., and Yamanaka, T. (2014). Growth characteristics of *ectomycorrhizal* seedlings of *Quercus glauca*, *Quercus salicina*, and *Castanopsis cuspidata* planted on acidic soil. *Trees* 28, 569–583. doi: 10.1007/s00468-013-0973-y.
41. Kiers, E. T., Duhamel, M., Beesetty, Y., Mensah, J. A., Franken, O., Verbruggen, E., Fellbaum, C. R., Kowalchuk, G. A., Hart, M. M., Bago, A., Palmer, T. M., West, S. A., Vandenkoornhuyse, P., Jansa, J., & Bücking, H. (2011). Reciprocal rewards stabilize cooperation in the mycorrhizal symbiosis. *Science*, 333(6044), 880–882. <https://doi.org/10.1126/science.1208473>.
42. Laishram, B., Devi, O. R. and Ngairangbam, H. 2023. Insight into Microbes for Climate SmartAgriculture. *Vigyan Varta* 4(4): 53-56.
43. Lehmann, A., and Rillig, M. C. (2015). Arbuscular mycorrhizal contribution to copper, manganese and iron nutrient concentrations in crops—a meta-analysis. *Soil Biol. Biochem.* 81, 147–158. doi: 10.1016/j.soilbio.2014.11.013.
44. Lehmann, A., Veresoglou, S. D., Leifheit, E. F., and Rillig, M. C. (2014). Arbuscular mycorrhizal influence on zinc nutrition in crop plants: a meta-analysis. *Soil Biol. Biochem.* 69, 123–131. doi: 10.1016/j.soilbio.2013.11.001.
45. Lin, C., Wang, Y., Li, M., Li, Q., Xiao, W., & Song, X. (2020). Effects of nitrogen deposition and phosphorus addition on arbuscular mycorrhizal fungi of Chinese fir (*Cunninghamia lanceolata*). *Scientific Reports*, 10(1). <https://doi.org/10.1038/s41598-020-69213-6>.
46. Liu, C., Ravnskov, S., Liu, F., Rubæk, G. H., and Andersen, M. N. (2018). Arbuscular mycorrhizal fungi alleviate abiotic stresses in potato plants caused by low phosphorus and deficit irrigation/partial root-zone drying. *J. Agric. Sci.* 156, 46–58. doi: 10.1017/S0021859618000023.
47. Lu, F., Lee, C., and Wang, C. (2015). The influence of arbuscular mycorrhizal fungi inoculation on yam (*Dioscorea* spp.) tuber weights and secondary metabolite content. *Peer J.* 3, 12–66. doi: 10.7717/peerj.1266.
48. Mbuthia, L. W., Acosta-Martínez, V., DeBruyn, J. M., Schaeffer, S. M., Tyler, D. D., Odoi, E. W., Mpheshea, M., Walker, F., & Eash, N. S. (2015). Long term tillage, cover crop, and fertilization effects on microbial community structure, activity: Implications for soil quality. *Soil Biology & Biochemistry*, 89, 24–34. <https://doi.org/10.1016/j.soilbio.2015.06.016>.

49. Mitra, D., Navendra, U., Panneerselvam, U., Ansuman, S., Ganeshamurthy, A. N., and Divya, J. (2019). Role of mycorrhiza and its associated bacteria on plant growth promotion and nutrient management in sustainable agriculture. *Int. J. Life Sci. Appl. Sci.* 1, 1–10.
50. Miyauchi, Shingo; Kiss, Enikő; Kuo, Alan; et al. (2020). "Large-scale genome sequencing of mycorrhizal fungi provides insights into the early evolution of symbiotic traits". *Nature Communications*. **11** (1): 5125. Bibcode:2020NatCo..11.5125M. doi:10.1038/s41467-020-18795-w. PMC 7550596. PMID 33046698.
51. Nell, M., Wawrosch, C., Steinkellner, S., Vierheilig, H., Kopp, B., and Lössl, A. (2010). Root colonization by symbiotic arbuscular mycorrhizal fungi increases sesquiterpenic acid concentrations in *Valeriana officinalis* L. *Planta Med.* 76, 393–398. doi: 10.1055/s-0029-1186180.
52. Ortas, I. (2012). The effect of mycorrhizal fungal inoculation on plant yield, nutrient uptake and inoculation effectiveness under long-term field conditions. *Field Crops Res.* 125, 35–48. doi: 10.1016/j.fcr.2011.08.005.
53. Pellegrino, E., and Bedini, S. (2014). Enhancing ecosystem services in sustainable agriculture: biofertilization and biofortification of chickpea (*Cicer arietinum* L.) by arbuscular mycorrhizal fungi. *Soil Biol. Biochem.* 68, 429–439. doi: 10.1016/j.soilbio.2013.09.030.
54. Prasad, R., Bholá, D., Akdi, K., Cruz, C., Sairam, K. V. S. S., Tuteja, N., et al. (2017). Introduction to mycorrhiza: historical development," in *Mycorrhiza*. Eds. A. Varma, R. Prasad, and N. Tuteja (Cham: Springer), 1–7. doi: 10.1007/978-3-319-53064-2\_1.
55. Provorov, N. A.; Shtark, O. Yu; Dolgikh, E. A. (2016). "[Evolution of nitrogen-fixing symbioses based on the migration of bacteria from mycorrhizal fungi and soil into the plant tissues]". *Zhurnal Obshchei Biologii.* **77** (5): 329–345. PMID 30024143.
56. Pugnier, Camille; Keller, Jean; Delaux, Pierre-Marc (2022-08-29). "Plant–microbe interactions that have impacted plant terrestrializations". *Plant Physiology.* **190** (1): 72–84. doi:10.1093/plphys/kiac258. PMC 9434271. PMID 35642902.
57. Püschel D, Janoušková M, Voříšková A, Gryndlerová H, Vosátka M, Jansa J. Arbuscular mycorrhiza stimulates biological nitrogen fixation in two Medicago spp. through improved phosphorus acquisition. *Front Plant Sci.* 2017;8: 390 10.3389/fpls.2017.00390.
58. Qiao X, Bei S, Li C, Dong Y, Li H, Christie P, Zhang F, Zhang J. Enhancement of faba bean competitive ability by arbuscular mycorrhizal fungi is highly correlated with dynamic nutrient acquisition by competing wheat. *Sci Rep.* 2015;5: 8122 10.1038/srep08122.
59. Remy, W.; Taylor, T. N.; Hass, H.; Kerp, H. (6 December 1994). "Four hundred-million-year-old vesicular arbuscular mycorrhizae". *Proceedings of the National Academy of Sciences.* **91** (25): 11841-11843.

60. Rillig MC, Mummey DL. Mycorrhizas and soil structure. *New Phytol.* 2006;171(1): 41–53. doi: 10.1111/j.1469-8137.2006.01750.x
61. Roupahel, Y., Franken, P., Schneider, C., Schwarz, D., Giovannetti, M., and Agnolucci, M. (2015). Arbuscular mycorrhizal fungi act as bio-stimulants in horticultural crops. *Sci. Hort.* 196, 91–108. doi: 10.1016/j.scienta.2015.09.002.
62. Sabia, E., Claps, S., Morone, G., Bruno, A., Sepe, L., and Aleandri, R. (2015). Field inoculation of arbuscular mycorrhiza on maize (*Zea mays* L.) under low inputs: preliminary study on quantitative and qualitative aspects. *Italian J. Agron.* 10, 30–33. doi: 10.4081/ija.2015.607.
63. Sbrana, C., Avio, L., and Giovannetti, M. (2014). Beneficial mycorrhizal symbionts affecting the production of health-promoting phytochemicals. *Electrophoresis* 35, 1535–1546. doi: 10.1002/elps.201300568.
64. Secilia J, Bagyaraj DJ. Bacteria and actinomycetes associated with pot cultures of vesicular–arbuscular mycorrhizas. *Can J Microbiol.* 1987;33: 1069–1073.
65. Siasou E, Standing D, Killham K, Johnson D. Mycorrhizal fungi increase biocontrol potential of *Pseudomonas fluorescens*. *Soil Biol Biochem.* 2009;41(6): 1341–1343.
66. Smith SE, Read D. The symbionts forming arbuscular mycorrhizas, in: Smith SE, Read D, editors. *Mycorrhizal symbiosis* (3rd edition). New York: Academic Press; 2008. pp. 13–41.
67. Smith, S. E., and Read, D. J. (1997). *Mycorrhizal symbiosis*. San Diego: Academic Press, 607.
68. Smith, S. E., Jakobsen, I., Grnlund, M., and Smith, F. A. (2011). Roles of arbuscular mycorrhizas in plant phosphorus nutrition: interactions between pathways of phosphorus uptake in arbuscular mycorrhizal roots have important implications for understanding and manipulating plant phosphorus acquisition. *Plant Physiol.* 156, 1050–1057. doi: 10.1104/pp.111.174581.
69. Trappe, J. M. (1987). "Phylogenetic and ecologic aspects of mycotrophy in the angiosperms from an evolutionary standpoint". In Safir, G. R. (ed.). *Ecophysiology of VA Mycorrhizal Plants*. Florida: CRC Press
70. Turrini, A., Bedini, A., Loor, M. B., Santini, G., Sbrana, C., Giovannetti, M., et al. (2018). Local diversity of native arbuscular mycorrhizal symbionts differentially affects growth and nutrition of three crop plant species. *Biol. Fertil. Soils* 54, 203–217. doi: 10.1007/s00374-017-1254-5.
71. Vallino, M., Fiorilli, V., & Bonfante, P. (2013). Rice flooding negatively impacts root branching and arbuscular mycorrhizal colonization, but not fungal viability. *Plant, Cell & Environment*, 37(3), 557–572. <https://doi.org/10.1111/pce.12177>.
72. Wagg, C., Barendregt, C., Jansa, J., and Heijden, M. G. A. (2015). Complementarity in both plant and mycorrhizal fungal communities are not necessarily increased by diversity in the other. *J. Ecol.* 103, 1233–1244. doi: 10.1111/1365-2745.12452.

73. Zaefarian, F., Rezvani, M., Ardakani, M. R., Rejali, F., and Miransari, M. (2013). Impact of mycorrhizae formation on the phosphorus and heavy-metal uptake of Alfalfa. *Comm. Soil Sci. Plant Anal.* 44, 1340–1352. doi: 10.1080/00103624.2012.756505.
74. Zeng, L., JianFu, L., JianFu, L., and MingYuan, W. (2014). Effects of arbuscular mycorrhizal (AM) fungi on citrus quality under nature conditions. *Southwest China J. Agric. Sci.* 27, 2101–2105. doi: 10.16213/j.cnki.scjas.2014.05.067.
75. Zhang, F., Jia-Dong, H. E., Qiu-Dan, N. I., Qiang-Sheng, W. U., and Zou, Y. N. (2018). Enhancement of drought tolerance in trifoliolate orange by mycorrhiza: changes in root sucrose and proline metabolisms. *Not. Bot. Horti. Agrobot. Cluj-Napoca* 46, 270. doi: 10.15835/nbha46110983.
76. Zhu, X. C., Song, F. B., Liu, S. Q., and Liu, F. L. (2016). Arbuscular mycorrhiza improve growth, nitrogen uptake, and nitrogen use efficiency in wheat grown under elevated CO<sub>2</sub>. *Mycorrhiza* 26, 133–140. doi: 10.1007/s00572-015-0654-3.