

## Comparative efficacy and economics of selected chemicals and biopesticides against pod borer [*Helicoverpa armigera* (Hubner)] on chickpea (*Cicer arietinum* L.)

### ABSTRACT

The research work was undertaken at Central Research Farm (CRF) Sam Higginbottom University of Agriculture Technology and Sciences, SHUATS, Naini, Prayagraj during winter season of 2023- 24. The treatments consists of eight including control viz, T<sub>1</sub>- *Beauveria bassiana* 1.15 % WP, T<sub>2</sub> Chlorantraniliprole 18.5 SC, T<sub>3</sub>-Emamectin benzoate 5 SG, T<sub>4</sub> - *Bacillus thuringiensis* 1x10<sup>9</sup> CFU/ml, T<sub>5</sub> -Azadirachtin 00.03% WSP, T<sub>6</sub> – NSKE 5% , T<sub>7</sub>- Spinosad 45 SC and T<sub>0</sub>-untreated control arranged in Randomized Block Design (RBD) with three replications targeting to evaluate the efficacy of selected insecticides on the larval population of *H. armigera* on Chickpea. The larval population of chickpea pod borer , *Helicoverpa armigera* on third, seven and fourteen days after spray revealed that among all the treatment T<sub>2</sub> Chlorantraniliprole 18.5 SC found superior with larval population of (1.05), and with highest cost benefit ratio (1:3.78 ), followed by Emamectin benzoate 5 SG with a larval population of (1.17) and cost benefit ratio (1:3.53), Spinosad 45 SC with a larval population of (1.24) and cost benefit ratio (1:3.16), *Beauveria bassiana* 1.15 % WP with a larval population of (1.32) and cost benefit ratio (1:3.00 ), *Bacillus thuringiensis* 1x10<sup>9</sup> CFU/ml with a larval population(1.35) and cost benefit ratio ( 1:2.76 ), Azadirachtin 00.03% WSP with a larval population(1.42) and cost benefit ratio (3.49, 1:2.25), NSKE 5% with a larval population (1.47) and cost benefit ratio ( 1:1.77), NSKE 5% was least effective among the treatments and control plot T<sub>0</sub> with a larval population (2.07) and cost benefit ratio (1:1.51) .

**Keywords:** Biopesticides, chemicals, chickpea, cost benefit ratio, efficacy, *Helicoverpa armigera*.

## INTRODUCTION:-

---

“Gram commonly known as a ‘chickpea’ or chana is a self-pollinating diploid ( $2n=2x=16$ ) plant. It is originated in South-eastern Turkey and spread to other parts of the world. A very important pulse crop grows as a seed of a plant named *Cicer arietinum* (L.) in the Leguminosae family. According to De Candolle, “Chanaka” which is the Sanskrit name of chickpea gives the indication of being cultivated in India from a very long duration compared to other countries in the world”. **(Kumar and Yadav, 2023)**.

“India ranks first in the production and consumption of chickpea (*Cicer arietinum* L.) in the world. Chickpea is a most important pulse crop of India, which is mostly grown under dry land condition with heavy cloudy soil. It is a rich source of nutritional values in the diet of Indian people because of containing 21.5 per cent protein, 64.5 per cent carbohydrates and 4.5 per cent fat which is comparatively deficient in the cereals and oilseeds. The people in their daily meals use its green leaves and pods as green vegetables and germinated grains for breakfast and other delicious dishes”. **(Kumar et al., 2019)**.

“Several pests, mainly insects, attack chickpea. Sarwar, recorded “57 insect species, namely Lepidoptera as *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae), commonly known as gram pod borer is a major polyphagous and noctuid pest in Asia, causing heavy damage to agricultural, horticultural and ornamental crops”. “In India, the extent of losses due to *H. armigera* in chickpea is up to 27.9 per cent in North West Plain Zone, 13.2 per cent in North East Plain Zone, 24.3 per cent in Central Zone and 36.4 per cent in South Zone. The crops have been noticed to suffer an avoidable loss of 9 to 60 per cent by this insect. In Uttar Pradesh, alone 15.3 per cent of the chickpea crop worth ₹.462.5 million is lost annually due to *H. armigera* attack, 17.2 per cent in Karnataka and 28.5per cent in Delhi”. **(Bhati et al., 2023)**.

## MATERIALS AND METHODS:-

---

The experiment was conducted at the experimental research plot of the Department of Entomology, Central Research Farm, Sam Higginbottom University of Agriculture Technology and Sciences, during the winter season of 2023-24. The Research field is situated at 25°27 North Latitude 80°50 East Longitudes and at an altitude of 98 meter above sea level. The climate is typically semi- arid and sub-tropical. The maximum temperature reaches up to 47°C in summer and drops down to 2.5°C in winter. The experimental design was Randomized Block Design with eight treatments, each replicated thrice. The plot size was ( 2m × 1m) with a spacing of (30×10 cm). The treatments included - *Beauveria bassiana* 1.15 % WP (Bevroz) , Chlorantraniliprole 18.5 SC (Coragen) , Emamectin benzoate 5 SG (Proclaim), *Bacillus thuringiensis* 1×10<sup>9</sup> CFU/ml (Thuricide), Azadirachtin 00.03% WSP (Neemaura), NSKE 5% (Neemicide), Spinosad 45 SC (Tracer), and a control.

The numbers of larva were counted on five randomly selected plants in each plot. The pre-treatment count was made a day before the spray whereas, the post-treatment counts were made on 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> day after each spray. The larval population over control against gram pod borer was calculated by considering the mean of three observations recorded at 3<sup>rd</sup>, 7<sup>th</sup>, and 14<sup>th</sup> day after spray

## RESULT AND DISCUSSION

---

### 3.1 Efficacy of *Helicoverpa armigera* after first spray

#### Third days after spraying:

The data of gram pod borer (Table 1) after three spray days revealed that all treatments was significantly superior over the control. Among all the treatments, the lowest number of larval populations was recorded in Chlorantraniliprole 18.5 SC (1.40) found superior over other treatments followed by Emamectin benzoate (1.53), Spinosad 45 SC (1.60), *Beauveria bassiana* (1.67), *Bacillus thuringiensis*  $1 \times 10^9$  CFU/ml (1.67), Azadirachtin 00.03% WSP (1.80) and Neem seed kernel extract 5% (1.80) is found to be least effective among all the treatments as compared to control (2.00).

#### Sevan days after spraying:

The data of gram pod borer (Table 1) after three spray days revealed that all treatments was significantly superior over the control. Among all the treatments, Chlorantraniliprole 18.5 SC (1.07) found superior over other treatments followed by Emamectin benzoate (1.13), Spinosad 45 SC (1.20), *Beauveria bassiana* (1.27), *Bacillus thuringiensis*  $1 \times 10^9$  CFU/ml (1.27), Azadirachtin 00.03% WSP (1.33 ) and Neem seed kernel extract 5% (1.40) is found to be least effective among all the treatments as compared to control (2.07).

#### Fourteen days after spraying:

The data of gram pod borer (Table 1) after three spray days revealed that all treatments was significantly superior over the control. Among all the treatments, Chlorantraniliprole 18.5 SC (0.67) found superior over other treatments followed by Emamectin benzoate (0.87), Spinosad 45 SC (0.93), *Beauveria bassiana* (1.00), *Bacillus thuringiensis*  $1 \times 10^9$  CFU/ml (1.07), Azadirachtin 00.03% WSP (1.13), and neem seed kernel extract 5% (1.20) is found to be least effective among all the treatments as compared to the control (2.13).


**Overall mean of first spray:**

The data on the larval population of gram pod borer (Table 1) on mean (3<sup>rd</sup>, 7<sup>th</sup>, 14<sup>th</sup> DAS) days after spray revealed that all the treatments was significantly superior over control. Among all the treatments, Chlorantraniliprole 18.5 SC (1.05) found superior over other treatments followed by Emamectin benzoate (1.17), Spinosad 45 SC (1.24), *Beauveria bassiana* (1.32), *Bacillus thuringiensis*  $1 \times 10^9$  CFU/ml (1.35), Azadirachtin 00.03% WSP (1.42) and Neem seed kernel extract 5% (1.47) is found to be least effective among all the treatments as compared to control (2.07).

**Table .1 Effect of certain insecticides and biopesticides on the larval population of *Helicoverpa armigera* on chickpea during rabi season 2023 : 24**

Treatments		Number of larvae / 5 plants				
		1DBS	After spray			
			3 <sup>rd</sup> Day	7 <sup>th</sup> Day	14 <sup>th</sup> Day	Mean
<b>T<sub>0</sub></b>	Control	1.93	2.00	2.07	2.13	2.07
<b>T<sub>1</sub></b>	<i>Beauveria bassiana</i> 1.15 % WP	1.80	1.70	1.27	1.00	1.32
<b>T<sub>2</sub></b>	Chlorantraniliprole 18.5 SC	1.67	1.40	1.07	0.67	1.05
<b>T<sub>3</sub></b>	Emamectin benzoate 5 SG	1.73	1.50	1.13	0.87	1.17
<b>T<sub>4</sub></b>	<i>Bacillus thuringiensis</i> 1x10 <sup>9</sup> CFU/ml	1.80	1.70	1.27	1.07	1.35
<b>T<sub>5</sub></b>	Azadirachtin 00.03% WSP	1.73	1.80	1.33	1.13	1.42
<b>T<sub>6</sub></b>	NSKE 5%	1.80	1.80	1.40	1.20	1.47
<b>T<sub>7</sub></b>	Spinosad 45 SC	1.73	1.60	1.20	0.93	1.24
Overall Mean		1.77	1.69	1.34	1.13	1.39
F- test		NS	S	S	S	S
S. Ed. (±)			0.08	0.10	0.10	0.05
C. D. (P = 0.05)			0.185	0.228	0.224	0.255

-



**Figure 1. Efficacy of selected chemicals and biopesticides against larval population of gram pod borer (*H. armigera*) on chickpea**

**Table .2 Economics of cultivation**

<b>Sr. No:</b>	<b>Treatment</b>	<b>Yield of q/ha</b>	<b>B:C ratio</b>
<b>T<sub>0</sub></b>	Control	11.00	1:1.51
<b>T<sub>1</sub></b>	<i>Beauveria bassiana</i> 1.15 % WP	22.50	1:3.00
<b>T<sub>2</sub></b>	Chlorantraniliprole 18.5 SC	29.10	1:3.78
<b>T<sub>3</sub></b>	Emamectin benzoate 5 SG	26.65	1:3.53
<b>T<sub>4</sub></b>	<i>Bacillus thuringiensis</i> 1x10 <sup>9</sup> CFU/ml	21.24	1:2.76
<b>T<sub>5</sub></b>	Azadirachtin 00.03% WSP	17.08	1:2.25
<b>T<sub>6</sub></b>	NSKE 5%	13.50	1:1.77
<b>T<sub>7</sub></b>	Spinosad 45 SC	25.8	1:3.16

“The data on mean population after spray revealed that all the insecticides were found very effective and significantly superior over untreated control”. [18] Among all the treatments minimum number of larvae were found in T<sub>2</sub> Chlorantraniliprole 18.5 SC (1.05) as the similar findings was reported by **Jayanth and Kumar (2022), and Bhati et al., (2023)** to control *Helicoverpa armigera* larval population. T<sub>3</sub>- Emamectin benzoate 5SG (1.17) was found the next effective treatment with larval population (1.17). Abbas *et al.*, (2021), and **Reddy and Tayde (2023)** reported similar finding, for reducing the larval population of *Helicoverpa armigera*. T<sub>7</sub>-Spinosad 45 SC (1.24) was found the next best effective treatments which was similarly found by **Ravicharan and Tayde (2023)** who reported Spinosad 45 SC to be the next best and effective treatment in controlling larval population , T<sub>1</sub>- *Beauveria bassiana* 1.15 WP (1.32) was found the next most effective treatment with the same findings was done by **Sai et al.,(2021), Sireesha and Kumar (2022)** T<sub>4</sub>-*Bacillus thuringiensis* 1×10<sup>9</sup>CFU/ml was found the next best effective treatment with a larval population of (1.35) as the same findings was done by **Abbas et al., (2021) and Yerrabala et al., (2021)**, T<sub>2</sub> Azadirachtin 00.03% WSP was foundthe next effective treatment with a larval population of (1.42) as the same findings was done by **Santosh and Kumar (2022)** and **Gautam et al.,(2018)**,T<sub>6</sub> NSKE 5% was found the least effective treatment with a larval population of (1.47) and the same findings was done by **Machindra and Kumar (2022)**.

When the cost benefit ratio worked out, interesting result was achieved. Among all the treatments the higher cost benefit ratio was obtained from T<sub>2</sub> Chlorantraniliprole 18.5 SC (1:3.78), as the similar findings was done by **Barwa and Kumar (2022) (1:3.35)**, **Bhati et al.,(2023) (1:3.49)**, followed by the T<sub>3</sub> Emamectin benzoate 5SG exhibited a cost benefit ratio of (1:3.53) as the similar finding was done by ), **Bhati et al.,(2023) (1:2.66)**, followed by T<sub>7</sub> Spinosad 45 SC with a cost benefit ratio of (1:3.16) as the similar finding was done by **Chandra et al., (2017) (1:2.36)**, which was followed by T<sub>1</sub> *Beauveria bassiana* which exhibited cost benefit ratio of (1:3.00) which was supported by the finding of by **Anil and Kumar (2022) (1:2.96)**, followed by T<sub>4</sub> *Bacillus thuringiensis* exhibited cost benefit ratio of (1:2.76) it was supported by **Sireesha and Kumar (2022) (1:3.39)**, which was followed by T<sub>5</sub> Azadirachtin 00.03% WSP with a cost benefit ratio of (1:2.25) as the similar finding was done **Santosh and Kumar (2022) (1:2.41)**, Followed by T<sub>6</sub> NSKE 5% which obtained a cost benefit ratio of (1:1.77) which was supported by **Sharma and Tayde (2023) (1:1.75)**

## CONCLUSION

---

From research it was found that, spraying of insecticides significantly reduced the pod borer population in chickpea. The present findings conclude that the new generation insecticides like T<sub>1</sub> *Beauveria bassiana* 1.15 WP, T<sub>2</sub> Chlorantraniliprole 18.5% SC, T<sub>3</sub> Emamectin benzoate 5%SG, T<sub>4</sub> *Bacillus thuringiensis* 1×10<sup>9</sup> CFU/ml, T<sub>5</sub> Azadirachtin 00.03 WSP, T<sub>6</sub> NSKE 5%, T<sub>7</sub> Spinosad 45 SC. T<sub>2</sub> Chlorantraniliprole was found effective against lepidopteran caterpillar *Helicoverpa armigera* along with an additional yield level in chickpea. Further, it was observed that the cost benefit ratio was also high with Chlorantraniliprole 18.5% SC, Emamectin benzoate 5 SG and Spinosad 45% SC. Hence, it is suggested that the effective insecticides may be alternated in harmony with the existing integrated pest management programs in order to avoid the problems associated with insecticidal resistance, pest resurgence.

## REFERENCES

1. **Abbas, A., Wang, Y., Muhammad, U. and Fatima, A. (2021).** Efficacy of different insecticides against gram pod borer (*Helicoverpa armigera*) and their safety to the beneficial fauna. *International Journal of Biosciences*, **18**: 82-88.
2. **Anil, M. P. and Kumar, A. (2022).** Comparative efficacy and economics of certain chemicals and biopesticides against pod borer, *Helicoverpa armigera*(Hubner) in Chickpea at Naini, Prayagraj, India. *International Journal of Plant & Soil Science*, **34**(22), 269-276.
3. **Barwa, J. and Kumar, A. (2022).** Field efficacy of Chlorantraniliprole with some biopesticides against pod borer [*Helicoverpa armigera* (Hubner)] on chickpea (*Cicer arietinum* L.). *The Pharma Innovation Journal*, **11**(6): 1912-1916.
4. **Bhati, M. S., Rajnikant, T. A., Singh, S. P. and Chandar, A. S. (2023).** Effect of Selected Biopesticides and Chemicals against Pod Borer [*Helicoverpa armigera* (L.)] on Chickpea (*Cicer arietinum* L.). *International Journal of Plant and Soil Science*, **35**(18): 423-426.
5. **Chandra, GV., Kumar, A., Lavanya, V., and Sayad, R. (2017).** Efficacy of Certain Chemicals and Neem Products against *Helicoverpa armigera* (Hubner) on chickpea (*Cicer arietinum* L.). *Journal of Entomology and Zoology Studies* **5**(2): 01-05
6. **Gautam, M. P., Chandra, U., Yadav, S. K., Jaiswal, R., Giri, S. K. and Singh, S. N. (2018).** Studies on population dynamics of gram pod borer *Helicoverpa armigera* (Hubner) on chickpea (*Cicer arietinum* L.). *Journal of Entomology and Zoology Studies*, **6**(1): 904-906.
7. **Jayanth, T. and Kumar, A. (2022).** Field efficacy of selected insecticides with combination of neem oil against gram pod borer [*Helicoverpa armigera* (Hubner)]. *The Pharma Innovation Journal*, **11**(5): 465-469.
8. **Kumar, A. and Yadav, U. (2023).** Efficacy of different plant products against gram pod borer [*Helicoverpa armigera* (Hubner)] on chickpea, *Cicer arietinum* (L.) *The Pharma Innovation Journal* 2023; **12**(7): 1922-1927.
9. **Kumar, A., Tripathi, M. K., Chandra, U. and Veer, R. (2019).** Efficacy of botanicals and bio-pesticide against *Helicoverpa armigera* in chickpea. *Journal of Entomology and Zoology Studies*, **7**(1): 54-57.
10. **Machindra, J., P. and Kumar, A. (2022).** Field efficacy and economics of different chemicals

- and biopesticides against pod borer, *Helicoverpa armigera* (Hubner) on chickpea during *rabi* season. *The Pharma Innovation Journal* 2022, SP-11(7): 3970-3975.
11. **Ravicharan, C., and Tayde, A. R. (2023).** Field efficacy of Selected Insecticides against Pod Borer, *Helicoverpa armigera* (H.) in Chick Pea (*Cicer arietinum* Linnaeus). *Biological Forum – An International Journal*, **15**(6): 220-223
  12. **Reddy, D. V., and Tayde, A. R. (2023).** Comparative efficacy of different insecticides against fruit borer, *Helicoverpa armigera* (Hubner) on tomato, *Solanum Lycopersicon* (L.). *Annals of Plant Protection Sciences*, **31**(2): 122-126.
  13. **Sai, K. P., Tayde, A. R., Reddy, R. D. and Srivastava, V. K. (2021).** Management of gram pod borer *Helicoverpa armigera* (Hubner) in chickpea with bio-pesticides and combination. *Journal of Entomology and Zoology Studies*, **9**(1): 561-563.
  14. **Santhosh, K., and Kumar, A. (2022).** Comparative efficacy of selected insecticides and neem products against chickpea pod borer [*Helicoverpa armigera* (Hubner)]. *The Pharma Innovation Journal*. **11**(6): 1558-1562
  15. **Sharma, D., & Rajnikant, T. A. (2023).** Bioefficacy of Certain Chemicals and Biopesticides against Pod Borer [*Helicoverpa armigera* (Hubner)] on Chickpea (*Cicer arietinum* L.). *International Journal of Plant & Soil Science*, **35**(18), 49-54.
  16. **Sireesha, B. S. and Kumar, A. (2022).** Efficacy of selected insecticides against pod borer [*Helicoverpa armigera* (Hubner)] on green gram. *The Pharma Innovation Journal*, SP- **11**(8): 944-948.
  17. **Yerrabala, S., Kumar, H. S., and Yadav, U. (2021).** Comparative efficacy of *Bacillus thuringiensis* with botanicals and chemicals against gram pod borer *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) on cowpea [*Vigna unguiculata* (L.) Walp.]. *The PharmaInnovation Journal*, SP-**10**(5): 709-712.
  18. **Moosan M, Kumar A.** Efficacy and economics of different insecticides against gram pod borer, *Helicoverpa armigera* (Hubner). *The Pharma Innovation Journal*. 2022;**11**(7):4471-4.

