

# Edible Flowers in Herbal Beverage Development

---

## ABSTRACT

Herbal beverages with nutraceutical benefits are becoming more widely acknowledged as healthy dietary choices among mindful consumers. There are numerous types of herbal beverages prepared from different morphological plant elements. Edible flowers, with their unique flavors, distinct aromas, vivid colors, and nutraceutical properties, have gained predominance among all the other botanical ingredients utilized in the formulation of herbal beverages. This review elaborates on applications of edible flowers used in the preparation of herbal teas, infusions, and a few other novel developments in herbal beverages. Although several commercialized herbal beverages have been developed with edible flowers, many other edible flowers with medicinal properties remain unexplored. The review further elaborates on the pharmacotherapeutic properties of these flowers, emphasizing their potential to be used in herbal beverage formulations. Additionally, ensuring the food safety aspects of innovative herbal beverages derived from edible flowers by addressing their potential toxicity and microbiological quality is vitally important. Insights presented here pave the way for incorporating edible flowers in herbal beverages, encouraging nutritious and health-promoting beverage development and consumption.

*Keywords: Edible flowers, herbal beverages, phytochemicals, extraction, preservation, health benefits*

## 1. INTRODUCTION

The nutraceutical segment continues to expand as a preventive approach against non-communicable diseases like hypertension, cancer, and diabetes, ultimately enhancing human life expectancy. Nutraceutical products are consumed by approximately 50-70% of the developed countries and usage among women was higher compared to men [1]. Herbal remedies are widely gaining acceptance as a complementary medical option worldwide, including in the USA and Europe [2].

Intake of nonnutritive nutraceutical compounds from dietary sources available in different palatable forms such as pharmaceutical or food supplements, food additives, and herbal beverages exhibited a marked growth in recent years. Herbal beverages have been used in traditional medicine over decades and have become a popular global beverage choice now. Therefore, herbal beverages are accepted as therapeutic carriers and a viable solution for the consumption of valuable plant nutraceuticals [3].

Traditionally edible flowers are consumed as teas or infusions; however, recent research studies have exhibited their potential to be used in novel herbal beverage developments owing to their unique taste, aroma, and colour properties. Other than utilizing edible flowers alone in herbal beverages, blending them with other plant-based ingredients with medicinal and nutritional properties has generated more palatable beverage solutions.

## **2. HERBAL BEVERAGE DIVERSITY AND CONSUMPTION TRENDS**

The consumption of herbal beverages is embedded in the lifestyle of many Asians, Europeans, and Americans due to their health benefits, pleasant taste, and cultural influences. A wide range of herbal beverages can be identified and differentiated based on the preparation method and specific plant components utilized. In that context, herbal infusions play a vital role in traditional medicine where edible plant components with pharmacological values have been used as therapeutic agents over the years [4]. Furthermore, it has gained popularity among health-conscious consumers who are heavily concerned about the health benefits of the beverages they consume.

“Infusion” is when boiling water is poured over the plant material selected and allowed to settle for a few minutes followed by straining [5]. Preparation of infusions is usually completed at ambient temperature and the quality of the extraction depends on the bulk density of the herb used [6]. “Decoction” is a plant infusion specially made from the fibrous part of the plant material such as stem, bark, seed, rhizome, and woods [7]. If the infusion consists of two or more different plants it is known as “tisane” [8]. When the infusions are made specifically from the leaves of *Camellia Sinensis* it is named as “tea” [9].

The majority of herbal teas and infusions are prepared just before drinking and often served hot. In this context, herbal teas are available in various forms: whole dried plant components, dried flakes, granules within tea bags, or sold in bulk. The plant material undergoes initial processing, involving drying, comminution, and crushing. Alternatively, instant tea, a completely soluble powder, is a concentrated tea infusion that undergoes water removal through spray drying. To preserve flavor, aroma, and nutritional quality, instant teas may undergo methods like vacuum drying and freeze-drying as well [10]. However, a recent trend shows a surge in the popularity of ready-to-drink bottled herbal beverages [4]. For instance, iced tea, crafted from diluted herbal plant extracts, is marketed in bottles or cans for immediate consumption. Additionally, the realm of herbal beverage products sees innovation with carbonated herbal teas, presenting a promising avenue for commercial success. In fortified herbal beverages, a variety of ingredients are incorporated, spanning polyphenols, carotenoids (like astaxanthin, and lutein), Omega-3 oils ( $\alpha$ -linolenic, eicosapentaenoic, docosahexaenoic acids), sterols, stimulants, and botanicals (superfruits, teas, herbal extracts). These additions synergistically amplify health benefits while ensuring palatability and balance in formulations [11].

## **3. INCORPORATING PLANT ELEMENTS INTO HERBAL BEVERAGES**

There are many edible plant varieties but, in most species, the edible portion is restricted to one or few morphological plant parts. Phytochemical compounds can be isolated from some of these edible plant varieties, but they may most probably be not from the edible portion of the plant [12]. Different plant elements including leaves, seeds, fruits, stems, roots, peels, barks, and flowers, and their plant extracts are used in the preparation of herbal beverages. Flowers which are mostly recognized for their ornamental value due to their attractive colors, shapes, and exotic aroma are now gaining more attention for their medicinal properties in food and pharmaceutical applications. Some flowers are technically edible but less

palatable. Petals are the mostly consumed component in edible flowers although pollen, nectar, and other parts are also consumed [13]. Flowers, being more delicate but rich in bioactive compounds, flavor properties, characteristic aroma, and colour make a significant contribution mainly to herbal infusions and other novel applications in the herbal beverage industry.

There are 97 families, 100 genera, and 180 species of edible flowers found around the world excluding flowers based on vegetables such as broccoli, cauliflower, and artichoke [14]. Two types of edible flowers can be identified namely fruit flowers and non-fruit flowers. Fruit flowers such as Papaya, Apple, Peach, and Banana require to undergo a pre-heat treatment process before consumption. Non-fruit flowers such as Butterfly pea flower, Hibiscus, Rose, and Lotus are commonly categorized as ornamental flowers. Usually, this category of flowers has vividly colored petals & stamens [15]. There are several naturally occurring tree flowers such as Neem (*Azadirachta indica*), Sesbania (*Sesbania grandiflora*), Bauhinia (*Bauhinia racemosa*) & Butter tree flower (*Madhuca longifolia*) which are rich in phytochemical properties but still not widely explored for the applicability in the preparation of herbal beverages. Utilization of tree flowers in herbal beverages is restricted because of the short blooming period and availability in a limited geographical region. Furthermore, due to the short post-harvest life, these flowers are utilized mostly by the local tribes as a food ingredient or for medicinal purposes only during their respective flowering times [16].

#### 4. HERBAL BEVERAGE FORMULATIONS WITH EDIBLE FLOWERS

Edible flowers provide competitive advantages to the production of herbal beverages. Their distinctive flavors, vivid pigments, and inherent medicinal properties make them an excellent addition to various formulations. By incorporating edible flowers and their extracts, alongside other compatible ingredients, a diverse array of visually stimulating herbal beverages can be developed to cater to both gustatory and sensory experiences. These flowers harbor valuable bioactive compounds that hold the potential to positively impact overall well-being. Moreover, the integration of edible flowers into herbal beverages amplifies their phytochemical composition, thereby contributing to enhanced health benefits.

A functional beverage incorporated with *Clitoria ternatea* flower extract was developed with two other constituents, natural sweetener (Stevia extract) and a flavor (lime). This functional beverage possessed a significantly higher ( $p < 0.05$ ) antioxidant potential with respect to total phenolic content and oxygen radical absorbance capacity compared to pure *Clitoria ternatea* extract. The total flavonoid content of *Clitoria ternatea* extract incorporated functional beverage & extract was 43.67 mg quercetin equivalents (QE)/ liter of sample & 42.75 mg QE/L of sample respectively. Antioxidant activity of the functional beverage was quantified by ferric reducing antioxidant power, DPPH radical scavenging activity, and ABTS radical scavenging activity resulting values were 14.99 mg Trolox equivalents (TE) per liter of sample, 35.07 mg TE/L, and 185.81 mg TE/L respectively. As per the evaluation of quality parameters of the functional natural beverage it was concluded that the beverage can be effectively utilized to manage oxidative stress associated with chronic diseases [17].

“Kombucha” is a traditional fermented beverage that is made commonly using black tea, green tea, and oolong tea. However, a study was conducted replacing the original raw material with a *Clitoria ternatea* infusion as the base material in the preparation of Kombucha. Variations in Total flavonoid content & antioxidant activity of Butterfly pea Kombucha were examined. Total flavonoid content indicated a marked growth over the fermentation period, however, antioxidant activity values remained almost stable [18].

Powder drink from Butterfly pea flower was developed using co-crystallization, agglomeration, and drying methods by Marpaung et al.[19] and the prepared drink was evaluated over 28 days at three different storage temperatures. (27<sup>o</sup> C, 40<sup>o</sup> C, and 50<sup>o</sup> C). Although a significant decrease in the stability of Anthocyanins was seen as the storage temperature increased, the decrease in antioxidant activity of the powder drink was minimal.

More than 20,000 Chrysanthemum cultivars are found in the world and about 7000 cultivars are present in China [20]. Chrysanthemum tea is one of the most popular herbal infusions widely consumed mainly in China & Japan. Chrysanthemum flower with yellow petals is frequently used as a tea material, a coolant, and an antioxidant in the Korean herbal tea industry.

Two types of Chrysanthemum flower teas are produced and commercialized in China. Tea prepared from flower buds is named "Taiju" and another type of tea prepared from blooming flowers is named as "Hangbaiju". Keneko et al. [21] carried out an experiment using the aroma extract dilution analysis (AEDA) technique to identify the compounds responsible for characteristic floral/sweet odor in Chinese Chrysanthemum flower tea infusion. Studies revealed that flower buds are rich in four compounds (verbenone, ethyl 3-phenylpropanoate, propyl 3-phenylpropanoate, and ethyl cinnamate) which give four floral/sweet aroma peaks.

However, to develop chrysanthemum teas in different appealing colors and fragrances suitability of the new flower tea cultivar with a dark Purple petal over the commercial Yellow Chrysanthemum cultivar was evaluated under different infusing time and temperature conditions. Both cultivars exhibited strong ABTS radical scavenging activity and moderate DPPH radical scavenging activity. However Purple Chrysanthemum tea contained anthocyanins and linamarin, which were not detected in the Yellow Chrysanthemum tea and the content of chlorogenic acid, acacetin-7-O- $\beta$ -glucoside, and luteolin was higher compared to the Yellow Chrysanthemum tea. On the other hand, Yellow Chrysanthemum tea had higher luteolin-7-O- $\beta$ -glucoside, 3, 5-dicaffeoylquinic acid, apigenin-7-O- $\beta$ -glucoside, and apigenin contents compared to the Purple Chrysanthemum tea [22].

Tea aroma formation is a novel aspect in herbal beverage manufacturing to enhance the sensory properties. Therefore, tea volatiles generation and retention, irrespective of the rigorous treatment conditions are important to deliver herbal teas with the finest quality. It was concluded that dry tea could adsorb substantial amounts of volatiles from jasmine flowers to improve the aroma profile of scented green tea [23].

Chamomile tea is one of the most popular herbal beverages all around the world and approximately a million cups of chamomile tea are consumed each day. German Commission E recommends the oral consumption of herbal infusions derived from Chamomile. Chamomile flowers can be used in the preparation of herbal beverages in their pure form after blending with other medicinal herbs. It is available as a powder and dried flower as a whole or flakes in tea bags. Chamomile tincture is prepared by mixing one part of chamomile flower in four parts of water having 12% grain alcohol. Herbal beers are made from the whole herb including the Chamomile flower. Chamomile wine, another derivative is prepared by adding dried Chamomile flowers to White wine and then followed by steeping and filtration [24, 25].

Lotus pollens are gaining much attention as an ingredient in natural health supplements. They have exhibited antioxidant properties when brewed in hot water. Lotus pollen tea was chosen as a viable consumer product considering the high pollen yield, potent antioxidant activity, favorable aroma, flavor, and color of *Nelumbo nucifera* tea [26]. Out of the four varieties of lotus evaluated (Sattabongkutt, Buntharik, Sattabutt, and Patum), Buntharik

reported the highest antioxidant capacity at a trolox equivalent of 499.48 µg/ml and total polyphenols equivalent of gallic acid of 63.26 µg/ml. The stamen of *Nelumbo nucifera* is used in dried or powdered form in herbal teas and as a main ingredient in traditional remedies in Asian countries such as China, Thailand, Japan, and India [27].

Rose petal teas are considered a caffeine-free beverage with high antioxidant capacity. Yakov et al. [28]. studied the antioxidant activity and total phenolic contents of Rose petal tea. Hot water infusions prepared from air-dried petals of twelve different Rose cultivars were analyzed and the study concluded that teas from different Rose cultivars possess significantly different sensory properties, and they are accepted to be used as an herbal beverage alone or in combination with other herbal materials.

Edible flowers can be used with other plant-based ingredients to produce herbal beverages with improved palatability other than using them alone in beverage preparations. To evaluate how different concentration levels (1.5%, 3%, 4%) of natural flavors (mint leaves, ginger, and lemon juice) affect the sensory and functional properties of the rose-extract-based drink. Water extract of Rose petals was obtained by maceration with 1% citric acid. Rose extract was concentrated by evaporating the filtrate of water extract in a rotary evaporator at 50-60<sup>0</sup> C until the desired volume was achieved. The functional beverage was produced by mixing the prepared rose extract, natural flavor materials, citric acid, and sugar and heating at 80-85<sup>0</sup>c. Before filling the mixture was cooled up to 60-65<sup>0</sup>C. A marked increase in the total anthocyanin content and antioxidant activity resulted from adding natural flavoring ingredients. The addition of mint leaves, ginger, and lemon increased the total anthocyanin content by 46.48%, 34.50%, and 16.07% respectively. Rose extract-based drink flavored with 3% mint leaves which gave 84.44% antioxidant activity and favorable flavor properties was selected as the best formulation [29].

Sylvia et al. [30] developed a functional fruit-herbal beverage and evaluated the physiochemical properties and consumer acceptance of the beverage. Fruit –herbal beverage was prepared with fruit juices (80%) obtained from Aronia (*Aronia melanocarpa Michx.*), Rugosa Rose (*Rosa rugosa Thumb.*), Acerola (*Malpighia glabra L.*), Sea buckthorn (*Hippophaë rhamnoides L.*), and Cranberry (*Oxycoccus palustris Pers.*, *Vaccinium Oxycoccus L.*) and supplemented with herbal extracts prepared from different combinations of Nettle leaves (*Urtica dioica*), Siberian ginseng (*Eleutherococcus senticosus*), Ginger (*Zingiber officinale*), Purple Coneflower (*Echinacea purpurea*), Aloe (*Aloe vera*), Horsetail (*Equisetum arvense*), Lingonberry (*Vaccinium vitis-idaea*), Silver birch (*Betula pendula*), and Chamomile (*Matricaria chamomilla*) (20%) in different combinations. The juice mixture sweetened with sucrose was heated up to 80<sup>0</sup>C, filtered, and then pasteurized at 85–87<sup>0</sup> C for 10 minutes. The study concluded that all the developed fruit herbal beverages showed a high antioxidant potential and were accepted by consumers in sensory evaluations. Highest carotenoid content, polyphenols content (807 mg GA/L), vitamin C content (3439 mg/l), and antioxidant activity (DPPH assay: 75.7 µM Trolox/g and ABTS assay: 63.2 µM Trolox/g) was reported in the formulation made with Rosehip–acerola, Siberian ginseng, Ginger, Purple Coneflower, and Aloe.

Oluwatoyin et al. [31] studied the synergistic effect of fruit/vegetable (Pineapple/Carrot/Orange) based beverages blended with *Hibiscus sabdariffa* extracts (HSE). The objective of the study was to formulate a functional beverage with optimized antioxidant properties. Juice formulation with 40% Pineapple, 16.5% Carrot, 17.2% Orange, and 26.3% HSE extract reported total phenolic content as 512.82 mg GAE 100 g<sup>-1</sup>, Vitamin C content as 3.37 mgg<sup>-1</sup>, and ability to inhibit DPPH as 51.34% and found to be the blend with optimum nutraceutical quality and sensory properties. Blends with increased ratios of

HSE exhibited the highest antioxidant activity and the study concluded all the ingredients used in the experiment are accepted to be utilized in herbal beverage preparations.

Another low-caloric beverage with high antioxidant capacity which can be used as an alternative option for popular industrialized beverages was produced and examined by Elizabeth et al. [32] This plant-based beverage included aqueous extracts of medicinal plants: Melissa (*Melissa officinalis*) and Lemon verbena (*Aloysia triphylla*), flowers: Chamomile (*Matricaria chamomilla*) and Bougainvillea (*Bougainvillea glabra*), fruits: (Guava, Pineapple, Strawberry, Orange, and Tangerine). Experimental results indicated that flower bougainvillea improved the sensory quality of the product by providing an appealing colour along with improved bioactive properties to the beverage.

A mixed beverage was prepared from Hibiscus and Coconut water using three different combinations, HA 1 (50% Hibiscus and 50% Coconut water), HA 2 (70% Hibiscus and 30% Coconut water), and HA 3 (90% Hibiscus and 10% Coconut water). Three mixes were pasteurized at 90°C for 3 min, hot filled into glass bottles, and stored at refrigeration temperature (4°C) for 35 days. Physicochemical, microbiological, sensory, and phytochemical properties of the mixed beverage were assessed throughout the storage period. The highest retention of total phenolic compounds (decreased from 177.74 to 153.71 mg GAE/g) and antioxidant activity (reduced from 174.23 to 20.81 µm Ferrous sulfate/g) was achieved in the HA3 beverage mix. Processing temperatures applied were capable of inactivating microorganisms and retardation of polyphenol oxidase enzymatic activity for 35 days. Most acceptable chemical and sensory features were achieved from the blend HA2, hence concluded to be the most viable formulation in the development of herbal beverages [33].

Dabbagh et al. [34] developed a Saffron-based probiotic beverage fermented by lactic acid bacteria. Air-dried Saffron petals were powdered, and water extract was prepared. Then it was sweetened with a mixture of sucrose, glucose, and fructose before sterilization at 121°C for 15 minutes. Then the prepared beverage was fermented by probiotic strains including *Lactococcus lactis*, *Lactobacillus plantarum*, *Lactobacillus brevis*, and *Lactobacillus casei*. Total phenolic compounds, DPPH scavenging capacity, and total monomeric anthocyanin of fermented saffron beverages ranged from 69.1-99.1 mg GAE/g, 105.7- 160.6 EC50 mg/100 g mass, 5.6-12.1 malvidin-3-O-glucoside equivalents/g dry sample respectively. Unfermented saffron beverage contained 51.8 mg GAE/g of total phenolic compounds and 18.2 malvidin-3-O-glucoside equivalents/g of total monomeric anthocyanins. Therefore, the study concluded that saffron-based fermented beverage is accepted as a health-promoting non-dairy beverage.

Ashwani et al. [35] conducted a study to investigate the physicochemical, phytochemical, sensorial, and storage properties of a cereals and milk-based functional beverage supplemented with Marigold powder and Rose Syrup Functional beverage produced from finger millet, oats, and double-toned milk. Marigold powder was prepared by grinding dehydrated flower petals at 40°C in a hot air cabinet dryer. Rose syrup, also known as rose petal preserves, was produced by macerating fresh rose petals with sugar for 2 days at room temperature and then heating the solution at 70-100°C for 1 hour until the complete dissolution of sugar to optimize the extraction of colour and flavor.

The bioactive potential of the functional beverage was significantly improved by blending with flower extracts. Anthocyanins & total phenolic content significantly increased by 78.82 & 119.18% in the beverage mixed with marigold powder and by 230.58% & 145.23% in the Rose syrup-supplemented functional beverage respectively. Beta carotene (0.37 mg/100 ml) was detected only in the functional beverage blended with Marigold powder. In Rose syrup supplemented beverage 12.65 % of DPPH radical scavenging activity was observed while

Marigold powder supplemented beverage reported 10.46 % of DPPH radical scavenging activity. Functional beverage supplemented with Rose syrup was the most acceptable formulation in sensory analysis. At refrigerated temperature shelf stability of the developed beverage was limited to 50 days [36].

A novel tea infusion was prepared with dried *Hibiscus calyces* petal powder, Ginger powder, and Cocoa powder mixed in different ratios and was analyzed by Folashade et al. [37] Beverage powder was packed in tea bags with an average weight of 1.5 g. To prepare the tea by hot water extraction tea blend was infused at 70°C for 1 minute and to prepare by cold water extraction tea bags were dipped in distilled water at 25°C for 5 minutes. Infusions prepared from both extraction methods had total phenolic content varied from 11.61 to 22.01 mg GAE/g; ferric reducing activity varied from 10.84 to 26.88 mg/g, vitamin C content varied from 1.77 to 4.82 mg/g and DPPH inhibition varied from 38.60 to 87.53%. The study concluded that the beverage blend has complimentary effects and beneficial antioxidant properties.

Rodrigues et al. [38] coupled Sea lavender (*Limonium algarvense*) and Green tea (*Camellia Sinensis*) to produce an herbal beverage with improved inhibitory properties towards enzymes related to Alzheimer's (acetyl- and butyrylcholinesterase) and type-2 diabetes mellitus ( $\alpha$ -amylase and  $\alpha$ -glucosidase). Powders obtained from dried plant material were mixed in different combinations to prepare the infusions for further analysis. Both the plant species are rich in antioxidant properties, however, the experimental results concluded that when they are coupled together radical-scavenging and anti-lipid peroxidation activities of the prepared herbal beverage improved synergistically. Although Lavender had a stronger inhibition capacity regarding anti-cholinesterase activity compared to Green tea, the herbal mixture exhibited a greater synergistic effect. Green tea is rich in epicatechin and caffeic acid was the main contributor to  $\alpha$ -glucosidase inhibition, but the herbal blend induced an antagonistic effect.

## **5. PHYTOCHEMICALS AND PHARMACOTHERAPEUTIC POTENTIAL OF EDIBLE FLOWERS IN HERBAL BEVERAGES**

The prime advantage of incorporating edible flowers into herbal beverages lies in their abundance of bioactive compounds, encompassing carotenoids, phenolic acids, flavonoids, coumarins, alkaloids, polyacetylenes, saponins, and terpenoids [39]. Consumer interest is shifting towards safe natural origin products other than relying on conventional pharmaceuticals to cure the pathologic conditions negatively affecting the health status [40]. Furthermore, recent research explored the synergistic potential of phytochemicals found in herbal beverages on effective management of medical conditions when used parallel to conventional medicines [41].

Edible flowers deliver a plethora of health benefits for consumers, offering antioxidant, antibacterial, antiviral, anti-inflammatory, antifungal, antidiabetic, neuroprotective, antiallergic, antithrombotic, and vasodilatory actions. Additionally, they exhibit antimutagenic, anticarcinogenic, and anti-aging effects, making them valuable contributors to overall well-being [4, 42]. Both in vitro and in vivo studies are carried out to assess their pharmacotherapeutic properties, clinical efficacy, and safety concerns related to toxicity and microbiological quality. Table 1 presents a concise overview of the essential bioactive compounds and therapeutic effects found in common edible flowers, often utilized in the preparation of herbal beverages.

**Table 1. Major bioactive compounds & pharmacotherapeutic properties of edible flowers used in herbal beverages.**

<b>Crop name</b>	<b>Major bioactive compounds</b>	<b>Pharmacotherapeutic properties</b>	<b>Ref</b>
<i>Hibiscus calyces</i>	Polyphenols (Delphinidin 3-sambubioside, 3-caffeoylquinic acid) Anthocyanins (Cyanidin-3-O-sambubioside, delphinidin-3-O-sambubioside)	Antibacterial, antioxidant, nephro- and hepato-protective, renal/diuretic, anti-diabetic and anti-hypertensive effects, lowering blood pressure in pre and mildly hypertensive adults, lowering blood pressure in type II diabetic patients with mild hypertension, increasing antioxidant capacity of both the enzymatic and non-enzymatic systems in the plasma of the MSF (Marfan syndrome) patients, reducing total cholesterol, low-density lipoprotein-cholesterol (LDL-C), triglycerides, increasing high-density lipoprotein-cholesterol (HDL-C).	[43-58][59]
<i>Rosa spp.</i>	Anthocyanins (Cyanidin 3, 5-di-O-glucoside, peonidin 3-O-sophoroside, peonidin 3, 5-di-O-glucoside, peonidin 3-O-glucoside), Carotenoids (Luteoxanthin, violaxanthin, zeaxanthin, $\beta$ -Carotene, lutein epoxide, lutein, antheraxanthin, neoxanthin), Flavanols (Kaempferol 3-O-rhamnoside, quercetin 3-O-glucosid)	Antioxidants, anticancer, anti-inflammatory, antimutagenic, and antidepressant properties, moderate anti-HIV activity, anti-viral, anti-diabetic activities, antifungal properties, neuropharmacological, alleviate constipation.	[60-70]
<i>Nelumbo nucifera</i>	Flavonoids (Anthocyanin, myricetin 3-O-glucoside, quercetin, kaempferol, isohamnetin, diosmetin)	Strong antibacterial activity against <i>Escherichia coli</i> , <i>Bacillus Subtilis</i> , <i>Staphylococcus aureus</i> and moderate inhibition against <i>Klebsiella pneumonia</i> and <i>Pseudomonas aeruginosa</i> , prevention of degenerative diseases, treating bleeding disorders, fever, diarrhea, hyperdipsia, cholera, and hepatopathy, improving circulatory system, decrease blood glucose and lipid levels, treatment of diabetes mellitus.	[71-77]
<i>Chrysanthemum morifolium</i>	Carotenoids (Lutein, zeaxanthin, $\beta$ -cryptoxanthin, 13-cis- $\beta$ -carotene, $\alpha$ -carotene, trans- $\beta$ -carotene, and 9-cis- $\beta$ -carotene.)	Antioxidant, anti-inflammatory, antitumor, antiosteoporotic, effects, easing stress and anxiety, protection against oxidative damage, enhancing the immune function, improving eye health, cardiovascular health, dispelling pathogenic heat, improving liver function and detoxification, antiviral activities against Herpes simplex virus type 1 (HSV 1).	[22, 78-81]

<i>Calendula officinalis</i> L.	Carotenoids (Lutein, Lycopene, beta carotene, Zeaxanthin, rubixanthin, flavoxanthin, neoxanthin, auroxanthin, luteoxanthin and 9Z-antheraxanthin, violaxanthin), terpenoids and terpenes (bisabolol, faradiol, chamazulene, arnidiol and esters), flavonoids, (quercetin, isorhamnetin and kaempferol aglycones) and polyunsaturated fatty acids, (calendic acid), Coumarins (scopoletin, umbelliferone, esculetin)	Angiogenic, vascular regeneration, analgesic, antibacterial, antiprotozoal activity, antifungal, antioxidant and immunomodulatory effects, wound healing properties, anti-inflammatory, cytotoxic, antiseptic, anti-viral, hepatoprotective, antidiabetic, anthelmintic activity.	[82-86]
<i>Magnolia grandiflora</i> L.	Flavanols (Kaempferol, quercetin, isorhamnetin) alkaloids, sesquiterpenes, phenolic constituents, glycosides)	Treatment of diarrhea, abdominal diseases, rheumatic arthritis, heart disturbances, high blood pressure, epilepsy, infertility and fever, anti-inflammatory properties, anti-melanogenic and antioxidant properties	[87, 88]
<i>Clitoria ternatea</i>	Polyphenols (ternatin anthocyanins), flavanol glycosides of kaempferol, rutin, quercetin, and myricetin,	Antimicrobial, antioxidant, anti-inflammatory, anti-diabetic properties, tranquilizing effect, antipyretic activities, anti-proliferative properties inhibiting cancer cell lines, diuretic, nootropic, anti-asthmatic, analgesic, antilipidemic, anti-arthritis, antioxidant, and wound-healing properties.	[89-95]
<i>Matricaria chamomilla</i>	Terpenoids (sesquiterpenes and $\alpha$ bisabolol), flavonoids (apigenin, quercetin, patuletin, and lute, coumarins (herniarin and umbelliferoneolin), cinnamic acid derivatives (ferulic acid and caffeic acid)	Antioxidant, antimicrobial, antidepressant, anti-inflammatory, antidiarrheal activities, angiogenesis activity, anti-carcinogenic, hepatoprotective, and antidiabetic properties and beneficial impacts on knee osteoarthritis, ulcerative colitis, premenstrual syndrome, and gastrointestinal disorders, wound healing properties, protective effect on hematological parameter disorders, erythrocytes-induced oxidative stress after the acute alcohol administration, neuroprotective agent against oxidative brain damage, anti-allergic activity, alleviating depression and sleep quality problems in post-partum women, beneficial effects on glycemic control and serum lipid profile in type 2 diabetes mellitus patients, relieving the intensity of premenstrual syndrome associated symptomatic psychological pains.	[96-106]

## 5. SAFETY ASPECTS OF THE CONSUMPTION OF EDIBLE FLOWERS

Although there are many edible flowers rich in phytonutrients and safe for human consumption, few of them are not recommended for consumption. Some edible flowers contaminated by dimethate, sulphite, and pathogenic microorganisms (*Salmonella spp*) have been identified by Rapid Alert System for Food and Feed (RASFF) during the last years [107]. Hazardous agents may originate from the plant itself or at the various steps in the food chain.

Pathogenic bacteria and residues from agricultural production are the two main external impurities found in edible flowers. The microbiological quality of the edible flowers depends on several factors such as soil quality, irrigation waters, fertilization, crop protection techniques, and harvesting equipment. Subsequently, after the harvesting, edible flowers may be contaminated during processing, transport, and distribution, by processing water or ice, and due to lack of personal hygiene of the food handler [108]. In an experiment carried out by Aleksandra et al.[109] microbial quality of 5 selected edible flowers namely Nasturtium (*Tropaeolum L.*), Calendula (*Calendula officialis L.*), Daylily (*Hemerocallis L.*), Daisy (*Bellis*), Pink (*Dianthus caryophyllus L.*) were assessed. At the end of the experiment, it was concluded none of the flowers were contaminated by *Salmonella sp.*, but *Staphylococcus aureus* was detected in all flower samples at  $>3 \log \text{CFU g}^{-1}$ . Total yeast and mould counts were less than  $6 \log \text{CFU g}^{-1}$  and *E. coli* was detected only in Nasturtium flowers.

Toxic compounds and their possibility to cause allergic reactions are one of the main risks associated with the consumption of edible flowers. The availability of toxic compounds depends on factors such as the type of flower, the cultivation of the flowers, and the presence of toxic compounds in the floral tissues. Edible flowers may contain phytotoxins and ingestion of them can lead to health implications such as nausea, vomiting diarrhea, and even DNA damage in exceptional situations. Thujone, three alkenylbenzenes-methyl eugenol, 1, 8 cineole (eucalyptol), and safrole are the three phytotoxins frequently reported in edible flowers [110].

In a study conducted by Akindahunsi et al. [111]wistar albino rats were orally administrated with an aqueous fraction of an aqueous-alcoholic extract of *Hibiscus sabdariffa L. calyces* to examine the toxicological impact. Results of the study demonstrated that prolonged usage of 15 doses of 250 mg/kg of this extract caused liver injury and mild effects were given at 1-10 dose levels. Also, the average consumption of 150–180 mg/kg per day was considered to be safe. Another study conducted by Sireeratawong et al. [112] revealed that at the doses of 50, 100, and 200 mg/kg body weight for 270 days, *H. sabdariffa* extract did not produce chronic toxicity in rats. In the amount of 5,000 mg/kg body weight oral administration of a single dose of *H. sabdariffa* extract also did not generate acute toxicity in rats.

Akbari M et al. [113] assessed the toxic effects of *Rosa damascena* using an animal model using exponential doses of the infusion made with petals (90-1440 mg/kg/d) over a period of 10 days. The Study of the results concluded that minimal nephrotoxic or hepatotoxic effects.

A chronic toxicity study was carried out by feeding *Chrysanthemum morifolium* extract to rats at dose levels of 320, 640, and 1280 mg/kg body weight /day for 26 weeks. No toxicological changes in body weight, hematologic examination, blood chemical examination, organ weight, and microscopic histopathologic examination were reported in rats at the end of the study [114].

The concentration of metals and heavy metals in tissues is another risk associated with the usage of edible flowers in food applications. However, the possibility for metal poisoning as a result of the consumption of edible flowers is low hence edible flowers contain a relatively low concentration of metals and heavy metals (Cd, Co, Ni, Pb, and V) [115, 116].

Edible flowers harvested from wild plants may pose certain safety concerns. One potential issue is the risk of contamination with toxic plant species that grow near the edible ones. Additionally, wild plants, especially those in non-agricultural areas, may accumulate chemical impurities from the soil or airborne pollutants. Certain regions may suffer from high concentrations of heavy metals in the soil, which can be absorbed by these plants, leading to potential safety risks when consumed by humans. Furthermore, impurities from the air, such as those arising from automotive or industrial pollution, can also affect the safety of wild edible flowers. Therefore, the location where these flowers are harvested may play a significant role in determining their safety for consumption [108]. While numerous studies have been carried out to assess the safety aspects of edible flowers, there remains a need for more extensive research concerning the presence of hepatotoxic pyrrolizidine alkaloids in both nectar and pollen [117].

## 6. CONCLUSION

Although a wide array of edible flowers exists in various regions globally, only a handful of them have been subject to thorough investigation and exploration. It is required to conduct extensive research in this field to unveil the health advantages of these underutilized or undiscovered edible flowers. These findings can be leveraged for the development of functional beverages aimed at mitigating the risk of non-communicable diseases. Furthermore, forthcoming research should center on the bioavailability and bio accessibility of edible flowers subjected to processing and preservation methods during the preparation of herbal beverages.

## REFERENCES

1. Vilela A, Pinto T. Grape Infusions: The Flavor of Grapes and Health-Promoting Compounds in Your Teacup. *Beverages*. 2019;5(3):48.
2. Izzo AA, Hoon-Kim S, Radhakrishnan R, Williamson EM. A Critical Approach to Evaluating Clinical Efficacy, Adverse Events, and Drug Interactions of Herbal Remedies. *Phytother Res*. 2016;30(5):691-700.
3. Poswal FS, Russell G, Mackonochie M, MacLennan E, Adukwu EC, Rolfe V. Herbal Teas and their Health Benefits: A Scoping Review. *Plant Foods Hum Nutr*. 2019;74(3):266-76.
4. Chandrasekara A, Shahidi F. Herbal beverages: Bioactive compounds and their role in disease risk reduction - A review. *J Tradit Complement Med*. 2018;8(4):451-8.
5. Jäger S, Beffert M, Hoppe K, Nadberezny D, Frank B, Scheffler A. Preparation of herbal tea as infusion or by maceration at room temperature using mistletoe tea as an example. *Sci Pharm*. 2011;79(1):145-55.
6. Suna S, Tamer CE, Özcan-Sinir G. Trends and Possibilities of the Usage of Medicinal Herbal Extracts in Beverage Production. *Natural Beverages*. 2019.
7. Guimarães R, Barros L, Carvalho AM, Ferreira IC. Infusions and decoctions of mixed herbs used in folk medicine: synergism in antioxidant potential. *Phytother Res*. 2011;25(8):1209-14.
8. Malik J, Szakova J, Drabek O, Balik J, Kokoska L. Determination of certain micro and macroelements in plant stimulants and their infusions. *Food Chemistry*. 2008;111(2):520-5.

9. Ni H, Jiang Q-X, Zhang T, Huang G-L, Li L-J, Chen F. Characterization of the Aroma of an Instant White Tea Dried by Freeze Drying. *Molecules*. 2020;25(16):3628.
10. Mason TJ, Yiyun Z. Enhanced extraction of tea solids using ultrasound. *Ultrasonics*. 1994;32(5):375-7.
11. Gruenwald J. 5 - Fortification of beverages with products other than vitamins and minerals. In: Paquin P, editor. *Functional and Speciality Beverage Technology*: Woodhead Publishing; 2009. p. 92-106.
12. Pratt DE. Natural Antioxidants from Plant Material. *Phenolic Compounds in Food and Their Effects on Health II*. ACS Symposium Series. 507: American Chemical Society; 1992. p. 54-71.
13. Rodrigues H, Cielo DP, Gómez-Corona C, Silveira AAS, Marchesan TA, Galmarini MV, et al. Eating flowers? Exploring attitudes and consumers' representation of edible flowers. *Food Research International*. 2017;100:227-34.
14. Lu B, Li M, Yin R. Phytochemical Content, Health Benefits, and Toxicology of Common Edible Flowers: A Review (2000-2015). *Crit Rev Food Sci Nutr*. 2016;56 Suppl 1:S130-48.
15. Mlcek J, Rop O. Fresh edible flowers of ornamental plants – A new source of nutraceutical foods. *Trends in food science & technology*. 2011;22(10):561-9.
16. Pinakin DJ, Kumar V, Suri S, Sharma R, Kaushal M. Nutraceutical potential of tree flowers: A comprehensive review on biochemical profile, health benefits, and utilization. *Food Res Int*. 2020;127:108724.
17. Lakshan SAT, Jayanath NY, Abeysekera W, Abeysekera W. A Commercial Potential Blue Pea (*Clitoria ternatea* L.) Flower Extract Incorporated Beverage Having Functional Properties. *Evid Based Complement Alternat Med*. 2019;2019:2916914.
18. Dwiputri MC, Feroniasanti YML. Effect of Fermentation to Total Titrable Acids, Flavonoid and Antioxidant Activity of Butterfly Pea Kombucha. *Journal of Physics: Conference Series*. 2019.
19. Marpaung AM, Lee M, Kartawiria IS, editors. *The Development of Butterfly pea (Clitoria ternatea) Flower Powder Drink by Co-crystallization*2020.
20. Wang F, Zhang F-J, Chen F-D, Fang W-M, Teng N-J. Identification of *Chrysanthemum* (*Chrysanthemum morifolium*) Self-Incompatibility. *The Scientific World Journal*. 2014;2014(1):625658.
21. Kaneko S, Chen J, Wu J, Suzuki Y, Ma L, Kumazawa K. Potent Odorants of Characteristic Floral/Sweet Odor in Chinese *Chrysanthemum* Flower Tea Infusion. *J Agric Food Chem*. 2017;65(46):10058-63.
22. Han AR, Kim HY, So Y, Nam B, Lee IS, Nam JW, et al. Quantification of Antioxidant Phenolic Compounds in a New *Chrysanthemum* Cultivar by High-Performance Liquid Chromatography with Diode Array Detection and Electrospray Ionization Mass Spectrometry. *Int J Anal Chem*. 2017;2017:1254721.
23. Shen J-X, Rana MM, Liu G-F, Ling T-J, Gruber MY, Wei S. Differential Contribution of Jasmine Floral Volatiles to the Aroma of Scented Green Tea. *Journal of Food Quality*. 2017;2017:5849501.
24. Srivastava JK, Shankar E, Gupta S. Chamomile: A herbal medicine of the past with bright future. *Mol Med Rep*. 2010;3(6):895-901.
25. Srivastava JK, Gupta S, editors. *Health Promoting Benefits of Chamomile in the Elderly Population*2009.
26. Hongsamoot D, Bumroongsook S. A FEASIBILITY STUDY ON ORGANIC LOTUS POLLEN TEA AS CONSUMER PRODUCT. *Emirates Journal of Food and Agriculture*. 2015;27(5):441-6.
27. Tungmunnithum D, Renouard S, Drouet S, Blondeau JP, Hano C. A Critical Cross-Species Comparison of Pollen from *Nelumbo nucifera* Gaertn. vs. *Nymphaea lotus* L. for Authentication of Thai Medicinal Herbal Tea. *Plants (Basel)*. 2020;9(7).

28. Vinokur Y, Rodov V, Reznick N, Goldman G, Horev B, Umiel N, et al. Rose petal tea as an antioxidant-rich beverage : Cultivar effects. *Journal of Food Science*. 2006;71:23.
29. Saati EA, Ramadhan AH, Lutfi M, Wahyudi VA, Manshur HA. Utilization of rose flower extract as antioxidant rich-drink. *IOP Conference Series: Earth and Environmental Science*. 2020;458(1):012035.
30. Skąpska S, Marszałek K, Woźniak Ł, Szczepańska J, Danielczuk J, Zawada K. The Development and Consumer Acceptance of Functional Fruit-Herbal Beverages. *Foods*. 2020;9(12):1819.
31. Ogundele OM, Awolu OO, Badejo AA, Nwachukwu ID, Fagbemi TN. Development of functional beverages from blends of Hibiscus sabdariffa extract and selected fruit juices for optimal antioxidant properties. *Food Sci Nutr*. 2016;4(5):679-85.
32. Contreras-López E, Ramírez-Godínez J, García-Martínez MM, Gutiérrez-Salomón AL, González-Olivares LG, Jaimez-Ordaz J. Low-Calorie Beverages Made from Medicinal Plants, Flowers and Fruits: Characteristics and Liking of a Population with Overweight and Obesity. *Applied Sciences*. 2021;11(9):3766.
33. Carvalho de Castro JM, Nascimento Alves CA, de Lima Santos K, de Oliveira Silva E, Maria da Silva Araújo Í, Barros de Vasconcelos L. Elaboration of a mixed beverage from hibiscus and coconut water: An evaluation of bioactive and sensory properties. *International Journal of Gastronomy and Food Science*. 2021;23:100284.
34. Dabbagh Moghaddam A, Garavand F, Razavi SH, Dini Talatappe H. Production of saffron-based probiotic beverage by lactic acid bacteria. *Journal of Food Measurement and Characterization*. 2018;12:2708-17.
35. Kumar A, Kaur A, Tomer V, Gupta K, Kaur K. Effect of Rose Syrup and Marigold Powder on the Physicochemical, Phytochemical, Sensorial and Storage Properties of Nutricereals and Milk-Based Functional Beverage. *Journal of the American College of Nutrition*. 2021;40(2):133-40.
36. Ashwani K AK, Joshi VK. *Rosa damascena* : Quality Evaluation and Process Optimization for the Development of Rose Syrup. *Intl J Food Ferment Technol* 2017;7(2):279-85.
37. Awe FB, Fagbemi TN, Ifesan BO, Badejo AA. Antioxidant properties of cold and hot water extracts of cocoa, Hibiscus flower extract, and ginger beverage blends. *Food Research International*. 2013;52:490-5.
38. Rodrigues MJ OMBPP, Neves V, Ovelheiro A, Pereira CA, Neng NR  
Coupling Sea Lavender (*Limonium Algarvense* Erben) and Green Tea (*Camellia Sinensis* (L.) Kuntze) to Produce an Innovative Herbal Beverage with Enhanced Enzymatic Inhibitory Properties. *S Afr J Bot*. 2019;120:87-94.
39. Kumari P, Ujala, Bhargava B. Phytochemicals from edible flowers: Opening a new arena for healthy lifestyle. *Journal of Functional Foods*. 2021;78:104375.
40. Daliu P, Santini A, Novellino E. A decade of nutraceutical patents: where are we now in 2018? *Expert Opinion on Therapeutic Patents*. 2018;28(12):875-82.
41. Malongane F, McGaw LJ, Mudau FN. The synergistic potential of various teas, herbs and therapeutic drugs in health improvement: a review. *J Sci Food Agric*. 2017;97(14):4679-89.
42. B. Prabawati N, Oktavirina V, Palma M, Setyaningsih W. Edible Flowers: Antioxidant Compounds and Their Functional Properties. *Horticulturae*. 2021;7(4):66.
43. Piovesana A, Rodrigues E, Noreña CPZ. Composition analysis of carotenoids and phenolic compounds and antioxidant activity from hibiscus calyces (*Hibiscus sabdariffa* L.) by HPLC-DAD-MS/MS. *Phytochem Anal*. 2019;30(2):208-17.
44. Grajeda-Iglesias C, Figueroa-Espinoza MC, Barouh N, Baréa B, Fernandes A, de Freitas V, et al. Isolation and Characterization of Anthocyanins from *Hibiscus sabdariffa* Flowers. *J Nat Prod*. 2016;79(7):1709-18.
45. Riaz G, Chopra R. A review on phytochemistry and therapeutic uses of *Hibiscus sabdariffa* L. *Biomed Pharmacother*. 2018;102:575-86.

46. Ali BH, Al Wabel N, Blunden G. Phytochemical, pharmacological and toxicological aspects of *Hibiscus sabdariffa* L.: a review. *Phytother Res.* 2005;19(5):369-75.
47. Da-Costa-Rocha I, Bonnlaender B, Sievers H, Pischel I, Heinrich M. *Hibiscus sabdariffa* L. - a phytochemical and pharmacological review. *Food Chem.* 2014;165:424-43.
48. Etheridge CJ DE. *Hibiscus* Tea and Health: A Scoping Review of Scientific Evidence. *Nutr Food Technol.* 2020;6(2).
49. Pepe G, Giusti B, Sticchi E, Abbate R, Gensini GF, Nistri S. Marfan syndrome: current perspectives. *Appl Clin Genet.* 2016;9:55-65.
50. Soto ME, Zuñiga-Muñoz A, Guarner Lans V, Duran-Hernández EJ, Pérez-Torres I. Infusion of *Hibiscus sabdariffa* L. Modulates Oxidative Stress in Patients with Marfan Syndrome. *Mediators Inflamm.* 2016;2016:8625203.
51. McKay DL, Chen CY, Saltzman E, Blumberg JB. *Hibiscus sabdariffa* L. tea (tisane) lowers blood pressure in prehypertensive and mildly hypertensive adults. *J Nutr.* 2010;140(2):298-303.
52. Herrera-Arellano A, Flores-Romero S, Chávez-Soto MA, Tortoriello J. Effectiveness and tolerability of a standardized extract from *Hibiscus sabdariffa* in patients with mild to moderate hypertension: a controlled and randomized clinical trial. *Phytomedicine.* 2004;11(5):375-82.
53. Herrera-Arellano A, Miranda-Sánchez J, Avila-Castro P, Herrera-Alvarez S, Jiménez-Ferrer JE, Zamilpa A, et al. Clinical effects produced by a standardized herbal medicinal product of *Hibiscus sabdariffa* on patients with hypertension. A randomized, double-blind, lisinopril-controlled clinical trial. *Planta Med.* 2007;73(1):6-12.
54. Haji Faraji M, Haji Tarkhani A. The effect of sour tea (*Hibiscus sabdariffa*) on essential hypertension. *J Ethnopharmacol.* 1999;65(3):231-6.
55. Mozaffari-Khosravi H, Jalali-Khanabadi BA, Afkhami-Ardekani M, Fatehi F. Effects of sour tea (*Hibiscus sabdariffa*) on lipid profile and lipoproteins in patients with type II diabetes. *J Altern Complement Med.* 2009;15(8):899-903.
56. Mozaffari-Khosravi H, Jalali-Khanabadi BA, Afkhami-Ardekani M, Fatehi F, Noori-Shadkam M. The effects of sour tea (*Hibiscus sabdariffa*) on hypertension in patients with type II diabetes. *J Hum Hypertens.* 2009;23(1):48-54.
57. Mohagheghi A, Maghsoud S, Khashayar P, Ghazi-Khansari M. The effect of *Hibiscus sabdariffa* on lipid profile, creatinine, and serum electrolytes: a randomized clinical trial. *ISRN Gastroenterol.* 2011;2011:976019.
58. Gurrola-Díaz CM, García-López PM, Sánchez-Enríquez S, Troyo-Sanromán R, Andrade-González I, Gómez-Leyva JF. Effects of *Hibiscus sabdariffa* extract powder and preventive treatment (diet) on the lipid profiles of patients with metabolic syndrome (MeSy). *Phytomedicine.* 2010;17(7):500-5.
59. Hopkins AL, Lamm MG, Funk JL, Ritenbaugh C. *Hibiscus sabdariffa* L. in the treatment of hypertension and hyperlipidemia: a comprehensive review of animal and human studies. *Fitoterapia.* 2013;85:84-94.
60. Cendrowski A, Ścibisz I, Mitek M, Kieliszek M, Kolniak-Ostek J. Profile of the Phenolic Compounds of *Rosa rugosa* Petals. *Journal of Food Quality.* 2017;2017:7941347.
61. Wan H, Yu C, Han Y, Guo X, Luo L, Pan H, et al. Determination of Flavonoids and Carotenoids and Their Contributions to Various Colors of Rose Cultivars (*Rosa* spp.). *Frontiers in Plant Science.* 2019;10.
62. Sarangowa O, Kanazawa T, Nishizawa M, Myoda T, Bai C, Yamagishi T. Flavonol glycosides in the petal of *Rosa* species as chemotaxonomic markers. *Phytochemistry.* 2014;107:61-8.
63. Boskabady MH, Shafei MN, Saberi Z, Amini S. Pharmacological effects of *rosa damascena*. *Iran J Basic Med Sci.* 2011;14(4):295-307.

64. Choi EM, Hwang JK. Investigations of anti-inflammatory and antinociceptive activities of *Piper cubeba*, *Physalis angulata* and *Rosa hybrida*. *J Ethnopharmacol.* 2003;89(1):171-5.
65. Mahmood N, Piacente S, Pizza C, Burke A, Khan AI, Hay AJ. The anti-HIV activity and mechanisms of action of pure compounds isolated from *Rosa damascena*. *Biochem Biophys Res Commun.* 1996;229(1):73-9.
66. Tripathi SC, Dixit SN. Fungitoxic properties of *Rosa chinensis* Jacq. *Experientia.* 1977;33(2):207-9.
67. Nyeem AB, Alam M, Awal, Mostafa M, Uddin SJ, Islam N, et al., editors. CNS DEPRESSANT EFFECT OF THE CRUDE ETHANOLIC EXTRACT OF THE FLOWERING TOPS OF *ROSA DAMASCENA* 2006.
68. Gholamhoseinian A, Fallah H, Sharifi far F. Inhibitory effect of methanol extract of *Rosa damascena* Mill. flowers on alpha-glucosidase activity and postprandial hyperglycemia in normal and diabetic rats. *Phytomedicine.* 2009;16(10):935-41.
69. Sadraei H, Asghari G, Emami S. Effect of *Rosa damascena* Mill. flower extract on rat ileum. *Res Pharm Sci.* 2013;8(4):277-84.
70. Hajhashemi V, Ghannadi A, Hajiloo M. Analgesic and Anti-inflammatory Effects of *Rosa damascena* Hydroalcoholic Extract and its Essential Oil in Animal Models. *Iran J Pharm Res.* 2010;9(2):163-8.
71. Deng J, Chen S, Yin X, Wang K, Liu Y, Li S, et al. Systematic qualitative and quantitative assessment of anthocyanins, flavones and flavonols in the petals of 108 lotus (*Nelumbo nucifera*) cultivars. *Food Chem.* 2013;139(1-4):307-12.
72. Limwachiranon J, Huang H, Shi Z, Li L, Luo Z. Lotus Flavonoids and Phenolic Acids: Health Promotion and Safe Consumption Dosages. *Compr Rev Food Sci Food Saf.* 2018;17(2):458-71.
73. Chen S, Fang L, Xi H, Guan L, Fang J, Liu Y, et al. Simultaneous qualitative assessment and quantitative analysis of flavonoids in various tissues of lotus (*Nelumbo nucifera*) using high performance liquid chromatography coupled with triple quad mass spectrometry. *Anal Chim Acta.* 2012;724:127-35.
74. Venkatesh B, Dorai A, editors. Antibacterial and Antioxidant potential of White and Pink *Nelumbo Nucifera* Gaertn Flowers 2011.
75. Paudel KR, Panth N. Phytochemical Profile and Biological Activity of *Nelumbo nucifera*. *Evid Based Complement Alternat Med.* 2015;2015:789124.
76. Tungmunnithum D, Pinthong D, Hano C. Flavonoids from *Nelumbo nucifera* Gaertn., a Medicinal Plant: Uses in Traditional Medicine, Phytochemistry and Pharmacological Activities. *Medicines (Basel).* 2018;5(4).
77. Huralikuppi JC, Christopher AB, Stephen PM. Anti-diabetic effect of *Nelumbo nucifera* (Gaertn): Part I preliminary studies in rabbits. *Phytotherapy Research.* 1991;5(2):54-8.
78. Park CH, Chae SC, Park SY, Kim JK, Kim YJ, Chung SO, et al. Anthocyanin and Carotenoid Contents in Different Cultivars of *Chrysanthemum* (*Dendranthema grandiflorum* Ramat.) Flower. *Molecules.* 2015;20(6):11090-102.
79. Shahrajabian MH. A REVIEW OF CHRYSANTHEMUM, THE EASTERN QUEEN IN TRADITIONAL CHINESE MEDICINE WITH HEALING POWER IN MODERN PHARMACEUTICAL SCIENCES. *Applied Ecology and Environmental Research.* 2019.
80. Lal M, Chandraker SK, Shukla R. 4 - Antimicrobial properties of selected plants used in traditional Chinese medicine. In: Prakash B, editor. *Functional and Preservative Properties of Phytochemicals: Academic Press; 2020.* p. 119-43.
81. Sassi AB, Harzallah-Skhiri F, Bourgougnon N, Aouni M. Antimicrobial activities of four Tunisian *Chrysanthemum* species. *Indian J Med Res.* 2008;127(2):183-92.
82. Pavelková P, Krmela A, Schulzová V. Determination of carotenoids in flowers and food supplements by HPLC-DAD. *Acta Chimica Slovaca.* 2020;13(1):6-12.

83. Bhawe A, Schulzová V, Mrnka L, Hajšlová J. Influence of Harvest Date and Postharvest Treatment on Carotenoid and Flavonoid Composition in French Marigold Flowers. *J Agric Food Chem*. 2020;68(30):7880-9.
84. Bakó E, Deli J, Tóth G. HPLC study on the carotenoid composition of *Calendula* products. *J Biochem Biophys Methods*. 2002;53(1-3):241-50.
85. Silva D, Ferreira MS, Sousa-Lobo JM, Cruz MT, Almeida IF. Anti-Inflammatory Activity of *Calendula officinalis* L. Flower Extract. *Cosmetics*. 2021;8(2):31.
86. Shahane K, Kshirsagar M, Tambe S, Jain D, Rout S, Ferreira MKM, et al. An Updated Review on the Multifaceted Therapeutic Potential of *Calendula officinalis* L. *Pharmaceuticals*. 2023;16(4):611.
87. Wang N, Zhang C, Bian S, Chang P, Xuan L, Fan L, et al. Flavonoid Components of Different Color *Magnolia* Flowers and Their Relationship to Cultivar Selections. *HortScience horts*. 2019;54(3):404-8.
88. Huang HC, Hsieh WY, Niu YL, Chang TM. Inhibition of melanogenesis and antioxidant properties of *Magnolia grandiflora* L. flower extract. *BMC Complement Altern Med*. 2012;12:72.
89. Jeyaraj EJ, Lim YY, Choo WS. Extraction methods of butterfly pea (*Clitoria ternatea*) flower and biological activities of its phytochemicals. *J Food Sci Technol*. 2021;58(6):2054-67.
90. Nair V, Bang WY, Schreckinger E, Andarwulan N, Cisneros-Zevallos L. Protective Role of Ternatin Anthocyanins and Quercetin Glycosides from Butterfly Pea (*Clitoria ternatea* Leguminosae) Blue Flower Petals against Lipopolysaccharide (LPS)-Induced Inflammation in Macrophage Cells. *J Agric Food Chem*. 2015;63(28):6355-65.
91. G.D N, Mohd Salleh R, Ong MT. Chemical composition and anti-proliferative properties of flowers of *Clitoria Ternatea*. *International Food Research Journal*. 2013;20:1229-34.
92. Chusak C, Thilavech T, Henry CJ, Adisakwattana S. Acute effect of *Clitoria ternatea* flower beverage on glycemic response and antioxidant capacity in healthy subjects: a randomized crossover trial. *BMC Complement Altern Med*. 2018;18(1):6.
93. Leong CR, Azizi MAK, Taher A, Wahidin S, Lee KC, Tan W-N, et al. Anthocyanins from *Clitoria ternatea* Attenuate Food-Borne *Penicillium expansum* and its Potential Application as Food Biopreservative. *Natural product sciences*. 2017;23:125-31.
94. Goh S, Kwong PJ, Ng CL, Ng W-J, Ee KY. Antioxidant-rich *Clitoria ternatea* L. flower and its benefits in improving murine reproductive performance. *Food Science and Technology*. 2021.
95. Multisona RR, Shirodkar S, Arnold M, Gramza-Michalowska A. *Clitoria ternatea* Flower and Its Bioactive Compounds: Potential Use as Microencapsulated Ingredient for Functional Foods. *Applied Sciences*. 2023;13(4):2134.
96. Sebai H, Jabri MA, Souli A, Rtibi K, Selmi S, Tebourbi O, et al. Antidiarrheal and antioxidant activities of chamomile (*Matricaria recutita* L.) decoction extract in rats. *J Ethnopharmacol*. 2014;152(2):327-32.
97. Miraj S, Alesaeidi S. A systematic review study of therapeutic effects of *Matricaria recutita* chamomile (chamomile). *Electron Physician*. 2016;8(9):3024-31.
98. Motealleh B, Zahedi P, Rezaeian I, Moghimi M, Abdolghaffari AH, Zarandi MA. Morphology, drug release, antibacterial, cell proliferation, and histology studies of chamomile-loaded wound dressing mats based on electrospun nanofibrous poly( $\epsilon$ -caprolactone)/polystyrene blends. *J Biomed Mater Res B Appl Biomater*. 2014;102(5):977-87.
99. Jabri MA, Sani M, Rtibi K, Marzouki L, El-Benna J, Sakly M, et al. Chamomile decoction extract inhibits human neutrophils ROS production and attenuates alcohol-induced haematological parameters changes and erythrocytes oxidative stress in rat. *Lipids Health Dis*. 2016;15:65.

100. Ranpariya VL, Parmar SK, Sheth NR, Chandrashekhar VM. Neuroprotective activity of *Matricaria recutita* against fluoride-induced stress in rats. *Pharm Biol.* 2011;49(7):696-701.
101. Chandrashekhar VM, Halagali KS, Nidavani RB, Shalavadi MH, Biradar BS, Biswas D, et al. Anti-allergic activity of German chamomile (*Matricaria recutita* L.) in mast cell mediated allergy model. *J Ethnopharmacol.* 2011;137(1):336-40.
102. Bulgari M, Sangiovanni E, Colombo E, Maschi O, Caruso D, Bosisio E, et al. Inhibition of neutrophil elastase and metalloprotease-9 of human adenocarcinoma gastric cells by chamomile (*Matricaria recutita* L.) infusion. *Phytother Res.* 2012;26(12):1817-22.
103. Chang SM, Chen CH. Effects of an intervention with drinking chamomile tea on sleep quality and depression in sleep disturbed postnatal women: a randomized controlled trial. *J Adv Nurs.* 2016;72(2):306-15.
104. Rafraf M, Zemestani M, Asghari-Jafarabadi M. Effectiveness of chamomile tea on glycemic control and serum lipid profile in patients with type 2 diabetes. *J Endocrinol Invest.* 2015;38(2):163-70.
105. Sharifi F, Simbar M, Mojab F, Majd HA. Comparison of the effects of *Matricaria chamomila* (Chamomile) extract and mefenamic acid on the intensity of premenstrual syndrome. *Complement Ther Clin Pract.* 2014;20(1):81-8.
106. Sah A, Naseef PP, Kuruniyan MS, Jain GK, Zakir F, Aggarwal G. A Comprehensive Study of Therapeutic Applications of Chamomile. *Pharmaceuticals (Basel).* 2022;15(10).
107. Purohit SR, Rana SS, Idrishi R, Sharma V, Ghosh P. A review on nutritional, bioactive, toxicological properties and preservation of edible flowers. *Future Foods.* 2021;4:100078.
108. Matyjaszczyk E, Śmiechowska M. Edible flowers. Benefits and risks pertaining to their consumption. *Trends in Food Science & Technology.* 2019;91:670-4.
109. Wilczyńska A, Kukułowicz A, Lewandowska A. Preliminary assessment of microbial quality of edible flowers. *LWT.* 2021;150:111926.
110. Lucarini M, Copetta A, Durazzo A, Gabrielli P, Lombardi-Boccia G, Lupotto E, et al. A Snapshot on Food Allergies: A Case Study on Edible Flowers. *Sustainability.* 2020;12(20):8709.
111. Akindahunsi AA, Olaleye MT. Toxicological investigation of aqueous-methanolic extract of the calyces of *Hibiscus sabdariffa* L. *J Ethnopharmacol.* 2003;89(1):161-4.
112. Sireeratawong S, Itharat A, Khonsung P, Lertprasertsuke N, Jaijoy K. Toxicity studies of the water extract from the calyces of *Hibiscus sabdariffa* L. in rats. *Afr J Tradit Complement Altern Med.* 2013;10(4):122-7.
113. Akbari M, Kazerani HR, Kamrani A, Mohri M. A preliminary study on some potential toxic effects of *Rosa damascena* Mill. *Iranian Journal of Veterinary Research.* 2013;14(3):232-6.
114. Li L, Gu L, Chen Z, Wang R, Ye J, Jiang H. Toxicity study of ethanolic extract of *Chrysanthemum morifolium* in rats. *J Food Sci.* 2010;75(6):T105-9.
115. Grzeszczuk M ME, Stefaniak A, Wysocka G. Mineral Composition of Some Edible Flowers. *J Elem.* 2017;1/2018.
116. Drava G, Iobbi V, Govaerts R, Minganti V, Copetta A, Ruffoni B, et al. Trace Elements in Edible Flowers from Italy: Further Insights into Health Benefits and Risks to Consumers. *Molecules.* 2020;25(12):2891.
117. Benvenuti S, Mazzoncini M. The Biodiversity of Edible Flowers: Discovering New Tastes and New Health Benefits. *Front Plant Sci.* 2020;11:569499.