

Original Research Article Novel, Green Synthesis and Antimicrobial activity of Amine-Boranes

ABSTRACT

Amine-Boranes (AB) are an elite class of reagent and belongs to the Borane-Lewis family, has found extensive use in chemical synthesis. Interesting aspect of AB reagents are to display chemo selectivity among the functional groups. The present investigation highlights on the innovative, green synthesis and characterization of the three of unique AB complexes, including triethylamine-borane (aliphatic), aniline-borane (aromatic), and (aromatic substituted) N,N-dimethylaniline-borane (DMAB) under an open air condition. The synthesized AB reagents were characterized by employing the ^{11}B NMR technique, the chemical shift values were consistent with previous investigations. The microbiological efficacy of three AB reagents has been stated for the first instance by using disc diffusion method against four bacterial strains: *Escherichia coli*, *Salmonella typhimurium*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*.

Keywords: Amine-Borane, ^{11}B -NMR, *P. aeruginosa*, *S. typhimurium*, *S. aureus*, *E. coli*

1. INTRODUCTION

Although AB reagents have been existing and well-studied for decades and might be used in place of frequently used borane reagents, their stability and inertness make them less effective for synthetic processes. prefers to react with nucleophilic sites and reagents [1-6]. Without a doubt, they might serve as substitutes for frequently used borane reagents, and in comparison, to other electrophilic boron reagents, the majority of the complexes demonstrate stability, inertness, and slow reactivity has been documented by various researchers [7-9] and the multitude applications of AB complexes are displayed (Fig. 1).




Fig. 1. Multitude applications of Amine.Borane complexes (Kulkarni & Ramachandran, 2017)

When it comes to organic synthesis, AB reagents have the following benefits over other electrophilic boron reagents: 1. They are easy to work with and convenient; 2. They exhibit chemo- and regio-selectivity; 3. They are stable and interdependent[10-12]. Dimethylamine-Borane and Morpholine-Borane complexes are frequently employed as reducing agents in the electronics sector for the fabrication of circuit boards and electroplating applications due to their solubility and mild reducing capabilities[13]. In accordance with recent study reports, transition metal catalysts are currently applied in an attempt to enhance their reactivity in hydrogenation reactions[14-17].

In the study by PV Ramachandran *et al* (2020), the AB reagents have a dual function in direct amidation, whereby they may activate both aromatic and aliphatic carboxylic acids and then deliver amines to produce corresponding functionalized amides with a yield of 99%. The utilization of these AB complexes was reported in hydrogen storage fuel cells, due to its stability and high gravimetric content of hydrogen[18-19].

AB reagents reducing ability can be affected by the steric effects and electronic characteristics of the groups bonded to the nitrogen atom. The incorporation of acetic acid, mineral acid, Lewis acid, and high reaction temperature is necessary to activate the stable AB complexes such as Trimethylamine-borane, Pyridine-borane[20-21].

The nature of bond present in AB's is coordinate bond which is semi-dative polar bond as shown in the Fig. 2. The reactivity of AB complexes depends on the nature and bond strength of N-B bond. If AB is less stable then it is less reactive, inert and easy to prepare. The reverse is true if the AB adduct is unstable and difficult to prepare.




Fig. 2. Dative bond formation in an Amine-Borane complex (Kulkarni & Ramachandran, 2017)

In the previous research studies, we select a moderately reactive DMAB reagent, microwave irradiation and ultrasound sonication techniques were chosen for activating the AB complex *in situ*. The ease of recovery of the amine after the reaction and the possibility of recycling it make the AB as environmentally friendly reagents of choice. We first reported the preparation, properties and synthetic utility of DMAB (N,N-dimethyl aniline.borane) under microwave and ultrasound conditions[22-26].

2. EXPERIMENTAL WORK

2.1 Materials and Methods

The chemicals used for the experiment are purchased from Fluka and Aldrich and the technical bulletin procedure of Sigma-Aldrich is followed to handle the air-sensitive reagents[27]. Liquid substrates and solvents were distilled prior to use. NMR spectra for product were recorded on Bruker AVANCE spectrometer, 400 MHz for ^{11}B NMR. Solvents used to dissolve the products are CDCl_3 , CD_3OD and DMSO and the chemical shift values were referenced to TMS and coupling constants were calculated in hertz (Hz). IR and MS spectra was recorded on FTIR, EI-Shimadzu-GC-MS Shimadzu spectrometer.

2.2 Test microorganisms

The strains of bacteria that were employed were *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli*, and *Salmonella typhimurium*. Gentamicin was used as a positive control in the antimicrobial analysis utilising the antibiotics disc. The pathogenic strains were collected from the Department of Microbiology, Devi Ahilya Vishwavidyalaya (DAVV), Indore, MP. The bacteria were kept on nutrient agar slants at 4°C and cultivated in nutrient agar medium.

2.3 Processes involved in synthesising AB reagents


The novel method is used for preparing the DMAB reagent is as follows, taking 3.80 ml of N,N-dimethyl aniline (0.03 moles) and adding it to a 100 ml two-necked RB flask in addition to 15 ml of chloroform, 10 ml of water, and 5 mg of phase transfer catalyst while stirring. After 15 minutes of cooling the reaction mixture in an ice bath, 2.3g of sodium borohydride (0.06 moles) was added portion-wise, and the temperature was kept below 10°C . After that, the reaction ingredients continue

to remain to stir for furthermore two hours. The complexation propensity of an amine can be monitored using ^{11}B -NMR spectroscopy, and the reaction mixture is filtered under vacuum once complexation is accomplished. The filtrate is blended with 10 millilitres of 2N HCl, and the DMAB complex can be extracted with chloroform. After that, the collected extractions using chloroform get dried over anhydrous sodium sulphate, and the solvent is removed under vacuum to yield the DMAB reagent.

The DMAB stock solution is kept in an amber colour bottle and further confirmed by ^{11}B NMR spectra, which show chemical shift values of -4 ppm (decoupled) and -4 to -9 ppm (coupled).

The other two AB reagents (triethylamine and aniline) are prepared using the same methodology as described above and the ^{11}B NMR spectra of the synthesized AB reagents are presented as (Fig. 4-6).

^{11}B -NMR spectra and gasometer analysis were used to periodically determine the stability of the three AB reagents.



Scheme 1. Preparation of DMAB

2.4 Characterization of AB complex

The characterization of the synthesized AB reagent is necessary, and this can be done with the support of the ^{11}B -NMR spectral technique, and it is necessary for studying and comprehending the AB reagents. The ^{11}B - chemical shift values can be used as criteria to measure the complexation strength of the N-B dative bond. Table 1. shows that there is a minimum fluctuation in ^{11}B - chemical shift values, despite the fact that aliphatic and aromatic amines differ significantly in their basicity.

Table 1. Typical ^{11}B -NMR spectral values for various AB reagents

No	Amine.Borane (AB) complex	^{11}B -NMR peak value (ppm)	Nature of the signal
1	Trimethylamine.Borane	-13 coupled -10 to -13 decoupled	Quartet
2	Triethylamine.Borane	-11 coupled	Quartet

3	Pyridine.Borane	-8 to -11 decoupled -12 coupled	Quartet
4	N-ethyl-N-methyl aniline.Borane	-5 coupled -5 to -9 decoupled	Quartet
5	N,N-diethylaniline.Borane	-4 coupled -4 to -9 decoupled	Quartet

Further, Wang *et al.* (2020) reported that the borane adducts of aniline and aminopyridines demonstrated unique ^{11}B -NMR chemical shift values, due to paramagnetic anisotropic effects and along with hydrogen bonding [12]. The groups that are bound to the N-gen atom of amine group in the AB reagent dictate the type and position of the ^{11}B -NMR signal in the spectra. Table I summarizes the type and position of the ^{11}B -NMR signal of AB reagents.

2.5 Stability studies of ab reagents

While the much of AB complexes are widely recognised for their stability and inertness, reactive ABs are often less stable and more laborious to synthesize. ^{11}B -NMR spectrum investigations are used to examine the stability of the synthesized AB reagents.

2.5.1 Stability of DMAB reagent

A sample of aliquot (about 1ml) is withdrawn from the stock solution, which is kept in an environment of nitrogen, and it is examined to see if any additional signals above -4 to -7 ppm are present. The DMAB reagent is stable for up to eight days, according to the results of ^{11}B -NMR spectral studies. After that, a new signal at +20 ppm is detected, which is associated with boric acid. DMAB reagent has a 7-day shelf life, after which it begins to gradually degrade into boric acid. The stock solution becomes progressively darker in blue colour over the course of seven days and the figure demonstrates a variation in borane's molar concentration over the course of time.




Fig. 3. Stability studies of DMAB Borane **Fig. 4. ^{11}B -NMR spectra of Triethylamine.Borane**




Fig. 5. ^{11}B -NMR spectra of DMAB Borane **Fig. 6.** ^{11}B -NMR spectra of Aniline.Borane

2.6 ANTIMICROBIAL ACTIVITY:

2.6.1 Procedure of antibacterial activity of AB complexes^{28,29}

Microbes continue to be the most prevalent cause of death worldwide due to minimal or inadequate treatment. There is a pressing need at the moment for a novel class of antimicrobials with an alternative method of action. The antibacterial capability of synthesized AB reagents was investigated using the following species of bacteria: *Salmonella typhimurium*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*. AB samples were tested at a concentration level of 100 mg per millilitre, and the plates' medium was permitted to settle down for 10 to 20 minutes at room temperature for the compound to diffuse into the agar. In order to encourage the bacterial growth, the plates were eventually incubated for 24 hours at 35°C. The disc-diffusion method was used to test the antibacterial activity. Following incubation, a distinct zone of inhibition demonstrated the antibacterial activity, which was then measured using a ruler. The outcomes are shown in Table 2.

2.6.2 Assessment of Antibacterial effects using the agar disc diffusion method

Whatman No. 3 filter paper discs (diameter 6 mm) that had been sterilized were placed on the surface of a petri dish for inoculation. The three different AB reagents had been added to the Petri dishes after being independently dissolved in DMSO. In brief, each disc contained 100 μl of AB reagent as the test sample, 10 μg of gentamycin of the positive control, and 100 μl of DMSO as the negative control. A measuring tool can be used to measure the zone of inhibition, which represented the effectiveness of antibacterial agents. The zone of inhibition illustrated in Figure 7 and the Table II reveal the findings. The diameters of the inhibitory zones induced by the AB complexes were interpreted as

follows: resistant (12 mm or less), intermediate (13–15 mm), and susceptible (16 mm or above), following the standard.




Fig. 7. A- *S. typhimurium* B- *P. aeruginosa* C- *S. aureus* D- *E. coli*

Table 2. Antibacterial properties of AB reagents

No	AB reagent name	<i>E. coli</i>	<i>S. typhimurium</i>	<i>S. aureus</i>	<i>P. aeruginosa</i>
1	Triethylamine.Borane	12	16	14	15
2	Aniline.Borane	16	14	17	18
3	N,N-Dimethylaniline.Borane (DMAB complex)	21	18	17	22
4	Gentamicin*	26	24	25	27
5	DMSO solvent**	-	-	-	-

The measurements are expressed in mm; *(+ve control; **(-)ve control; No inhibition Zone; Solvent

3. RESULTS AND DISCUSSION

With the assistance of the ^{11}B -NMR technique, stability inquiries of the synthesized AB reagents demonstrate that Triethylamine.Borane is inert and stable while Aniline.Borane is reactive.

3.1 Evaluation of AB reagents' antimicrobial properties

The four known bacterial pathogens *S. typhimurium*, *P. aeruginosa*, *S. aureus*, and *E. coli* were employed to examine the three AB compounds antibacterial properties and percentage inhibition determined using the standard approach. The results are shown in Table II, and it was discovered that DMAB complex demonstrated a significant inhibition zone against *E. coli* and *P. aeruginosa*. The type of substituent present on AB reagents has a consequence on the degree of activity. The highest zone of inhibition was observed against *S. aureus* and *P. aeruginosa* with Aniline.Borane reagent. Triethylamine- Borane complex exhibited the largest zone of inhibition against *S. typhimurium* and followed by *P. aeruginosa*.

4. CONCLUSIONS

In the past few decades, lead bioactive compounds possessing nitrogen have become appealing synthetic targets owing to their structural diversity and biological significance. In consonance with current investigation outcomes, AB reagents will be the promising lead molecules for drug development in the future research. A great deal of work has gone into developing a new, easy-to-use method for synthesizing the AB complexes using sodium borohydride and chloroform.

The *in vitro* evaluation of three AB reagents have demonstrated intriguing effectiveness against bacterial pathogens and DMAB complex exhibited excellent bacterial inhibition. The subsequent research project will be emphasized on the generation of substituted, pharmacologically active Aniline.Borane, reagent. The results of characterization, molecular docking, simulation, and *in vivo* research investigations will be presented in the near future.

REFERENCES:

1. Reddy DO. A short chronological Review on the Synthesis of Amine-Boranes. Chemical Review and Letters. 2020; 4: 184-91. doi: <https://doi.org/10.22034/crl.2020.238.849.1070>
2. Brown H C. Organic synthesis via Boranes Vol.1. Aldrich chemical company, Milwaukee. 1997.
3. Meller A. Gmelin Handbook of Inorganic and Organometallic Chemistry, 4th Supplement, Vol. 3, Springer, Berlin, 1992. (ISBN: 3540936734)
4. Marjorie AY, Marty BJD. and Ciuculescu-Pradines CD. Chapter 6: Amines and Amine-boranes. Series: Nanoscience and Nanotechnology, 2021. doi: <https://doi.org/10.1039/9781839163623-00130>
5. Burnham BS. Amine-Boranes, Encyclopedia of Metalloproteins. Editors: Kretsinger RH, Uversky VN, Permyakov EA. Springer New York, 2013.
6. Long LH. In A Comprehensive Treatise on inorganic and theoretical chemistry, Supplement Vol.5, Part B1, Longman, London, 1981.
7. Okuda J, Mashima K, Shirase S, Mukherjee D. Chemo-selective Reduction of Tertiary Amides to Amines Catalyzed by Triphenylborane. Angewandte Chemie International Edition. 2016; 55: 13326-31. doi: <https://doi.org/10.1002/anie.201605236>
8. Bellham P, Hill MS, Köhn KG. Stoichiometric and catalytic reactivity of tert-butylamine-borane with calcium silylamides. Organometallics. 2014; 33(20): 5716-21. doi: <https://doi.org/10.1021/om5.00467b>
9. Ramachandran PV, Kulkarni AS, Zhao Y, Mei GJ. Amine-boranes bearing borane-incompatible functionalities: application to selective amine protection and surface functionalization. Chemical Communication. 2016; 52: 11885-88.
10. Ramachandran PV, Kulkarni AS. Open-Flask Synthesis of Amine-Boranes via Tandem Amine-Ammonium Salt Equilibration-Metathesis. Inorganic Chemistry. 2015; 54: 5618-20.
11. Bailey CL, Joh AY, Hurley Z Q, Anderson C L, Singaram B. Controlled Reduction of Tertiary Amides to the Corresponding Alcohols, Aldehydes, or Amines Using Dialkylboranes and Amino borohydride Reagents. Journal of Organic Chemistry. 2016; 81(9): 3619-26. doi: <https://doi.org/10.1021/acs.joc.6b00276>
12. Wang X, Yang Y, Song Y, Zhao S, Qu JH. Research Progress on N-heterocyclic Carbene Boranes. Scholars International Journal of Chemistry and Material Sciences. 2020; 3(10): 152-63. doi: <https://doi.org/10.36348/sijcms.2020.V03i1.0.002>
13. G.O. Mallory, J.B. Hajdu, Electroless plating, Fundamentals and Applications. Noyes Publications/William Andrew, Orlando, Florida, 1990. <https://trove.nla.gov.au/work/6453483>
14. Rossin A, Peruzzini M. Ammonia-Borane and Amine-Borane Dehydrogenation Mediated by Complex Metal Hydrides. Chemical Reviews. 2016; 116: 8848-72. doi: <https://doi.org/10.1021/acs.chemrev.6b00043>
15. Patni NA, Mantri SA, Kundu D. Ionic liquid promoted dehydrogenation of Amine.Boranes: A review. International Journal of Hydrogen Energy. 2021; 46(21): 11761-81, doi: <https://doi.org/10.1016/j.ijh.2021.01.032>

16. Couturier M, Andresen BM, Dubé P, Brenek SJ, Negri JT. The use of borane–amine adducts as versatile palladium catalysed hydrogen-transfer reagents in methanol. *Tetrahedron Letters*. 2001; 42 (1): 2763-66.
17. Lin R, Scherschel NF, Zeller M, Hamlin SG, Snyder M, Son S, Ramachandran PV, Piercey DG. Synthesis and Energetic Characterization of Borane-Amines on High-Nitrogen Heterocycles. *ACS Omega*. 2024; 9: 14241-48. doi: <https://doi.org/10.1021/acsomega.3c09934>
18. Ramachandran PV, Kulkarni AS. Water-promoted, safe and scalable preparation of Ammonia-Borane. *International Journal of Hydrogen Energy*. 2017; 42(2): 1451-55. doi: <https://doi.org/10.1016/j.ijhydene.2016.06.231>
19. Ramachandran PV, Hamann HJ, Choudhary S. Amine-Boranes as Dual-purpose reagents for direct Amidation of Carboxylic acids. *Organic Letters*. 2020; 22 (21): 8593-97. doi: <https://doi.org/10.1021/acs.orglett.0c03184>
20. Kelly HC, Mario B. Giusto MB, Marchelli FR. Amine-Borane Reductions in Aqueous Acid Media. *Journal of American Chemical Society*. 1964; 86(18): 3882-84. doi: <https://doi.org/10.1021/ja01072a063>
21. Hullio AA, Mastoi GM. Nicotine-Borane-based task specific ionic liquid: an alternative to Pyridine-Borane for efficient reductive Amination. *Student's Journal of Chemistry*. 2013; 1(2): 57-67.
22. Jayakumar SV, Ahamed MFA. Ultrasound Mediated Functional Group Reduction and Chemo-selective Studies by DMAB. *Chemical Science Transactions*. 2019; 8: 418-24. doi: <https://doi.org/10.7598/cst2019.1581>
23. Jayakumar SV, Srinivas KA, Hirianna SG, Pati HN. Microwave accelerated hydroboration of alkenes with N,N-Dimethylaniline-Borane. *Rasayan Journal of Chemistry*. 2008; 1: 326-30.
24. Jayakumar SV, Ahamed MFA. Ultrasound Mediated Hydroboration of Alkenes, Alkynes Employing: N,N-Dimethylaniline-Borane (DMAB). *Der Pharma Chemica*. 2018; 10(3): 90-93.
25. Jeyakumar SV, Ganesh SK. Microwave Mediated Reduction of Selected Functional Groups Employing: N,N-Dimethylaniline.borane. *Orbital: Electronic Journal of Chemistry*. 2014; 6: 56-60.
26. Jayakumar SV, Saxena A, Tomar DR. Microwave Mediated Modification of Selected Limonoids with DMAB Reagent and Antibacterial Studies. *Macromolecular Symposia*. 2024; 413: 2300005-15. doi: <https://doi.org/10.1002/masy.202300005>
27. Sigma-Aldrich, Handling Air-Sensitive Reagents, Technical Bulletin AL-134, Sigma-Aldrich.com
28. Mhaibes MR. Antimicrobial and Antioxidant activity of heterocyclic compounds derived from new Chalcones. *Journal of Medicinal and Chemical Sciences*. 2023; 6(4): 931-37. doi: https://doi.org/10.26655/JM_CHEMSCI.2023.4.25
29. Zayda GM, Abdel-Rahman HAA, El-Essawy EF. Synthesis and Antibacterial activities of different five-membered heterocyclic rings incorporated with Pyrido-thienopyrimidine. *ACS Omega*. 2020; 5: 6163-68. doi: <https://dx.doi.org/10.1021/acsomega.0c00188>