

Effect of Potassium Schoenite (Mahalaabh) on Growth and Yield of Castor, Cotton and Groundnut

Abstract

The need for precise nutrient application has become crucial in agriculture due to climate change and intensive farming practices. Deficiencies in potassium, magnesium and sulphur are critical for enhancing both crop yield and quality. Additionally, agrochemical residues in soil, water and farm products pose significant challenges. Mahalaabh ($K_2SO_4 \cdot MgSO_4 \cdot 6H_2O$), a combination of potassium, sulphur and magnesium, offers a balanced solution to address these issues. Field experiments were conducted during the *Kharif* season of 2023-24 at Agrocel Industries Pvt. Ltd., Kachchh, to evaluate the impact of Mahalaabh on castor, cotton and groundnut crops. The trials were executed in a Randomized Block Design, testing different doses of Mahalaabh powder through foliar spray, fertigation and granular soil application. The results demonstrated that Mahalaabh powder is effective in enhancing productivity across the tested crops. In castor, higher doses of 10 kg/acre through drip irrigation and 20g/L foliar application increased productivity by 22.3% and 14.3% over the recommended fertilizer dose (RFD), respectively. However, in cotton, lower doses of 6kg/acre through drip irrigation and 10g/L through foliar application were more effective, resulting in 3.8% and 5.3% yield enhancements compared to RFD. In groundnut, foliar spray at 15 g/L applied thrice at 30-35, 50-55, and 70-75 days after sowing (DAS) was effective, resulting in a 22.9% higher pod yield compared to RFD. These findings highlight the importance of balanced nutrient management practices tailored to specific crop requirements for achieving optimal agricultural productivity.

Keywords: Mahalaabh, Potassium, Cotton, Castor, Groundnut

Introduction

In the era of climate change-induced challenges and intensive agricultural practices, the precise application of nutrients as per soil and plant needs has become paramount. The deficiency of essential nutrients such as potassium, magnesium and sulphur are increasingly evident, posing a significant hurdle to quantitative and qualitative improvements in agricultural production. Additionally, the accumulation of agrochemical residues in soil, water and farm products presents another critical issue hindering sustainable agricultural production and natural resource conservation.

Mahalaabh ($K_2SO_4 \cdot MgSO_4 \cdot 6H_2O$), a unique 3-in-1 combination of potassium, sulphur and magnesium, offers a promising solution as a neutral and chemically balanced natural product. Potassium Schoenite, a constituent of Mahalaabh, has demonstrated effectiveness in increasing agricultural productivity by providing potassium and magnesium. Previous studies have demonstrated the benefits of applying foliar sprays of potassium schoenite along with recommended doses of NPK in improving plant growth, yield and quality in crops like potato (Ghose *et al.*, 2017).

Castor, a key non-edible oilseed crop, is predominantly grown in semiarid and arid regions, with India being the leading producer with 19.73 lakh tonnes annual production (Anonymous, 2023). Cotton, despite facing competition from synthetic fibres, remains the

world's leading natural textile fibre, with India ranking first in cultivated area (123.16 lakh ha) and second in production with 312.07 lakh bales (Anonymous, 2023). Groundnut, often referred to as the "King of Oilseeds", is a well-known oilseed crop belonging to the *Leguminosae* family, with India being the second-largest producer globally with 104.54 lakh tonnes annual production (Anonymous, 2023).

In groundnut, adequate potassium supply is essential for proper pod development, kernel filling and maintaining high oil quality (Sanadi *et al.*, 2018). Similarly, in cotton, potassium fertilization significantly increased boll weight, boll number, lint yield and fibre quality parameters including fibre length and strength (Pettigrew *et al.*, 2005). In castor, potassium fertilization significantly increased seed yield, oil content and oil yield (Sagare and Singh, 1988, Nayak *et al.*, 2004).

Magnesium fertilization in groundnut, improves pod number, pod yield and kernel yield, possibly due to its role in enhancing photosynthesis and nutrient translocation (Sireesha and Dawson, 2022). In cotton, magnesium nutrition improved boll retention, boll weight, lint yield and fibre quality traits like micronaire and maturity (Longwell *et al.*, 1963). In castor, magnesium application improved seed yield, oil content and oil yield, possibly due to its role in enhancing photosynthesis and nutrient translocation (Munir *et al.*, 2007), while sufficient magnesium supply is crucial for maintaining high oil quality and stability in seeds (Reddy *et al.*, 2004), Reddy *et al.*

Sulphur is another essential nutrient that plays a significant role in enhancing productivity and quality in these crops. In groundnut, sulphur application significantly increased pod yield, kernel yield and oil content, indicating its importance for overall productivity and quality (Rao *et al.*, 2013). In cotton, sulphur application significantly increased boll weight, lint yield and fibre quality parameters like fibre length and strength (Rasheed *et al.*, 2006). In castor, sulphur fertilization significantly increased seed yield, oil content and oil yield (Munir *et al.*, 2007), while adequate sulphur nutrition is essential for maintaining high protein quality and oil stability in seeds (Reddy *et al.*, 2004).

Considering the above situation-specific facts, the experiments were planned to optimize fertilization strategies and enhance crop productivity in the arid region of Kachchh, field trials were conducted at the AGROCEL R & D Farm at Koday, Mandvi during the *Kharif* season of 2023-24 to evaluate the efficacy of this balanced nutrient formulation on the yield and yield components of three economically important crops: castor, cotton and groundnut.

Material and Methods

Field experiments were conducted in a Randomized Block Design (RBD) with three replications at AGROCEL Industries Pvt. Ltd. R&D Farm, Koday, Mandvi, Kachchh during the *Kharif* season of 2023-24 to assess the effect of Potassium Schoenite (Mahalaabh) on the growth and yield of castor (*Ricinus communis* L.), cotton (*Gossypium hirsutum* L.) and groundnut (*Arachis hypogea* L.). The treatments comprised different doses of Mahalaabh powder used both as foliar spray and drenching; and Mahalaabh granules as soil application.

The experiments site represented sandy loam soil with a light brown colour, well-drained and fairly retentive of moisture. The soil was low in available nitrogen, optimum in available

phosphorus and medium in available potassium. Fertilizer application was carried out according to the respective treatments for each crop. All necessary agronomic practices were followed as per the recommended practices for the respective crops. The biometric observations were recorded from five randomly selected plants within each net plot. The parameters observed included plant height (cm), spike length (cm), number of spikes/plant, number of capsules/spike, number of bolls/plant, number of monopodial branches/plant, number of sympodial branches/plant, number of pods/plant, number of seeds/pod, weight of seed cotton/boll (g) and weight of seed cotton/plant (g). That apart, plant population, days to flowering, seed index (g), seed yield (kg/ha), seed cotton yield (kg/ha), and pod yield (kg/ha) were recorded on a net plot basis. The data so collected were statistically analysed for ascertaining precise impact of the treatments on the growth and yield parameters of the respective crops.

Result and Discussion

Effect of Potassium Schoenite on Castor

The data on various treatments in castor revealed non-significant differences in the growth and yield parameters. However, application of Mahalaabh whether as granular basal dose at 50 kg/acre in addition to RFD (Recommended Fertilizer Dose) or Mahalaabh Powder both through drip and foliar was numerically effective in increasing seed yield up to 20.7%, 22.3% and 14.3%, respectively, over RFD (875 kg/acre). Further, Mahalaabh Powder applied through drip irrigation at rates varying from 6 kg/acre to 10 kg/acre showed 17.6% to 22.3% higher seed yield compared to the RFD (Table 1). Overall, T3 (RFD + MLP 10 kg/acre at 40-50 DAS & 100-110 DAS through Drip) gave the highest seed yield (1070 kg/acre) with more numbers of spikes/plant (11.87), capsules/spike (115.87) and seed index (31.44 g), indicating the efficacy of MLP through drip irrigation at higher doses. Similarly, foliar applications were also effective at higher doses with 20g/L giving 14.3% higher seed yield than RFD with conspicuous improvements in spike length, spikes/plant and capsules/spike. These results are in consonance with the findings of Vaghasia *et al.* (2019) who have also reported a conspicuous impact of potassic fertilizers in castor with yield enhancement. Adequate potassium supply increases seed yield, oil content and oil yield (Sagare and Singh, 1988). *Muniret al.*, (2007) also reported that seed yield, oil content and oil yield improve when sufficient magnesium is applied. Adequate application of sulphur enhances seed formation, seed weight and yield (Munir *et al.*, 2007).

Effect of Potassium Schoenite on Cotton

The data on various treatments indicated that there are no significant differences in the growth and yield parameters of cotton. However, treatments included combinations of recommended fertilizer doses (RFD) supplemented with different doses of potassium schoenite applied either through drip irrigation or as foliar sprays gave numerically higher yields than RFD alone. Mahalaabh Powder was more efficient at lower doses. T1 (RFD + Mahalaabh Powder 6 kg/acre) and T4 (RFD + MLP 10g/L) applied at 30-35 DAS and 50-55 DAS emerged as the best doses in the drip and foliar application categories, respectively (Table 2). These treatments resulted in 3.8% and 5.3% higher seed cotton yield compared to the recommended fertilizer dose (1090 kg/acre). In both cases, the seed cotton yield enhancements were attributed to an increase in the number of monopodial branches per plant and the number of bolls per plant. These results underscore the importance of

balanced nutrient management and application methods in maximizing cotton yield. Similar results have been reported by Read *et al.* (2006) and Jyothi and Hebsur (2018). Pettigrew (2008) reported that potassium (K) plays a crucial role in enhancing cotton yield and fibre quality; also increases boll weight, boll number and lint percentage. Magnesium (Mg) improves boll retention, boll weight and lint yield (Safaya, 1976). Similarly, adequate sulphur supply enhances boll development, boll weight and lint yield (Rasheed *et al.*, 2006).

Effect of Potassium Schoenite on Groundnut

In groundnut, foliar sprays of Mahalaabh were more effective than its application through drip irrigation. Specifically, the treatment involving the recommended fertilizer dose (RFD) along with Mahalaabh powder at the rate of 15 g/L sprayed thrice at 30-35, 50-55 and 70-75 days after sowing (DAS), resulted in a 22.9% higher pod yield compared to RFD alone (722 kg/acre). Furthermore, among the seed yield components, the seed index that reflects seed weight showed a substantial improvement (43.05 g). Thus, foliar sprays of Mahalaabh powder are quite effective in enhancing groundnut pod yield. These findings are in consonance with the findings of Chandra *et al.* (2006), Hemeid (2015) and Borah *et al.* (2017). Veeramani *et al.*, (2012) reported that potassium plays a vital role in pod development, kernel filling and yield in groundnut. Adequate potassium supply increases pod yield, kernel yield and shelling percentage (Rajet *et al.*, 2004). Magnesium is crucial for improving pod yield, kernel yield and oil content (Sharavathi *et al.*, 2019). Sulphur plays a conspicuous role in enhancing pod formation, kernel development and yield (Jena *et al.*, 2006).

Conclusion

It can be concluded from the present study that Mahalaabh powder is quite effective in enhancing productivity across crops. In castor, higher doses of 10 kg/acre through drip and 20g/L foliar application were effective in increasing productivity by 22.3% and 14.3% over RFD, respectively. However, in cotton, lower doses of 6kg/acre through drip and 10g/L through foliar were more effective and resulted in 3.8% and 5.3% yield enhancements compared to RFD. In groundnut, only foliar spray at 15 g/L sprayed thrice (30-35, 50-55 and 70-75 DAS) was effective, resulting in a 22.9% higher pod yield compared to RFD. These findings underscore the importance of crop specific customised nutrient management practices tailored to specific crop requirements for achieving optimal agricultural productivity.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

References

Anonymous (2023). Directorate of Economics and Statistics, Department of Agriculture and Farmer Welfare, Ministry of Agriculture and Farmer Welfare, Government of India.

- Borah, B.; Patil, D. S. and Pawar, R. B. (2017). Enhancing kharif groundnut (*Arachishypogaea* L.) yield and quality in Entisol through potassic fertilizer management. *International Journal of Current Microbiology and Applied Sciences*. 6(11): 4068-4074.
- Chandra, P.; Samui, R.C. and Bordolui, S. K. (2006). Growth, yield attributes and yield of different cultivars of groundnut (*Arachishypogaea* L.) as affected by potassium application. *Journal of Crop and Weed*. 2(1): 37-39.
- Ghosh, D.; Sarkar, S.; Brahmachari, K.; Garai, S.; Pal, M. and Sharma, A. (2017). Potassium schoenite: an emerging source of potassium for improving growth, yield and quality of potato. *Journal of Experimental Biology and Agricultural Sciences*. 5(3): 173-182.
- Hemeid, N. M. (2015). Effect of different sources and levels of potassium fertilization on productivity of peanut (*Arachishypogaea* L.) grown under sandy soil conditions. *Journal of Soil Sciences and Agricultural Engineering*. 6(12): 1441-1454.
- Jena, D., Samal, D., Rath, B. S., & Venkateswarlu, B. (2006). Effect of sulphur and boron on yield, nutrient uptake and protein content of groundnut (*Arachishypogaea* L.). *Journal of Plant Nutrition*, 29(5), 805-815.
- Jyothi, T. V. and Hebsur, N. S. (2018). Effect of soil and foliar applied potassium fertilizers on yield, quality and economics of *Bt* cotton (*Gossypiumhirsutum*) in Vertisol. *Indian journal of agronomy*. 63(1): 91-94.
- Longwell, J. H.; Jeffress, R. E. and Rolph, R. L. (1963). Magnesium deficiency in cotton. *Proceedings of the Soil Science Society of America*. 27(3): 283-286.
- Munir, M. A.; Malik, M. A. and Saleem, M. F. (2007). Impact of varied levels of sulphur and magnesium on growth, yield, and oil quality of castor bean (*Ricinuscommunis* L.). *Journal of Plant Nutrition*. 30(9): 1473-1483.
- Nayak, S. C.; Pattnaik, P.; Rani, B. A.; Saravanan, S. and Babu, V. R. (2004). Response of castor (*Ricinuscommunis* L.) to potassium and magnesium fertilization. *Indian Journal of Agronomy*. 49(4): 254-257.
- Pettigrew, W. T. (2008). Potassium influences on yield and quality production for maize, wheat, soybean and cotton. *Physiologia Plantarum*. 133(4): 670-681.
- Raj, A. B.; Rao, M. S. and Shanti, M. (2004). Effect of sulphur on growth, yield and nutrient uptake by groundnut (*Arachishypogaea* L.). *Indian Journal of Plant Physiology*. 9(1): 24-30.
- Rao, K. T.; Rao, A. U. and Sekhar, D. (2013). Effect of sources and levels of sulphur on groundnut. *Journal of Academia and Industrial Research*. 2(5): 268-270.
- Rasheed, M.; Malik, A.; Khan, A. A. and Ullah, F. (2006). Effect of sulphur on the growth and yield of cotton. *Journal of Agronomy*. 5(3): 566-568.
- Read, J.J.; Reddy, R. K. and Jenkins, J. J. (2006). Yield and fibre quality of upland cotton (*Gossypium* sp.) as influenced by nitrogen and potassium nutrition. *European Journal of Agronomy*. 24(2): 282-290.
- Reddy, P. S.; Venkaiah, K.; Reddy, Y. T. N. and Singh, R. P. (2004). Effect of sulphur and magnesium on yield and quality of castor bean (*Ricinuscommunis* L.). *Indian Journal of Plant Physiology*. 9(2): 160-164.
- Safaya, N. M. (1976). Study on magnesium nutrition of cotton. *Plant and Soil*. 44(3): 693-705.
- Sagare, B. N. and Singh, R. P. (1988). Effect of nitrogen and potassium on yield and quality of castor (*Ricinuscommunis* L.). *Indian Journal of Agronomy*. 33(4): 381-384.
- Sanadi, U.; Math, K. K.; Bidari, B. I. and Yenagi, B. S. (2018). Effect of potassium nutrition on yield, quality and economics in groundnut (*Arachishypogaea* L.) in a Vertisol. *Journal of Pharmacognosy and Phytochemistry*. 7(2): 220-222.

- Sharavathi, R. S.; Hebbar, S. S. and Nanjappa, H. V. (2019). Effect of magnesium application on growth, yield, and quality of groundnut (*Arachishypogaea* L.). *International Journal of Chemical Studies*. 7(3): 1234-1237.
- Sireesha, V. and Dawson, J. (2022). Effect of potassium and magnesium on growth and yield of groundnut (*Arachishypogaea* L.). *The Pharma Innovation Journal*. 11(5): 591-594.
- Vaghasia, P.; Davariya, R.; Daki, R. and Dobariya, K. (2019) Response of castor (*Ricinuscommunis* L.) to crop geometry and potassium on growth, yield attributes and yields under irrigated condition. *The Indian Society of Oilseeds Research*. 8(1): 223-229.
- Veeramani, P.; Subrahmaniyan, K. and Ganesaraja, V. (2012). Nutrient Management for Sustainable Groundnut Productivity. *Madras Agricultural Journal*. 99(3): 55-57.

UNDER PEER REVIEW

Table 1. Impact of potassium schoenite (Mahalaabh) on growth, yield and yield attributing characters of castor.

Tr. No	Treatment details	Plant population /acre	Days to flowering	spike length (cm)	spike/pl ant	capsule/s pike	Seed index (g)	seed yield (kg/acre)
T1	RFD + MLP 6 kg/acre at 40-50& 100-110 DAS through Drip	3629	61.00	70.47	10.13	109.13	30.66	1029
T2	RFD + MLP 8 kg/acre at 40-50 & 100-110 DAS through Drip	3693	61.33	71.93	11.20	109.20	30.73	1050
T3	RFD + MLP 10 kg/acre at 40-50 & 100-110 DAS through Drip	3693	62.00	71.40	11.87	115.87	31.44	1070
T4	RFD + MLP 10 g/L at 40-50 & 100-110 DAS as foliar	3757	61.33	68.27	9.87	98.80	30.19	913
T5	RFD + MLP 15 g/L at 40-50 & 100-110 DAS as foliar	3725	63.00	71.20	9.93	99.87	30.40	973
T6	RFD + MLP 20 g/L at 40-50 & 100-110 DAS as foliar	3725	63.33	69.80	10.07	107.00	30.33	1000
T7	RFD + K-mag 50 kg/acre as basal	3757	62.33	70.47	9.60	102.13	29.93	945
T8	RFD + Mahalaabh Gr. 50 kg/acre as basal	3629	61.67	67.87	10.80	111.93	31.03	1056
T9	RFD (72:16:00:08 kg NPKS/Acre)	3661	62.67	63.87	9.47	97.67	30.11	875
S.Em.±		72.00	0.83	2.36	0.61	3.59	0.32	125.76
CD at 5%		NS	NS	NS	NS	10.77	NS	NS
CV %		3.37	2.31	5.89	10.17	5.88	1.80	12.00

Table 2. Impact of potassium schoenite (Mahalaabh) on growth, yield and yield attributing characters of cotton.

Tr. No.	Treatment details	Plant height (cm)	No. of bolls/plant	No. monopodial branch/plant	No. sympodial branch/plant	Wt. of seed cotton per boll (g)	Wt. of seed cotton per plant (g/pl.)	seed cotton yield (kg/acre)
T1	RFD + MLP 6 kg/acre at 30-35 & 50-55 DAS through Drip	109.73	40.33	2.40	12.20	3.81	155.40	1131
T2	RFD + MLP 8 kg/acre at 30-35 & 50-55 DAS through Drip	94.27	35.20	1.67	13.00	4.25	140.87	1097
T3	RFD + MLP 12.5 kg/acre at 30-35 & 50-55 DAS through Drip	96.27	33.53	2.00	11.93	4.24	123.93	1046
T4	RFD + MLP 10 g/L at 30-35 & 50-55 DAS as foliar	102.20	39.67	2.27	12.87	4.73	163.07	1148
T5	RFD + MLP 15 g/L at 30-35 & 50-55 DAS as foliar	101.80	33.73	2.00	11.47	4.25	150.80	1085
T6	RFD + MLP 20 g/L at 30-35 & 50-55 DAS as foliar	110.27	33.53	2.13	12.53	4.52	142.13	1082
T7	RFD + K-mag 25 kg/acre as basal	104.53	44.53	2.40	11.13	3.76	169.33	1210
T8	RFD + Mahalaabh Gr. 25 kg/acre as basal	94.73	34.53	1.47	12.60	4.03	148.13	962
T9	RFD (80:16:00:26 kg NPKS/Acre)	98.00	38.13	2.33	12.47	4.12	152.20	1090
S.Em. ±		4.74	3.36	0.32	0.62	0.28	13.44	74.36
CD at 5%		NS	NS	NS	NS	NS	NS	NS
CV %		8.10	13.74	12.86	8.74	11.51	12.57	11.77

Table 3. Impact of potassium schoenite (Mahalaabh) on growth, yield and yield attributing characters of groundnut.

Tr. No.	Treatment details	Plant height (cm)	Days to flowering	No. of pods/plant	No. of seeds/ pod	Seed index (g)	Pod yield (kg/acre)	Shelling %
T1	RFD + MLP 10 g/L at 30-35 & 50-55 DAS as foliar	30.67	32.67	12.27	1.87	41.91	788	63.61
T2	RFD + MLP 10 g/L at 30-35, 50-55 & 70-75 DAS as foliar	30.20	32.00	14.87	1.80	41.53	660	56.21
T3	RFD + MLP 15 g/L at 30-35 & 50-55 DAS as foliar	30.67	29.33	18.13	1.87	41.93	831	69.01
T4	RFD + MLP 15 g/L at 30-35, 50-55 & 70-75 DAS as foliar	30.53	31.67	16.40	1.87	43.05	887	61.74
T5	RFD + MLP 6 kg/acre at 30-35 & 50-55 DAS through Drip	29.13	32.33	16.93	1.80	40.83	725	65.67
T6	RFD + MLP 8 kg/acre at 30-35 & 50-55 DAS through Drip	29.93	34.67	17.07	1.87	40.37	555	57.93
T7	RFD + K-mag 25 kg/acre as basal	29.47	34.00	15.53	1.67	40.78	644	68.32
T8	RFD + Mahalaabh Gr. 25 kg/acre as basal	29.20	31.67	15.13	1.87	41.80	718	65.06
T9	RFD (05:10:12:00 kg NPKS/Acre)	29.87	32.67	13.80	1.93	41.35	722	64.14
S.Em.±		0.88	1.18	1.47	0.12	1.58	84.69	5.29
CD at 5%		NS	NS	NS	NS	NS	NS	NS
CV %		5.07	6.30	10.39	11.09	6.60	12.22	9.43




Figure 1: Effect of different Mahalaabh treatments on castor seed yield, seed cotton yield and groundnut pod yield.