

Original Research Article

Identifying the traditional and existing tree –crop combination of agroforestry system in Southern part of Chhattisgarh

Abstract

Present study was conducted in Bastar district of Chhattisgarh for identify the traditional and existing agroforestry practices by farmers. The data were collected through field visit and filled out of pre-prepared questionnaires in four blocks of Bastar district, namely Jagdalpur, Bastar, Tokapal and Bakawand. The survey revealed six different types of agroforestry system practices i.e., agrisilvihorticulture, agrisilviculture, silvihorticulture, agrisilvipastoral, silvihortipastoral, and agrisilvihortipastoral while the 67% of farmers were practices agrisilvihorticulture-type agroforestry systems. Farmers of Bastar were found using 29 different tree crops, 17 varieties of horticultural crops, 20 vegetable varieties, and 11 agricultural crops in their fields.

Key words: Agroforestry, Cropping Patterns, Sustainability, Livelihood security, Tree crop interaction, Bastar

Introduction

India contains one sixth of world population is about 1.43 billion peoples [1]. Growing population creates huge pressure on demand of foods, water, and land and also quality of air [2, 3] resulting deteriorates the natural resources. Deforestation due to the construction of roads and buildings, expansion of industries [4] and extensive agriculture are the major causes of disturbing environmental sustainability. On the other hand, the agroforestry system is playing a crucial role in minimizing food shortages through different products and providing livelihood security [5-8].

Agroforestry is a sustainable land use approach that involves consciously combining agricultural output in the lower story alongside a woody component i.e. home garden, hedge, multistoried cropping, windbreak, shelterbelt, border planting, and strip planting [9]. It defined as an agroforestry system is a collective land use where field crops are planted alongside trees and woody perennial plants [10, 11]. It has been multifunctional benefits such as control of soil erosion [12, 13], improve air quality, rich organic matter in soil [14], restore available ground water, shade for beneath growing crops, fodder for livestock, sustain the environment through conservation of biodiversity [15-18]. Adaptation of suitable agroforestry system have been depending upon the various factors i.e. topography of the farm, physiochemical properties of soil [19], agro-climatic features of the regions, availability of labour and irrigation facilities, and size of the farm etc. [20-22]. Different types of agroforestry systems practiced in the country namely agrisilviculture system

(combination of agricultural field crops + trees), silvipastoral system (livestock production + trees), agrisilvihorticulture system (field crops + horticultural crops + trees), agrisilvipastoral system (agricultural field crops + livestock production + trees), agrisilvihortipastoral system (agricultural field crop+ horticulture crops + animal husbandry + trees), and silvihorticulture system (horticultural crops + trees) [23-25].

This study was conducted in Bastar situated in southern part of Chhattisgarh (India). Bastar is world famous for its traditional tribal culture and handmade art-craft [26]. It covers 403,003 hectare of total geographical area whereas 20.83 % forest land, 17.63% uncultivated land, 10.69% agriculture land and 5.10% wasteland. The tropical moist deciduous type of forest specially dominated by Sal trees (*Shorea robusta*), while other tree species *Bauhiniavariegata*, *Tectona grandis*, *Pterocarpus marsupium*, *Terminalia tomentosa*, *Anogeissus latifolia*, *Madhuca indica*, *Diospyros melanoxylon*, *Embilica officinalis*, *Cleistanthuscollinus*, *Tamirindus indica*, *Acacia nilotica*, *Acacia catechu*, *Mangifera indica*, *Terminalia arjuna*, and *Dendrocalamus* species are present on either bund of the agricultural fields or **boundary** of the farm. Bastar region mainly covers entisol, inceptisol and alfisol types of soil and nutrient such as nitrogen, phosphorus, potassium were ranged from 188.6-276.64, 7.62-10.72, 121-242.5, kg/ha [27-29]. During monsoon season rice, maize and millets crops are major crops grown in rainfed and irrigated condition while post-monsoon and/or pre-monsoon season wheat, maize, **moongbeen**, vegetables like cucurbits, papaya, **chilli**, tomato, brinjal were grown.

Agroforestry system pushed in past three decades to developing nation as a way to raising livelihood income of the growers [5, 30]. There is no previously study in the area on which species of tree and crops combination practiced for cultivation, and how much income generated from the forest produces. Hence, this research designs for identification of tree-crop combination practiced as an agroforestry system by the farmers of Bastar.

Materials and Methods

The study was carried during the period of 2016-17 in Bastar district (Chhattisgarh), which were situated at 19°10'71" N latitude and 81°95'35" E longitude at an altitude of 850 meter with average annual rainfall and temperature was 1250 mm and 18.9°C - 34.1°C.

The data were collected by pre-prepared questionnaire with regular field visit of the villages [31] and farming system. Twelve villages were randomly selected in each blocks of Bastar district (Table 1). 25 farmers of each block were randomly selected on the basis of availability farming system with different tree-crop combination on the field. The questionnaire was contended different aspect of

agroforestry system such as available tree species in the farm, crops grown, mixed cropping techniques, utilization of tree products, income from tree produces etc.

Table 1 Blocks and villages were selected for surveyed during the study.

Sl. No.	Blocks	Villages
1	Jagdapur	Pandripani, Parpa, Pamela, Karkapal, Podaguda, Aasna, Nagrnar, Neeyanar, Ghatpadmur, Sargipal, Lamni, Burungpal
2	Bastar	Marlenga, Chokar, Mahupal, Mongpal, Bastar, Bhirlinga, Mohpal Barai, Kawadgaon, Kudkanar, Ghatlohanga, Kaviashana, Parchanpal
3	Bakawand	Bakawand, Sivnagudam, Karpawand, Devbeejaguda, Chhotedevda, Banras, Dongrepal, Sonpur, Kodawand, NarawandMatiguda, Kosmi
4	Tokapal	Tekameta, Bademorthpal, Parakuriyaguda, Bhaluguda, Kuranga, Karanji, Aavarabhata, Kodipara, Kalepal, Telimarenga, Matigudapara, Karanji, Chhotemorathpal

Results and Discussion

The result was revealed in agroforestry system practiced by the farmers of different blocks. The maximum number of farmer practiced agrisilvihorticulture system about 67 % while the 23% aagrisilviculture system, 7% silvihorticulture system, 1% agrisilvipastoral, 1% silvihortipastoral, 1% agrisilvihortipastoral system were recorded (Fig. 1). Adaptation of suitable agroforestry system have been depending upon the various factors i.e. topography of the farm, physiochemical properties of soil, agro-climatic features of the regions, availability of labour and irrigation facilities, and size of the farm etc [20]. The dependency of farmers on agrisilvihorticulture system was highest due to the their diversified produces such as tree provides wood for their house construction and making agriculture implements, crops provided food grains, and horticulture system provides fruits, flowers, vegetables etc for selling [32]. The Bastar area employs a wide range of agro - forestry approaches. The selection of tree species varies from based on site to site, need of the farmer, and their socioeconomic considerations [23, 33]. This gives several examples of where farmers in the area have successfully used traditional agro - forestry techniques and indigenous techniques and agroforestry systems plays crucial role in ensuring the sustainability in resources for livelihood [34].




Figure 1. Different agroforestry systems were used by farmers of Bastar.




Figure 2 Frequency of tree species occurred in different agroforestry system

3.2 Tree-crop combination practiced in Agroforestry System

The study was revealed the six type of agroforestry system practiced by the farmers in Bastar such as agrisilvihorticulture, agrisilviculture, silvihorticulture, agrisilvipastoral, silvihortipastoral, and agrisilvihortipastoral. The maximum variation in species of agroforestry system was recorded in agrisilvihorticulture. Whereas 29 tree species, 17 horticultural crops, 20 vegetables crop species and 11 agricultural crops were used in tree-crop combination system. *Mangifera indica* and *Eucalyptus*

tree species was reported almost all type of agroforestry system while the other trees such as *Phoenix sylvestris*, *Acacia nilotica*, *Schlecharia oleosa*, *Caryotaurens*, *Cocunucifra*, *Terminalia tomentosa*, *Psidium guajava*, *Ziziphus marutiana*, *Moringa olifera*, and *Bombax ceiba* practiced in scattered form or on the bunds in an agroforestry system. Frequency of tree species present in an agroforestry system were also recorded (Fig. 2). This study was reported the combination of agriculture crops, horticulture crops and tree species practiced on the field of farmers in Bastar region as shown in Table 2 and Fig. 3. Agroforestry system provide wide range of benefits to the farmers such as food for life, woods for furniture and construction home, fruits for nutrition, etc. Similarly investigation were reported on the horticulture based agroforestry models whereas horticulture species includes *Emblica officinalis*, *Psidium guajava*, *Punica granatum*, *Aegle marmelos*, *Z. mauritiana* combination with trees species includes *T. grandis*, *G. arborea*, *A. lebeck*, *T. arjuna* and *A. nilotica* on the field [34]. Studied in Bastar region on different aspect of agroforestry system and reported the agrisilviculture system were highly practiced with the combination of tree species like *Acacia auriculiformis*, *Psidium guava*, *Tectona grandis*, *Cocos nucifera*, with the horticultural fruits crops like *Punica granatum*, *Anacardium occidentale*, *Emblica officinalis*, *Manilkara achras*, *Mangifera indica*, *Psidium guava*, *Terminalia Arjuna*, *Tamarindus indica*, *Terminalia belerica*, *Litchi chinensis* [23]. The agroforestry systems have used to intensify in the farming system by maximizing of resources. The agroforestry system was a very promising method for Mediterranean areas with intercropping, such as growing olives with grain crops like barley and mixes of barley and leguminous plants [35]. Landscapes for agriculture have always included windbreaks as a vital component. Among its many functions are crop protections from wind, conserving habitats for biodiversity, safeguarding against soil erosion, and creating microclimates that mitigate the impact of adverse weather conditions [36, 37].

Table 2 Tree- crop combination practiced by the farmers of Bastar in various Agroforestry systems.

Sl. No.	Agroforestry System	Combination of trees, horticultural crops, agricultural crops and fodder crops on the same piece of land in same time
1	Agrisilviculture system	<i>Tectona grandis</i> + <i>Oryza sativa</i> / <i>Zea mays</i>
		<i>Tamarindus indicus</i> + <i>Eucalyptus tereticornis</i> + <i>Manihot esculenta</i>
		<i>Phoenix sylvestris</i> + <i>Oryza sativa</i>
		<i>Acacia nilotica</i> + <i>Oryza sativa</i>
		<i>Eucalyptus tereticornis</i> + <i>Oryza sativa</i>

		<i>Azadirachta indica</i> , + <i>Tamarindus indica</i> + <i>Saccharum officinarum</i>
		<i>Acacia auriculiformis</i> + <i>Zea mays</i> / <i>Oryza sativa</i> / <i>Saccharum officinarum</i>
		<i>Tamarindus indica</i> + <i>Zea mays</i> / <i>Zingiber officinales</i>
2	Silvihorticulture system	<i>Tectona grandis</i> + <i>Mangifera indica</i> / <i>Capsicum annum</i>
		<i>Syzygium cuminii</i> + <i>Capsicum annum</i>
		<i>Eucalyptus tereticornis</i> + <i>Momordica charantia</i> + <i>Abelmoscus esculentus</i>
		<i>Acacia auriculiformis</i> + <i>Abelmoscus esculentus</i>
		<i>Anacardium occidentale</i> + <i>Abelmoscus esculentus</i>
		<i>Gmelina arborea</i> + <i>Abelmoscus esculentus</i> / <i>Capsicum annum</i>
		<i>Caryotaurens</i> + <i>Cocus nucifera</i> + <i>Vigna anguiculata</i>
		<i>Tectona grandis</i> + <i>Tamarindus indicus</i> + <i>Leucaena luecocephala</i> + <i>Schleichera oleosa</i> + <i>Semecarpus anacardium</i> + <i>Litchi chinensis</i> + <i>Manilkara zapota</i> + <i>Mangifera indica</i> + <i>Syzygium cuminii</i> + <i>Psidium guajava</i> + <i>Carissa carandus</i> + <i>Capsicum annum</i> + <i>Abelmoscus esculentus</i>
		<i>Mangifera indica</i> + <i>Abelmoscus esculentus</i> + <i>Polianthes tuberosa</i> <i>Eucalyptus tereticornis</i> + <i>Acacia auriculiformis</i> + <i>Caryotaurens</i> + <i>Tamarindus indicus</i> + <i>Diospyros melonoxylon</i> + <i>Mangifera indica</i> + <i>Anacardium occidentale</i>
3	Agrisilvipastoral System	<i>Eucalyptus tereticornis</i> + <i>Leucaena leucocephala</i> + <i>Oryza sativa</i> / <i>Zea mays</i> + <i>Ipomoea batatas</i> + Nepiar grass
4	Silvihortipastoral system	<i>Eucalyptus tereticornis</i> / <i>Tamarindus indicus</i> + <i>Terminalia tomentosa</i> + <i>Mangifera indica</i> + <i>Manilkara zapota</i> + <i>Psidium guajava</i> + <i>Cynadondactylon</i> / Nepiar grass

5	Agrisilvihortipastoral system	<i>Azadirachta indica</i> + <i>Eucalyptus tereticornis</i> / <i>Diospyros melonoxylon</i> + <i>Leucaena leucocephala</i> (Fodder) + <i>Ziziphus mauritiana</i> + <i>Atrocarpus heterophyllus</i> / <i>Syzygiumcumini</i> + <i>Allium cepa</i> + <i>Abelmoscus esculentus</i> / <i>Phaseolus species</i> + <i>Oryza sativa</i> / <i>Zea mays</i> + <i>Saccharum officinarum</i> / <i>Cynadondactylon</i> (Fodder)
		<i>Tectona grandis</i> / <i>Gmelina arborea</i> + <i>Acacia auriculiformis</i> + <i>Mangifera indica</i> + <i>Moringa oleifera</i> + <i>Cocus nucifera</i> / <i>Vigna anguilata</i> + <i>Oryza sativa</i> / <i>Zea mays</i> + <i>Cynadondactylon</i> / <i>Pennisetum glaucum</i>
6	Agrisilvihorticulture system	<i>Tamarindus indicus</i> + <i>Capsicum annum</i> + <i>Zea mays</i> + <i>Helianthus annus</i>
		<i>Tamarindus indicus</i> + <i>Eucalyptus tereticornis</i> + <i>Acacia auriculiformis</i> + <i>Manngifera indica</i> + <i>Atrocarpus heterophyllus</i> + <i>Syzygiumcuminii</i> + <i>Zea mays</i>
		<i>Tamarindus indicus</i> + <i>Gmelina arborea</i> / <i>Tectona grandis</i> + <i>Cocus nucifera</i> + <i>Atrocarpus heterophyllus</i> + <i>Mangifera indica</i> + <i>Manilkara zapota</i> + <i>Oryza sativa</i> / <i>Zea mays</i>
		<i>Gmelina arborea</i> + <i>Tectona grandis</i> + <i>Eucalyptus tereticornis</i> + <i>Bombax ceiba</i> + <i>Mangifera indica</i> + <i>Anacardium occidentaleis</i> + <i>SyzygiumCuminii</i> + <i>Oryza sativa</i>



***Tectona grandis* with *Abelmoscus esculentus* in Neeyanar**



***Mangifera indica* with *Tagest species* in Pandripani**



Acacia auriculiformis with *Zea mays* in Tekameta



Tamarindus indicus with *Capsicum annum* in Aashana



Eucalyptus teretecornis with Bitter gourd in Ghatpadmur



Acacia auriculiformis with *Abelmoscus esculentus* in Karanji



Acacia auriculiformis with *Oryza sativa* in Tekameta




Mangifera indica with *Cymbopogon flexuosus* in Kodipara



Mangifera indica with *Solanum melongina* in Pandripani



Tectona grandis with *Capsicum annum* in Bastar



Anacardium occidentale with *Cymbopogon flexuosus* in Kodipara



Tamarindus indica with *Zingiber officinalis* in Kodipara



Tectona grandis with *Mangifera indica* in Mohpal Barai



Mangifera indica with *Zea mays* in Bastar



Mangifera indica with *Curcuma longa* in Sargipal



Syzygium cuminii with *Capsicum annum* in Pandripani

Figure 3 Agroforestry system existed in field of Bastar farmers.

Benefits from tree in an agroforestry system

This study found that varied uses of trees occurred on agricultural farms, whether they were naturally occurring or planted (Table 3). Farmers receive benefits from agroforestry strategies in combination with a variety of produce. During the study, it was found that most of the farmers in Bastar district expressed their desire to plant various types of tree species in their fields as per their needs and utility namely *Tamarindus indica*, *Mangifera indica*, *Atrocarpus heterophyllus*,

Anacardium occidentale, *Caryotaurens*, *Madhuca indica* and *Diospyros melonoxylon*. As estimated by CAFRI, Jhansi, there are 1,023 million ha of agroforestry globally, and 13.75 million ha of agroforestry are found in India. It indicates that hardly 10% of the country's agricultural areas are covered with agroforestry [38]. An agroforestry system produces high-quality fruits, vegetables, and lumber [39]. It is believed to be able to interact with ecosystem functions and contribute to sustainable agriculture [40]. Silvopasture is a viable agroforestry farming substitute that combines the benefits of trees (timber production) with the production from livestock, including meat and dairy [41]. Similarly, agrisilviculture models provide a higher quality of both agriculture grain and timber production, resulting in farmers receiving a higher profit per unit of land [42]. Because of the wide range of items produced, it also offers nutritional security. The normative return from cultivating vegetables is greater than that of ordinary field crops [43].

Table 3 Farmers of Bastar used of tree species for different purposes.

Tree species	Common name	Uses	Marketable value (In INR)*
<i>Mangifera indica</i>	Mango	Fuel wood, fruit	80-100/ kg
<i>Leucaena leucocephala</i>	Subabul	Fodder, fuel wood	-
<i>Syzygium Cumini</i>	Jamun	Fruit	Dried seeds- 42/ kg
<i>Bombax ceiba</i>	Semul	Timber, fibre	-
<i>Acacia auriculiformis</i>	Australian babul	Timber, fuel wood	-
<i>Diospyros melonoxylon</i>	Tendu	Fruit, timber, fuel wood, tendu patta	4000/standard bag
<i>Azadirachta indica</i>	Neem	Timber, medicinal value, Seeds	Seeds- 27/ kg
<i>Ziziphus mauritiana</i>	Ber	Timber, fruit, fodder	40-80/ kg
<i>Terminalia tomentosa</i>	Saja (Aasan)	Timber, silk rearing, fuel wood	Silk rearing- 400/ 100 pieces
<i>Tectona grandis</i>	Teak	Timber	-
<i>Gmelina arborea</i>	Khamhar	Timber, fodder	-
<i>Eucalyptus tereticornis</i>	Nilgiri	Timber, oil extraction	-
<i>Cocos nucifera</i>	Coconut	Fruit	50/ piece

<i>Moringa oleifera</i>	Drum stick	Fruit	50-80/ kg
<i>Atrocarpus heterophyllus</i>	Jack fruit	Fruit, fuel wood	50-60/ kg
<i>Tamarindus indicus</i>	Imli	Fruit pulp, fuel wood, seeds, flower	Seeds- 36/ kg Flower- 63/ kg Pulp- 35-40/ kg
<i>Psidium guajava</i>	Guava	Fruit	50-80/ kg
<i>Schleichera oleosa</i>	Kusum	NWFP, Lac production	Lac- 300-400/ kg
<i>Semecarpus anacardium</i>	Bhelva	Fuelwood, fruit	9/ kg
<i>Manilkara zapota</i>	Sapota	Fruit	80-100/ kg
<i>Litchi chinensis</i>	Litchi	Fruit	80-100/ kg
<i>Phoenix sylvestris</i>	Chhind or silver date palm	Fruit, beverage	Beverage- 40.00/ litre
<i>Madhucaindia</i>	Mahua	Seeds, Flower, beverage	Seeds -29/ kg Flower-30/ kg Beverage – 60/ litre
<i>Caryotaurens</i>	Sulfi	Beverage	80-10/ litre
<i>Shorea robusta</i>	Sal	Seeds	20/ kg

*Source of marketable value – local market and Chhattisgarh minor forest produce limited [44].

Conclusion

Agroforestry is a dynamic and sustainable land management system that involves deliberately growing of woody perennials along with agricultural crops on farmlands to secure both direct and indirect benefits for farmers. This study reported the identification and documentation of patterns of tree-crop combinations used in the agroforestry system by the farmers of Bastar District. The agrisilviculture agroforestry system was mostly practiced throughout the Bastar district, and *Tectona grandis*, *Shorea robusta*, *Eucalyptus tereticornis*, *Phenix sylvestris*, *Mangifera indica*, *Syzygiumcumini*, *Gmelina arborea*, *Pongamia pinnata*, *Tamarindus indica*, *Bambosabambos*, *Anacardium occidentale*, *Terminalia arjuna*, *Terminalia balerica*, *Madhuca indica*, *Acacia auriculiformis*, and *Acacia nilotica* tree species were grown by the farmers on their field.

References

1. Jose, S., Gulati, A., Khurana, K. Achieving nutritional security in India: Vision 2030. Indian Council for Research on International Economic Relations. 2020.
2. Islam, S. M. F., Karim, Z. World's demand for food and water: The consequences of climate change. *Desalination-challenges and opportunities*, 2019; 1-27.
3. **Tiwari**, Deepti, Manoj Kumar Patel, Payal Priyadarsini, Ankita, Shatakashi Mishra, Vijai P., UP Singh. Physiological Responses of Stress Tolerant Rice Varieties across Different Cropping Systems under Rainfed Stress. *International Journal of Environment and Climate Change*. 2024; 14 (5):56-66. <https://doi.org/10.9734/ijecc/2024/v14i54170>
4. Okechukwu Ucheje, Okoroafor, Obiageli, Jacinta Okolo, Okonma, Mark Chukwueloka. Effect of Climate Change on Nigerian Economic Sustainability. *International Journal of Environment and Climate Change*. 2024; 14 (5):67-78. <https://doi.org/10.9734/ijecc/2024/v14i54171>
5. Kuyah, S., Sileshi, G. W., Luedeling, E., Akinnifesi, F. K., Whitney, C. W., Bayala, J., Mafongoya, P. L. Potential of agroforestry to enhance livelihood security in Africa. *Agroforestry for Degraded Landscapes: Recent Advances and Emerging Challenges*. 2020; 1: 135-167 https://doi.org/10.1007/978-981-15-4136-0_4
6. Pius, Moses Nnah, S. A. Yelwa, A. Jibrillah, U. Umar. Time Series Analysis of Climate Change, Drought and Food Security Nexus in Kaduna State Nigeria. *Journal of Geography, Environment and Earth Science International*. 2023; 27 (12):76-88. <https://doi.org/10.9734/jgeesi/2023/v27i12738>
7. Kavitha , S., G. Samuel, I. Sreenivasa Rao, M. Goverdhan, D. Srinivasa Chary. Integrated Farming System for a Sustainable Livelihood. *International Journal of Environment and Climate Change*. 2024; 14 (6):165-69. <https://doi.org/10.9734/ijecc/2024/v14i64218>
8. Kamesh, Singh, B.P., Misra, S., Verma, K.K., Singh C.K., Kumar, R. An emerging adsorption technology and its applicability on trees as an adsorbent for the remediation of water pollution: A review. *Ecology Environment and Conservation*. 2023; 29(02):627-640 DOI:10.53550/EEC.2023.v29i02.014
9. Mosquera-Losada MR, Borek R, Balaguer F, Mezzrila G, Ramos-Font E. Agroforestry as a mitigation and adaptation tool, EIP-Agri Focus Gropu agroforestry. 2017 https://ec.europa.eu/eip/agriculture/sites/default/files/fg22_mp9_cc_adaptation_mitigation_2017_en.pdf
10. Gupta, S. R., Dagar, J. C., Teketay, D. Agroforestry for rehabilitation of degraded landscapes: achieving livelihood and environmental security. *Agroforestry for Degraded Landscapes:*

Recent Advances and Emerging Challenges. 2020; 1, 23-68. https://doi.org/10.1007/978-981-15-4136-0_2

11. Lojka, B., Teutscherová, N., Chládová, A., Kala, L., Szabó, P., Martiník, A., Lawson, G. Agroforestry in the Czech Republic: what hampers the comeback of a once traditional land use system?. *Agronomy*, 2021; 12 (1), 69. <https://doi.org/10.3390/agronomy12010069>
12. Béliveau, A., Lucotte, M., Davidson, R., Paquet, S., Mertens, F., Passos, C. J., Romana, C. A. Reduction of soil erosion and mercury losses in agroforestry systems compared to forests and cultivated fields in the Brazilian Amazon. *Journal of environmental management*. 2017; 203, 522-532. (Soil erosion) <https://doi.org/10.1016/j.jenvman.2017.07.037>
13. Kale, V. B. Estimation and Assessment of Soil Erosion Risk of Nun Watershed Using Remote Sensing and GIS. *Journal of Geography, Environment and Earth Science International*. 2019;19 (1):1-14. <https://doi.org/10.9734/jgeesi/2019/v19i130078>
14. Lakshman, C.H. Raval, H. K. Patel, Suwa Lal Yadav, Devilal Birla, Priyanka Choudhary, Manju Choudhary. Effect of Organic Sources on Properties of Soil on the Cultivation of Greengram (*Vigna Radiate L.*) in Sandy Loam Soil of Middle Gujrat Region. *International Journal of Environment and Climate Change*. 2024; 14 (7):76-82. <https://doi.org/10.9734/ijecc/2024/v14i74253>
15. Udawatta, R. P., Rankoth, L. M., Jose, S. Agroforestry and biodiversity. *Sustainability*. 2019; 11 (10), 2879. <https://doi.org/10.3390/su11102879>
16. Patel, Krishna Kumar, Alok Kumar Pandey, Ajay Kumar Baheliya, Veerendra Singh, Rishikesh Yadav, Shubham Pandey, Adesh, Vikash Yadav. A Comparative Study of the Effects of Biofertilizers and Chemical Fertilizers on Soil Physical and Biological Properties under Chickpea Crop (*Cicer Arietinum L.*). *International Journal of Environment and Climate Change*. 2024; 14 (3):72-80. <https://doi.org/10.9734/ijecc/2024/v14i34020>
17. Tirkey , Sandhya, Brijendra Pratap Singh, Kamesh, Shailly Misra, Pinaki Prasad Singh, Ruby, Tulika J. Evaluation of Physio-Chemical Properties and WQI of the Drinking Water in Urban Area of Bilaspur, Chhattisgarh, India. *Asian Journal of Environment & Ecology*. 2024; 23 (7):189-99. <https://doi.org/10.9734/ajee/2024/v23i7574>
18. Shivam Dinkar, Brijendra Pratap Singh, Kamesh, Shailly Misra, Chandresh Kumar Singh, Tulika Johri, Ramesh and Rajesh K. Assessment of the Air pollution tolerance capacity of trees for development of greenery around the industrial area. *Ecology, Environment & Conservation*. 2024; 30- S165-S169. DOI:[10.53550/eec.2024.v30i01s.029](https://doi.org/10.53550/eec.2024.v30i01s.029)

19. Navyashree R, Mummigatti UV, Nethra P, Basavaraj B, Hanamaratti NG. Evaluating the Effects of Drought Stress on Biomass and Yield Traits in Sorghum Genotypes”. *International Journal of Environment and Climate Change*. 2024; 14(5):79-87. <https://doi.org/10.9734/ijecc/2024/v14i54172>
20. Pantera, A, Mosquera-Losada, MR, Herzog, F. Agroforestry and the environment. *Agroforest Syst*. 2021; 95: 767–774. <https://doi.org/10.1007/s10457-021-00640-8>
21. Burgess, A. J., Cano, M. E. C., Parkes, B. The deployment of intercropping and agroforestry as adaptation to climate change. *Crop and Environment*. 2022; 1(2), 145-160. <https://doi.org/10.1016/j.crope.2022.05.001>
22. Singh, V., Johar, V., Kumar, R., Chaudhary, M. Socio-economic and environmental assets sustainability by agroforestry systems: a review. 2021; 14 (4): 521-533. <http://dx.doi.org/10.30954/0974-1712.04.2021.6>
23. Hemrom, A, Nema, S, A study on traditional agroforestry practices existing at Bastar region of Chhattisgarh, *International Journal of Multidisciplinary Research and Development*, 2015; 2 (3): 56-64.
24. Singh, BK, Oraon, PR, Ranjan, S, Kumar, A, Suman, KK. Survey of existing farming system and agroforestry practices in the Lohardaga District, Jharkhand, India. *International Journal of Current Microbiology and Applied Sciences*. 2019;8 (9), 1625-1633. <https://doi.org/10.20546/ijcmas.2019.809.185>
25. Sahoo, G., Wani, A. M. Multifunctional agroforestry systems in India for livelihoods. *Annals of Horticulture*. 2019; 12 (2), 139-149. [10.5958/0976-4623.2019.00022.7](https://doi.org/10.5958/0976-4623.2019.00022.7)
26. Chandravanshi, J., Jain, N. Indigenous Art of Bastar District of Chhattisgarh: A Study of the Government Initiations for Artisans. 2023; 10 (1): 25 [10.5958/2583-3561.2023.00002.4](https://doi.org/10.5958/2583-3561.2023.00002.4)
27. Rao, S.P., Thomas, T., Singh, Z., Rachana Assessment of macronutrients in soils of Bastar Plateau Region, Chhattisgarh, India, *International Journal of Multidisciplinary Research and Development*. 2016; 3 (7): 34-38.
28. Mishra, P., Upadhyay, M., Khare, G. P. Estimation of Macronutrients in Soil of Bastar Plateau Region Chhattisgarh, India. *Journal of Chemistry and Chemical Sciences*. 2019; 9 (12): 326-330
29. Sahu, P., Chandrakar, T., Joshi, K. K., Mishra, R. Effect of organic and inorganic sources of nutrients on nutrient content and uptake of direct seeded rice in Inceptisols of Bastar plateau zone. *The Pharma Innovation*. 2020; 9(5): 285-289

30. Karthick, R., K. Arulmani, K. Ravi. Impact of Climate Change Assessments on Horticultural Crop Dynamics in Tamil Nadu, India. *International Journal of Environment and Climate Change*. 2024; 14 (3):114-25. <https://doi.org/10.9734/ijecc/2024/v14i34024>
31. Okamoto, I. Organizing community forestry in rural Myanmar: capability and functions of villages. In *Local Societies and Rural Development*. 2014; 67-100. Edward Elgar Publishing. <https://doi.org/10.4337/9781783474387.00010>
32. Nwosisi, S, Nandwani, D. Urban Horticulture: Overview of Recent Developments. In: Nandwani, D. (eds) *Urban Horticulture. Sustainable Development and Biodiversity*. 2018; 18. Springer, Cham. https://doi.org/10.1007/978-3-319-67017-1_1
33. Himshikha, Negi, AK, Poonia, PK, Kumar, M, Daneva, V. Evaluation of existing agroforestry systems in plain region of Uttarakhand, India. *Indian Journal of Agroforestry*, 2020; 22 (2), 44-50.
<https://www.indianjournals.com/ijor.aspx?target=ijor:ijaf&volume=22&issue=2&article=008>
34. Toppo, P, Raj, A, Jhariya, MK. Wild edible plants of Dhamtari district of Chhattisgarh, India. *Van Sangyan*, 2016; 3(4): 1-6.
35. Mantzanas K, Pantera A, Koutsoulis D, Papadopoulos A, Kapsalis D, Ispikoudis S, Fotiadis G, Sidiropoulou A, Papanastasis VP. Intercrop of olive trees with cereals and legumes in Chalkidiki, Northern Greece. *Agroforest Syst*. 2021 <https://doi.org/10.1007/s10457-021-00618-6>
36. Yukhnovskyi V, Polishchuk O, Lobchenko G, Khryk V, Levandovska S. Aerodynamic properties of windbreaks of various designs formed by thinning in central Ukraine. *Agroforest Syst*. 2021; 95, 855-865. <https://doi.org/10.1007/s10457-020-00503-8>
37. Dollinger J., Jose S. Agroforestry for Soil Health. *Agroforest Syst*. 2018; 92:213–219. <https://doi.org/10.1007/s10457-018-0223-9>
38. Chandra, K. K., Bhardwaj, A. K., Bhadouria, R. Wood production potential of trees in traditional agroforestry in eastern Chhattisgarh: implication for wood on every farm. *Indian Journal of Agroforestry*. 2022; 24 (1): 100-109. <https://epubs.icar.org.in/index.php/IJA/article/view/125568/47372>
39. Viswanath, S., Lubina, P.A. Traditional Agroforestry Systems. In: Dagar, J., Tewari, V. (eds) *Agroforestry*. Springer, Singapore. 2017; 91-119. https://doi.org/10.1007/978-981-10-7650-3_3

40. Gikas GD, Tsihrintzis VA, Sykas D. Effect of trees on the reduction of nutrient concentrations in the soils of cultivated areas. *Environ Monit Assess.* 2016; 188 (6):327. <https://doi.org/10.1007/s10661-016-5325-0>
41. Jose S., Dollinger J. Silvopasture: a sustainable livestock production system. *Agroforest Syst.* 2019; 93:1 <https://doi.org/10.1007/s10457-019-00366-8>
42. Tewari, J., Ram, M., Roy, M., Dagar, J.C. Livelihood Improvements and Climate Change Adaptations Through Agroforestry in Hot Arid Environments. In: Dagar, J., Singh, A., Arunachalam, A. (eds) *Agroforestry Systems in India: Livelihood Security & Ecosystem Services. Advances in Agroforestry.* 2014; 10. Springer, New Delhi. https://doi.org/10.1007/978-81-322-1662-9_6
43. Handa, A. K., Toky, O. P., Dhyani, S. K., Chavan, S. B., Toky, I. D. Innovative agroforestry for livelihood security in India. *World Agriculture.* 2016; 7, 7-16.
44. Chhattisgarh state minor forest produces <https://www.cgmpfed.org/new/>