

Perception and acceptance of Human Papilloma Virus Vaccine by Mothers' in Owerri, Nigeria: A cross sectional study

Abstract

Introduction: Uptake of human papilloma virus (HPV) vaccine among female children 9-14 years is critical to reducing the burden of HPV-related cancers in Nigeria. We assessed factors affecting caregivers' acceptance of HPV vaccination for their wards, using a structured, pretested, self-administered questionnaire in a face-to-face interview amongst caregivers attending a well child/immunization clinic in Owerri Nigeria.

Methodology: Participants were recruited by convenience sampling method between August and October 2023. We analyzed the cross-sectional data from 267 caregivers of caregivers aged 20–57 years in a Nigerian Teaching hospital. Statistical analysis included descriptive statistics and bivariate analysis to investigate the factors associated with HPV vaccination uptake.

Among the respondents, 132 (49.4%) of them had heard about HPV and 52 (19.5%) stated that it could prevent cervical cancer. Bivariate analysis was used to assess the relationship(s) between the caregivers' perception of the likelihood of their wards getting vaccinated, knowledge of HPV vaccine and social demographics. Tertiary education in the mother (OR 2.25, 95% CI 1.27 to 4.02) and father (OR 2.25, 95% CI 1.28 to 3.85) were significantly associated with their knowledge about HPV vaccine and their wards probability of taking the HPV vaccine. The participants' intention to give HPV vaccination to their children was 77.5%.

Our findings suggest that interventions tailored to enhance education, and support among caregivers could significantly increase HPV vaccine uptake among girls in Nigeria.

Conclusion: Being well-educated appears to guarantee a good knowledge of HPV vaccine but not its prevention of cervical cancer. Our respondents' intention to give HPV vaccine to their children was high. Concerted efforts need to be made by government and health workers to create awareness about HPV infection and its vaccine to caregivers and enhance HPV vaccine uptake.

Keywords: Knowledge, Attitude, Human Papilloma Virus, Vaccination.

INTRODUCTION:

Persistent Human Papilloma Virus (HPV) infection causes nearly all cases of cancer of the cervix; it is also responsible for a number of anogenital and oropharyngeal cancers.¹The HPV is mainly transmitted sexually,with most infectionsacquiresoon after becoming sexually active;² however, mother to child transmission can occur during childbirth.³ Over 100 HPV serotypes exist; persistent infection with high-risk HPV serotypes 16 and 18 have been found to be responsible for approximately 70% of all cases of cancer of the cervix worldwide.⁴

Cervical cancer is the fourth most common cancer type in women worldwide, accounting for 604,127 cases in 2020, while it is also the fourth cause of cancer death in women globally with341,831 deaths.⁵In Nigeria, it is the second most common type of cancer amongst 15 to 44-year-oldwomen, with about 14,943 new cases diagnosed annually⁴and 8000 deaths⁶and beingthe second most frequent cause of cancer deaths.⁷Cancer of the cervix and diseases associated with HPV,⁸ can be prevented by HPV vaccination programmes.Vaccination against HPV is alluded to be one of the most valuable practices against HPV,^{9,10}with available data confirming its safety profile.^{11,12} Unfortunately, the uptake of both HPV vaccination and cervical screening services are still rudimentary in Nigeria.¹³

Nigeria, along with various other countries, has added HPV vaccination – particularly, the bivalent HPV vaccine (Gardasil) for subtypes 16 and 18 – to its National Programme of Immunization (NPI) for children 9–14years old and has made this free for its citizens.⁶ It is not yet available for boys in the NPI although itis available in private health facilities for girls, boys and women, but at a largely unaffordable cost to majority of Nigerian.¹⁴

To achieve optimum cost-effectiveness of HPV vaccination, more than 70% of the female population must be vaccinated.¹⁵Unfortunately, the knowledge of HPV as a cause of cervical cancer and HPV vaccination as a prevention against cervical cancer, including the uptake of vaccination by the target population of young people, is still abysmally low.² These low rates can be due to socioeconomic, cultural and religious factors and inequitable access.¹⁶⁻¹⁸Studies have shown that parents and caregivers of young children have poor knowledge of cervical cancer prevention and screening for their wards in Nigeria.^{19, 20}To reverse this trend of low rates of vaccine acceptance and uptake, parents need to key into the concept of vaccination against HPVas acceptance by parents with appropriateknowledge is the main driver towards increasing vaccination coverage rates.

Therefore, to advance awareness and knowledge of HPV infection and HPV vaccination among caregivers of children (parents), we sought to assess the knowledge of human papilloma virus and attitude of mothers towards uptake of HPV vaccine for their children.

Methodology

We conducted a descriptive, cross-sectional, health institution-based survey involving 267 female caregivers at the Well-Child and Immunization clinic of the Federal Teaching Hospital (FTH), Owerri, between August and October 2023. The FTH is a 624-bed tertiary hospital that offers maternal and child health services and serves as a referral centre for general hospitals, maternity facilities and private hospitals within Owerri and its adjoining states. A convenience sampling technique was utilized in obtaining data from all consecutive parents who attended the well child clinic, using a questionnaire that covered baseline characteristics, knowledge and attitude about HPV and the HPV vaccine. Questionnaires were administered after obtaining informed consent from the parents, who were encouraged to ask for explanation on any of the questions from the researchers, but conversation amongst them was discouraged while completing the questionnaires. Question was interpreted into the local language if participants sought clarification, without further explanations to remove bias. Ethical approval was obtained from the Ethical Committee of the institution. The SPSS version 25.0 (IBM Corp., USA) statistical software was used for data entry, validation and analysis. Pearson's Chi-square (χ^2) statistic was used to examine the associations between categorical variables and effect estimates were presented as odds ratios along with their 95% confidence intervals. Relationships were said to be statistically significant when $p < 0.05$.

Results:

Questionnaires were administered to 302 respondents and 35 out of the 302 questionnaires were incompletely filled and not analysed further. Only 267 questionnaires were analysed, giving an effective response rate of 88.4%.

The key characteristics of the respondents are presented in Table 1. The mean age of the 267 participants was 31.0 ± 5.7 years; with a range of 20 – 52 years. One hundred and twenty-three (46.1%) respondents were aged 29 – 36 years and made up the modal age group. Most of the

respondents had received a university education (198, 74.2%) and a majority of them belonged to social classes II or III (180, 67.4%).

Analysis of the 267 caregivers revealed that 132 (49.4%) had heard about the HPV vaccine while 135 (50.6%) had not. The major source of information about the vaccine was from health workers in the hospital [79, 46.5%], the internet/social media [39, 23.0%], and the radio/TV [16(9.4%)] (Table 2). A significant proportion of the mothers, 52 (39.4%) stated that HPV vaccination prevents cervical cancer, 118 (89.2%) opined that it was given to prevent infection/disease and 96 (36.0%) did not know why it was given. One (0.4%) respondent declined comment.

When asked about being vaccinated, only twenty-three (9.6%) mothers had received the HPV vaccine while less than one-tenth [22, 8.2%] of the respondents affirmed that their children had received the vaccine. The vaccine was mainly administered to their children at a cost in a government hospital [14, 53.8%]; this was followed by 8(30.8%) that received the HPV at a health centre while 4 (15.4%) admitted receiving it at a private health facility.

Of the respondents, 95 (35.6%) stated that boys could be vaccinated against HPV, 165 (61.8%) stated otherwise while 3 (1.1%) were undecided. Four (1.5%) respondents declined comment. A majority of respondents 207 (77.5%) stated that they wanted their children to receive the vaccine, 55 (20.6%) did not wish their children to be vaccinated while 3 (1.1%) were undecided. Two (0.7%) respondents did not state their preference for HPV vaccination of their children. The reasons those respondents who did not want their children vaccinated against HPV is expressed in Figure 1 below.

On bivariate analysis, age, occupation (father and mother), education (father and mother), religion, and state of domicile of the study participants were not found to be associated with the mothers' intention to have their children vaccinated with the HPV vaccine (Table 3).

Educational class significantly affected knowledge of HPV/vaccinations as respondents with a university degree or more were about 2.3 times more likely to know about it than their counterparts without a university degree; this was similar for both the mother's (OR 2.25,

95%CI: 1.27 to 4.02) and the father's (OR 2.25, 95%CI: 1.28 to 3.85) level of education. There was no significant difference in the knowledge of HPV/vaccine by age, state of domicile religion or occupation of the respondents (Table 4).

DISCUSSION

In Nigeria and many sub-Saharan countries, HPV infection/cancer of the cervix are still of serious public health concern, in spite of development of a safe and effective preventive vaccine. Broad acceptance of the vaccine by the all and sundry is yet to be attained, even as HPV infection/cancer of the cervix remains a foremost cause of morbidity and mortality.

This study demonstrated that HPV knowledge is fairly common (49.4%) amongst the participants (mothers), with a 9.6% uptake of HPV vaccine among the mothers and 8.2% uptake amongst their children. This finding, among literate women (74.2% of respondents) is a pointer to the fact that the state of affairs may be worse in the wider population. According to the analysis, only 19.5% were knowledgeable about HPV being the cause of cervical cancer, comparable to the 14.8%, 11.1% as documented in Benin² and Lagos,²¹ Nigeria. This contrasts with the 43% knowledge of HPV among Greek adolescents²² and 99.4% among mothers of adolescents.²³ The poor knowledge of HPV infection as a cause of cancer of the cervix displayed by the mothers could potentially jeopardize the HPV infection/ cervical cancer prevention endeavour, necessitating the need for early HPV /vaccine education be initiated/ enhanced and made to be all-inclusive (children, adolescents, young people and adult males and females).

Another striking finding relating to knowledge of HPV vaccination is that a significant association between a higher level of education and knowledge of HPV was demonstrated, as was reported by Makwe and coworkers.²⁴ In this study, parents with at least a university degree were about 2.3 times more likely to know about HPV/vaccine than their counterparts without a university degree.

It is not amazing that parents mentioned health workers in the hospital as their source of information about HPV vaccines. This is akin to the findings by Naoum P et al.²⁴ This finding accentuates the need for health workers in the hospital to be armed with more information, considering their role in the fight against HPV infection and cervical cancer.

Also noteworthy was the low uptake of HPV vaccine among parents (9.6%) and their children (8.2%). Several workers had reported similar low uptake values ranging from 2.5 – 6.9%.^{20, 24–26} This contrasts with high uptake rates of up to 47% reported from America.²⁷ Cost, availability of vaccines and inequity are some factors that may militate against HPV vaccine uptake in Nigeria.²⁸ Majority of mothers, 77.5% were willing to have HPV vaccine administered to their children; this is in consonance with study(s) by several workers.^{20,21} However, parents who were hesitant to have the HPV vaccine administered to their children cited a lack of knowledge (63%) as their main reason, contrasting with safety concerns/side effects documented in another study as the main reason for hesitancy;²⁹ while a fear of side effects (13%) was the second most common reason for the vaccine hesitancy identified in this survey. Other reasons adduced for vaccine hesitancy in this study were the fathers' decision (5%) and a belief of not having any need for it (4%). Accordingly, the largely optimistic outlook (77.5% willing to have HPV vaccine administered to their wards) exhibited by this study participants is quite encouraging and should be taken advantage of, to improve HPV vaccine uptake. Furthermore, lack of knowledge of the vaccine (63%) provides an opening for healthworkers and policy makers to exploit and accentuate advocacy and education of the populace for improved HPV vaccine uptake. This is so, because health workers (physician) recommendation has been found to be the factor with the highest positive association with the decision to vaccinate children with the HPV vaccine.²³

Pertaining mother's intention to have their children vaccinated with the HPV vaccine, the survey revealed that parents' age, occupation, education, religion, and state of domicile of the study participants were not found to be significantly associated. This is in contrast with findings by other workers that documented positive correlation between gender (sex) and vaccination intentions.³⁰ Despite the intention to receive HPV vaccine not demonstrating any significant association in this study, its (i.e. intention to vaccinate their children) high value amongst mothers suggests that free HPV vaccination/inclusion in the national programme on immunization, if implemented, holds promise of a huge success in aiding effort(s) aimed at reducing the burden of cervical carcinoma in Nigeria.

Conclusion

Given that HPV vaccination considerably reduces cervical cancer and death resulting therefrom, special consideration should be given to the process(s) that would enhance provision of valid information to parents/caregivers. This information; should be conveyed in a straightforward exhaustive and easy to understand manner to parents and general populace regarding recommendation for HPV vaccine and justification based on research. Instruction on HPV infection and cervical cancer should be encouraged at all levels of education in the Nigeria to boost awareness while serving as an underpinning for better knowledge of HPV/vaccine, cervical cancer and preventing deaths from cervical cancer. This will facilitate reduction in the enormous resources from the health budget that would have been spent on the management of cancer of the cervix.

Limitations

Recall bias may not be excluded as some of the questions required recall of past events, while social desirability bias cannot be excluded because of the urge to be seen to flow with the trend.

REFERENCES

1. deMartel C, PlummerM, Vignat J, Franceschi S. Worldwide burden of cancer attributable to HPV by site, country and HPV type. *Int J Cancer*. (2017) 141:664–70. doi: 10.1002/ijc.30716
2. Isara AR, Osayi N. Knowledge of Human Papillomavirus and Uptake of its Vaccine among Female Undergraduate Students of Ambrose Alli University, Ekpoma, Nigeria. *Journal of Community Medicine and Primary Health Care*. 33 (1) 64-75. doi.org/10.4314/jcmphc.v33i1.6
3. Trottier H, Mayrand MH, Coutlée F, Monnier P, Laporte L, Niyibizi J, et al. Human papillomavirus (HPV) perinatal transmission and risk of HPV persistence among children: design, methods and preliminary results of the HERITAGE study. *Papillomavirus Res*. (2016) 2:145–52. doi: 10.1016/j.pvr.2016. 07.001
4. Bruni L, Albero G, Serrano B, Mena M, Gómez D, Muñoz J *et al*. ICO/IARC Information Centre on HPV and Cancer (HPV Information Centre). Human Papillomavirus and related diseases in Nigeria. Summary Report 17 June 2019. [cited July 3, 2020]. Available from: <https://www.hpvcentre.net/statistics/reports/XWX.pdf>.

5. IARC. Cancer Today—Worldwide. Available online at: [https://gco.iarc.fr/ today/online-analysis-table?v=2018&mode=cancer&mode_population=continents&population=900&populations=&key=asr&sex=2&cancer=39&type=0&statistic=5&prevalence=0&population_group=0&ages_group%5B%5D=0&ages_group%5B%5D=17&group_cancer=1&incl](https://gco.iarc.fr/today/online-analysis-table?v=2018&mode=cancer&mode_population=continents&population=900&populations=&key=asr&sex=2&cancer=39&type=0&statistic=5&prevalence=0&population_group=0&ages_group%5B%5D=0&ages_group%5B%5D=17&group_cancer=1&incl) (accessed March 1,2024).
6. World Health Organization. www.afro.who.int/countries/nigeria/news/nigeria-vaccinate-77-million-girls-against-leading-cause-cervical-cancer
7. World Health Organization. Nigeria's call to action - Time to Eliminate Cervical Cancer in Nigeria. 2020. [cited August 20, 2020]. Available from: Available from: <https://www.afro.who.int/news/nigerias-call-action-time-eliminate-cervical-cancer-nigeria>.
8. Patel C, Brotherton JM, Pillsbury A, Jayasinghe S, Donovan B, Macartney K, et al. The impact of 10 years of human papillomavirus (HPV) vaccination in Australia: what additional disease burden will a nonavalent vaccine prevent? *Euro Surveill.* (2018) 23:1700737. doi: 10.2807/1560-7917.ES.2018.23.41.1700737
9. Markowitz LE, Dunne EF, Saraiya M, Lawson HW, Chesson H, Unger ER, et al. Quadrivalent human papillomavirus vaccine: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep.* (2007) 56:1–24. doi: 10.1037/e601292007-001 6.
10. Kash N, Lee MA, Kollipara R, Downing C, Guidry J, Tyring KS, et al. Safety and efficacy data on vaccines and immunization to human papillomavirus. *J Clin Med.* (2015) 4:614–33. doi: 10.3390/jcm4040614
11. Basu P, Banerjee D, Singh P, Bhattacharya C, Biswas J. Efficacy and safety of human papillomavirus vaccine for primary prevention of cervical cancer: a review of evidence from phase III trials and national programs. *South Asian J Cancer.* (2013) 2:187–92. doi: 10.4103/2278-330X.119877
12. Arbyn M, Xu L. Efficacy and safety of prophylactic HPV vaccines. A cochrane review of randomized trials. *Expert Rev Vaccines.* (2018) 17:1085–91. doi: 10.1080/14760584.2018.1548282

13. Ferlay J, Shin H, Bray F, Forman D, Mathers C, Parkin D. Estimates of worldwide burden of cancer in 2008. *Int J Cancer Prevention*. 2010; 127: 2893-2917.
14. Igomu T, Folorunsho-Francis A. HPV vaccine, the cervical cancer prevention most Nigerian women don't know. *Punch Newspapers*. [cited March 9, 2024]. Available at: <https://healthwise.punchng.com/hpvpvaccine-the-cervical-cancer-preventionmost-nigerian-women-dont-know>.
15. Canfell K, Chesson H, Kulasingam SL, Berkhof J, Diaz M, Kim JJ. Modeling preventative strategies against human papillomavirus-related disease in developed countries. *Vaccine*. (2012) 30:F157–67. doi: 10.1016/j.vaccine.2012.06.09.
16. Barnes KL, VanWormer JJ, Stokley S, Vickers ER, McLean HQ, Belongia EA, et al. Determinants of human papillomavirus vaccine attitudes: an interview of Wisconsin parents. *BMC Public Health*. (2018) 18:746. doi: 10.1186/s12889-018-5635-y
17. Maness SB, Thompson EL. Social determinants of human papillomavirus vaccine uptake: an assessment of publicly available data. *Public Health Rep*. (2019) 134:264–73. doi: 10.1177/0033354919838219
18. Thompson EL, Rosen BL, Maness SB. Social determinants of health and human papillomavirus vaccination among young adults, national health interview survey 2016. *J Community Health*. (2019) 44:149–58. doi: 10.1007/s10900-018-0565-2
19. Ifedioha CO, Azuike AC. Knowledge and attitudes about cervical cancer and its prevention among female secondary school students in Nigeria. *Trop Med Int Health*. 2018; 23(7): 714-723.
20. Azuogu BN, Umeokonkwo CD, Azuogu VC, Onwe OE, Okedo-Alex IN, Egbuji CC. Appraisal of willingness to vaccinate daughters with human papillomavirus vaccine and cervical cancer screening uptake among mothers of adolescent students in Abakaliki, Nigeria. *Niger J Clin Pract* 2019; 22: 1286-1291.
21. Makwe CC, Anorlu RI, Odeyemi KA. Human papillomavirus (HPV) infection and vaccines: Knowledge, attitude and perception among female students at the University of Lagos, Lagos, Nigeria. *Journal of Epidemiology and Global Health*. 2012; 2: 199-206.
22. Vaidakis D, Moustaki I, Zervas I, Barbouni A, Merakou K, Chrysi MS, et al. Knowledge of Greek adolescents on human papilloma virus (HPV) and vaccination: a national epidemiologic study. *Medicine*. (2017) 96:e5287. doi: 10.1097/MD.0000000000005287

23. Naoum P, Athanasakis K, Zavras D, Kyriopoulos J and Pavi E. Knowledge, perceptions and attitudes toward HPV vaccination: A survey on parents of girls Aged 11–18 Years Old in Greece. *Front. Glob. Womens Health* 3:871090. doi: 10.3389/fgwh.2022.871090
24. Oluwole EO, Idowu OM, Adejimi AA, Balogun MR, Osanyin GE. Knowledge, attitude and uptake of human papillomavirus vaccination among female undergraduates in Lagos State, Nigeria. *J Family Med Prim Care*. 2019; 8(11): 3627-3633.
25. Onowhakpor AO, Omuemu VO, Osagie OL, Odili CG. Human papillomavirus vaccination: Knowledge, attitude and uptake among female medical and dental students in a tertiary institution in Benin-City, Nigeria. *Journal of Community Medicine and Primary Health Care*. 2015; 28(2): 101-108.
26. El-Ola MJA, Rajab MA, Abdallah DI, Fawaz IA, Awad LS, Tamin HM *et al*. Low rate of human papillomavirus vaccination among schoolgirls in Lebanon: barriers to vaccination with a focus on mothers' knowledge about available vaccines. *Therapeutics and Clinical Risk Management*. 2018; 14: 617-626.
27. Barnard M, George P, Perryman ML, Wolff LA. Human papillomavirus (HPV) vaccine knowledge, attitudes, and uptake in college students: Implications from the Precaution Adoption Process Model. *PLoS ONE*. 2017; 12(8): e0182266.
28. Igomu T, Folorunsho-Francis A. HPV vaccine, the cervical cancer prevention most Nigerian women don't know. *Punch Newspapers*. [cited March 9, 2024]. Available at: <https://healthwise.punchng.com/hpv-vaccine-the-cervical-cancer-prevention-most-nigerian-women-dont-know/>.
29. Rositch AF, Liu T, Chao C, Moran M, Beavis AL. Levels of parental human papillomavirus vaccine hesitancy and their reasons for not intending to vaccinate: insights from the 2019 national immunization survey-teen. *J Adolesc Health*. (2022). doi: 10.1016/j.jadohealth.2022.01.223.
30. MohdNazriShafei, NurhafizahZainon, Nor FazlinaZulkifli, Mohd Ismail Ibrahim. Knowledge and perception on human papilloma virus infection and vaccination among medical students of a university in Malaysia. *Procedia - Social and Behavioral Sciences* 116 (2014) 2707 – 2710

Table 1: Sociodemographic characteristics of respondents

Variables	Frequency <i>n</i> (%)
Age range (years)	
20 – 28	98 (36.7)
29 – 36	123 (46.1)
37 – 44	40 (15.0)
45 – 52	6 (2.3)
Age of child	
<6 months	185 (69.3)
6 to <12 months	306 (6.4)
≥12 months	65 (24.3)
Religion	
Christianity	263 (98.5)
Islam	1 (0.4)
No response	3 (1.1)
Education level	
Primary	1 (0.4)
Secondary	68 (25.4)
Tertiary	198 (74.2)
Social class	

I	8 (3.0)
II	54 (20.2)
III	126 (47.2)
IV	35 (13.1)
V	39 (14.6)

Table 2. Source of knowledge of HPV vaccine among respondents

Source of knowledge of vaccine	n	% (n =170)
Hospital	79	46.5%
Internet	38	22.4%
Radio/TV	16	9.4%
Family and friends	16	9.4%
School	16	9.4%
Billboard	3	1.8%
Social media	1	0.6%
Church	1	0.6%

Table 3: Association between intention for child to have HPV vaccination and sociodemographic characteristics of respondents

Variable	Wants HPV vaccine for child/ward, yes N, %	Odds Ratio (95% CI)	p-value
Age			
≤30 years	112 (77.2)	1	
>30 years	95 (79.2)	1.12 (0.62 – 2.01)	0.706
Religion			
Christianity	203 (77.8)	1	
Islam	1 (100.0)	4.62 x 10 ⁸ (0.0 – ∞)	1.000
Education of father			
University education or more	153 (79.7)	1	
Less than a university education	54 (74.0)	0.72 (0.39 – 1.36)	0.315
Education of mother			
University education or more	157 (80.1)	1	
Less than a university education	50 (72.5)	0.65 (0.35 – 1.23)	0.187

Occupation of father (in 2 categories)				
	Class 4/5	24 (70.6)	1	
	Class 1 – 3	183 (79.2)	1.59 (0.71 – 3.55)	0.256
Occupation of mother (in 2 categories)				
	Class 4/5	56 (76.7)	1	
	Class 1 – 3	147 (78.6)	1.12 (0.59 – 2.13)	0.740
State of domicile				
	Imo	197 (79.4)	1	
	Outside Imo	9 (69.2)	0.58 (0.17 – 1.97)	0.379

Table 4: Association between knowledge about HPV and baseline socio-demographic variables

Variable		Heard of HPV vaccine, yes N, %	Odds Ratio (95% CI)	p-value
Age	≤30 years	72 (49.7)	1	
	>30 years	60 (49.2)	0.98 (0.61 – 1.59)	0.938
Religion	Christianity	130 (49.4)	1	
	Islam	1 (100.0)	1.65 x 10 ⁹ (0.0 – ∞)	1.000
Education of father	University education or more	106 (54.9)	1	
	Less than a university education	26 (35.1)	0.45 (0.26 – 0.78)	0.004
Education of mother	University education or more	108 (54.5)	1	
	Less than a university education	24 (34.8)	0.44 (0.25 – 0.79)	0.005
Occupation of father (in 2 categories)				
	Class 4/5	15 (41.7)	1	
	Class 1 – 3	117 (50.6)	0.70 (0.34 – 1.42)	0.316
Occupation of mother (in 2 categories)				
	Class 4/5	32 (43.2)	1	
	Class 1 – 3	97 (51.6)	0.72 (0.42 – 1.23)	0.223
State of domicile				
	Imo	122 (48.8)	1	0.062
	Outside Imo	10 (76.9)	3.50 (0.94 – 13.01)	




Figure 1: Reasons respondents did not want HPV vaccination given to their children