

1
2
3
4
5
6
7
8
9

IMPACT OF SEED HARDENING AND FOLIAR APPLICATION OF GROWTH SUBSTANCES ON MORPHOLOGICAL PARAMETERS OF GROUNDNUT (*Arachis hypogaea* L.)

ABSTRACT

An experiment was carried out during summer and *kharif*, 2022. A Factorial Randomized Block Design with three replications was used for an experiment that included seed hardening as one factor with nine levels while foliar spray of Chlorocholine Chloride @500 mg/L as another factor with two levels. Days to initiation of flowering, days to 50% flowering, days to maturity were significantly minimum with GA₃-150 mg/L seed hardening treatment. Meanwhile, plant height, number of primary branches per plant were also found significantly higher in seed hardening with GA₃-150 mg/L while these morphological parameters were found significantly lower after application of foliar spray of CCC @500 mg/L as compared to control which helps to increase the yield of groundnut.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

Keywords: [seed hardening; foliar spray; CCC; plant height; number of primary branches per plant; number of leaves per plant]

1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.) is popularly known as peanut, a self-pollinated crop and allotetraploid. It is a member of the order fabales and family Fabaceae. It is regarded as It ranks 13th among the principal economic crops. It is also called as wonder nuts, earth nuts, monkey nuts, goobers, pindea, manilla nuts and poor men's cashew nut [1].

Groundnut is also known as "KING OF OILSEEDS CROPS" on account of its diversified uses for food, feed, fodder and industrial purpose. It is valued as a rich source of energy contributed by oil (48–50%) and protein (25–28%) in the kernels [2]. In addition, the groundnut kernels contain many health enhancing nutrients such as minerals such as K, Na, Ca, Mn, Fe, and Zn among others, antioxidants, and vitamins and also rich in mono-unsaturated fatty acids. They contain antioxidants like p-coumaric acid and resveratrol, Vitamin E, and many important B-complex groups of thiamine, pantothenic acid, vitamin B6, folates, and niacin. Groundnut is a dietary source of biologically active polyphenols, flavonoids, and isoflavones.

Groundnut cultivation in India spans all three primary agricultural seasons: *kharif*, *rabi* and *summer*, primarily under rainfed conditions. Among these seasons, *kharif* cultivation alone constitutes a substantial 75% share of the total groundnut production [3].

The low productivity of crops in rainfed areas is contributed by the use of poor-quality seeds. The features like rapid and identical seedling emergence are the two essential prerequisites to increase seed yield and seed quality in a number of field crops [4].

38 The hardening resulting from pre-sowing treatments is due to a number of
39 physio-chemical changes within the cytoplasm including greater hydration of colloids,
40 higher viscosity and elasticity of the protoplasm, increase in hydrophilic and decrease in
41 lipophilic colloids, increase in the temperature required for protein coagulation and
42 increase in bound water content [5]. Seed hardening accelerated rapid germination,
43 better root development and rapid growth of seedlings which enables absorption of more
44 moisture. It induces drought tolerance by increasing the resistance to protoplasmic
45 dehydration in young seedlings subjected to moisture stress. Flowering is also slightly
46 accelerated in hardened seeds [6].

47 Plants developed from hardened seeds often exhibit a faster growth due to an
48 improved nutrient use efficiency besides higher relative growth rate. It is a well-
49 established fact that, pre-soaking seeds with optimal concentration of phytohormones
50 enhance their germination, growth and yield of some crop species under condition of
51 environmental stress by increasing nutrient reserves through increased physiological
52 activities and root proliferation [7].

53 Chlorocholine Chloride is gibberellin biosynthesis inhibitor involved in the
54 inhibition of cyclization of geranyl-geranyl pyrophosphate to copyallyl pyrophosphate.
55 Growth regulators which inhibit the biosynthesis of gibberellins have been shown to
56 enable the plants to impart tolerance against abiotic stress due to water [8].

57

58 **2. MATERIAL AND METHODS**

59

60 **2.1 Experimental Site**

61 The experiment is conducted at Regional Research Station, Anand Agricultural
62 University, Anand, India during Summer and *kharif*, 2022.

63 **2.2 Treatment Details**

64 Eighteen treatment combinations involving nine levels of seed hardening
65 treatments and two levels of foliar spray were incorporated in the study. Thus, eighteen
66 treatment combinations with two factors were embedded as factorial randomized block
67 design with three replications. Details of the treatments with their symbols are given as
68 under.

69 Factor-1: Seed Hardening (A)

70 A1 : CaCl₂1%

71 A2 : Ethrel-50 mg/L

72 A3 : Ethrel-100 mg/L

73 A4 : Ethrel-150 mg/L

74 A5 : GA₃-50 mg/L

75 A6 : GA₃-100 mg/L

76 A7 : GA₃-150 mg/L

77 A8 : Soaking in water

78 A9 : Control

79 Factor-2: Foliar Spray (B)

80 B1 : Control no foliar spray

81 B2 : CCC @500 mg/L

82 *Foliar spray of CCC was given at 35 and 55 DAS in all treatments

83 There were eighteen treatment combinations were evaluated in the present study viz.,
84 A1B1:CaCl₂1% seed hardening+Control (No foliar spray), A2B1:Ethrel-50 mg/L seed
85 hardening+Control (No foliar spray), A3B1:Ethrel-100 mg/L seed hardening+Control (No
86 foliar spray), A4B1:Ethrel-150 mg/L seed hardening+Control (No foliar spray), A5B1:
87 GA₃-50 mg/L seed hardening+Control (No foliar spray), A6B1:GA₃-100 mg/L seed
88 hardening+Control (No foliar spray), A7B1:GA₃-150 mg/L seed hardening+Control (No
89 foliar spray), A8B1:Water soking as seed hardening+Control (No foliar spray),
90 A9B1:Control (Without hardening)+Control (No foliar spray), A1B2:CaCl₂1% seed
91 hardening+CCC 500 mg/L foliar spray, A2B2:Ethrel-50 mg/L seed hardening+CCC 500
92 mg/L foliar spray, A3B2:Ethrel-100 mg/L seed hardening+CCC 500 mg/L foliar spray,
93 A4B2: Ethrel-150 mg/L seed hardening+CCC 500 mg/L foliar spray, A5B2: GA₃-50 mg/L
94 seed hardening+CCC 500 mg/L foliar spray, A6B2:GA₃-100 mg/L seed hardening+CCC
95 500 mg/L foliar spray, A7B2: GA₃-150 mg/L seed hardening+CCC 500 mg/L foliar spray.

96 **2.3 Methods of Seed Hardening and Foliar Application of Growth** 97 **Substances**

98 CaCl₂ 1% was prepared by dissolving 10 g of CaCl₂ in 1 liter of distilled water. Ethrel-
99 50 mg/L, Ethrel-100 mg/L and Ethrel-150 mg/L were prepared by dissolving 50, 100 and
100 150 mg of Ethrel in one liter of water respectively. GA₃-50 mg/L, GA₃-100 mg/L and GA₃-
101 150 mg/L were prepared by dissolving 50, 100 and 150 mg of GA₃ in one liter of water
102 respectively.

103 Seed hardening treatments were applied to Groundnut seeds, soaking them in double
104 volume solutions for four hours to prevent germination. After drying, seeds were ready
105 for sowing in the field and under laboratory conditions, ensuring their original moisture
106 level.

107 This experiment uses Chlorocholine Chloride (CCC) as a foliar spray. A stock solution
108 of 50 % CCC was prepared, and a final solution of 10 liters was prepared. Spraying was
109 carried out at 35 and 55 DAS in respective plots during both seasons.

110 111 **2.4 Morphological Parameters**

112 **2.4.1 Days to initiation of flowering (days)**

113 The number of days taken from sowing to the opening of first flower in an
114 experimental plot and was expressed in days.

115 116 **2.4.2 Days to 50% flowering (days)**

117 The number of days required for 50% flowering was recorded from the date of
118 sowing in all treatments in all three replications.

119 120 **2.4.3 Days to maturity (days)**

121 The number of days required from sowing to the date of 50% of plants became
122 dry were recorded as the date of physiological maturity in all treatments in three
123 replications.

124 125 **2.4.4 Plant height (cm)**

126 Plant height was recorded by non-destructive method. Plant height of groundnut
127 crop was measured from five plants selected randomly in each treatment for recording
128 observations of average plant height at 30, 50, 70, 90 DAS and at harvest from the base
129 of the plant (ground level) to the tip of upper most fully opened leaf and finally the mean

130 height of plant in centimeter (cm) at each period in each treatment was worked out and
131 were recorded for statistical analysis. These same five plants were also used for other
132 observations.

133

134 **2.4.5 Number of primary branches/plant**

135 Number of branches per plant was counted from previously selected five plants
136 from each net plot at 30, 50, 70, 90 DAS and at harvest of the crop. The mean number of
137 primary branches per plant was worked out and recorded separately for each treatment.

138

139 **2.4.6 Number of leaves/plant**

140 The number of green leaves from top to bottom of the plants was counted in the
141 randomly tagged five plants at 30, 50, 70, 90 DAS and at harvest. The average was
142 worked out and expressed as number of green leaves per plant.

143

144 **3. RESULTS AND DISCUSSION**

145

146 **3.1 Effect of Seed Hardening on Morphological Parameters**

147 **3.1.1 Effect of seed hardening on days to initiation of flowering (Days)**

148 The data summarized in Table 1 showed that the days to initiation of flowering
149 was influenced significantly by the use of different seed hardening treatments in
150 groundnut crop. The data regarding days to initiation of flowering (28.28, 27.53 and
151 27.91) showed that significantly in seed hardening treatment with GA₃-150 mg/L seed
152 hardening (A7) during summer and *kharif*-2022 and in pooled results, respectively.

153

154 **3.1.2 Effect of seed hardening on days to 50% flowering (Days)**

155 The data (Table 1) regarding days to 50% flowering (32.07, 32.60 and 32.33)
156 showed that significantly lower with GA₃-150 mg/L seed hardening (A7) during the
157 summer and *kharif*, 2022 and in pooled results, respectively. Whereas, higher days to
158 50% flowering (37.14, 37.11 and 37.12) was recorded with control (A9) in in both
159 seasons and in pooled results, respectively.

160

161 **3.1.3 Effect of seed hardening on days to maturity (Days)**

162 The significantly (Table 1) lower days to maturity (120.13, 120.72 and 120.43)
163 was recorded GA₃-150 mg/L seed hardening (A7) which was statistically at par with GA₃-
164 100 mg/L seed hardening in the summer and *kharif*, 2022 and pooled data, respectively.

165

165 **3.1.4 Effect of seed hardening on plant height (cm)**

166 The plant height (Table 2) at 30, 50, 70, 90 DAS and at harvest affects
167 significantly due to seed hardening treatment during summer and *kharif* 2022, as well as
168 in the pooled results. Among the treatments, GA₃-150 mg/L seed hardening (A7) showed
169 higher plant height (11.36, 14.44 and 12.90 cm) at 30 DAS, while Ethrel-50 mg/L seed
170 hardening (A2) showed that significantly higher plant height of 20.91 cm which was
171 statistically at par with GA₃-150 mg/L seed hardening, A7 (20.36 cm) during summer-
172 2022 while CaCl₂ 1% seed hardening (A1) showed higher plant height 22.44 cm which
173 was statistically at par with A7 (21.90 cm) during *kharif* season. Also plant height was
174 recorded

175
176

Table 1 Effect of seed hardening and foliar spray on days to initiation of flowering, days to 50% flowering and days to maturity in groundnut during summer and *kharif*, 2022 as well as in pooled analysis

	Days to Initiation of Flowering			Days to 50% Flowering			Days to Maturity		
	Summer	<i>kharif</i>	Pooled	Summer	<i>kharif</i>	Pooled	Summer	<i>kharif</i>	Pooled
Seed Hardening (A)									
A1	30.30	30.13	30.22	36.14	36.26	36.20	122.36	122.51	122.43
A2	30.05	30.00	30.03	35.81	35.21	35.51	121.95	122.12	122.04
A3	29.95	29.42	29.68	35.44	35.13	35.29	122.57	122.72	122.64
A4	28.93	28.53	28.73	34.05	34.07	34.06	121.12	121.59	121.36
A5	29.50	28.82	29.16	34.93	34.81	34.87	121.04	121.37	121.20
A6	28.45	28.35	28.40	33.35	33.50	33.42	120.78	120.96	120.87
A7	28.28	27.53	27.91	32.07	32.60	32.33	120.13	120.72	120.43
A8	30.65	30.62	30.63	36.51	36.24	36.37	122.99	123.30	123.14
A9	31.84	31.47	31.66	37.14	37.11	37.12	123.02	123.39	123.20
S.Em.(±)	0.278	0.246	0.185	0.731	0.887	0.575	0.945	0.438	0.521
C.D. (0.05)	0.798	0.707	0.523	2.101	2.549	1.622	NS	NS	1.469
Foliar Spray (B)									
B1	29.77	29.44	29.61	35.09	34.98	35.04	121.98	122.21	122.09
B2	29.77	29.42	29.60	35.00	35.00	35.00	121.57	121.94	121.76
S.Em.(±)	0.131	0.116	0.087	0.345	0.418	0.271	0.445	0.207	0.245
C.D. (0.05)	NS	NS	NS	NS	NS	NS	NS	NS	NS
Interaction									
AxS	-	-	NS	-	-	NS	-	-	NS
BxS	-	-	NS	-	-	NS	-	-	NS
AxB	NS	NS	NS	NS	NS	NS	NS	NS	NS
AxBxS	-	-	NS	-	-	NS	-	-	NS
C.V.(%)	2.28	2.05	2.17	5.11	6.21	5.68	1.90	0.88	1.48

177 *NS=Non-Significant, DAS- Days After Sowing

178 **Table 2** Effect of seed hardening and foliar spray on plant height in groundnut during summer and *kharif*, 2022 as well as in pooled

Plant Height (cm)				
30 DAS	50 DAS	70 DAS	90 DAS	At Harvest

	Summer	kharif	Pooled	Summer	kharif	Pooled	Summer	kharif	Pooled	Summer	kharif	Pooled	Summer	kharif	Pooled
Seed Hardening (A)															
A1	8.77	12.76	10.77	19.51	22.44	20.98	28.45	30.98	29.71	29.13	32.23	30.68	35.01	36.73	35.87
A2	8.06	8.47	8.27	20.91	18.46	19.69	29.57	29.27	29.42	30.65	30.83	30.74	34.67	34.58	34.63
A3	8.19	30 DAS	6.85	18.58	50 DAS	19.51	32.11	70 DAS	30.76	35.80	90 DAS	33.88	35.21	At Harvest	35.92
A4	Summer	kharif	Pooled	Summer	kharif	Pooled	Summer	kharif	Pooled	Summer	kharif	Pooled	Summer	kharif	Pooled
Seed Hardening (A)	13.86	11.67	10.74	19.02	21.20	20.11	25.87	30.74	28.30	31.93	36.45	34.19	35.36	33.68	34.52
A1A6	3.35	1.16	1.74	4.82	1.66	2.76	7.38	3.52	3.52	6.20	3.56	5.69	6.10	4.25	4.88
A2A7	11.36	15.44	12.90	20.36	31.00	31.73	31.15	35.33	32.34	37.22	37.26	37.25	38.43	49.64	44.03
A8	2.85	2.83	2.84	4.39	3.10	3.75	4.42	4.20	4.31	5.18	4.71	4.95	6.33	8.10	6.31
A9	7.76	10.92	9.34	19.78	19.87	19.83	28.73	31.12	29.92	32.59	32.44	32.51	34.87	32.81	42.25
A9	7.52	5.10	6.31	18.58	19.16	18.87	26.82	30.17	28.49	30.10	27.08	28.59	34.54	39.39	36.96
S.Em.(±)	0.210	0.229	0.155	0.381	0.355	0.260	1.084	0.744	0.657	1.066	1.081	0.759	1.038	1.122	0.764
C.D.(0.05)	0.603	0.657	0.438	1.096	1.019	0.735	3.115	2.138	1.855	3.063	3.106	2.142	2.983	3.226	2.157
Foliar Spray (B)															
B1	8.73	10.79	9.76	20.45	22.03	21.24	30.80	32.15	31.48	33.24	36.48	34.86	38.70	37.84	38.27
B2	8.75	10.70	9.73	18.06	18.65	18.36	27.03	29.00	28.02	29.79	31.56	30.67	36.17	32.70	34.44
S.Em.(±)	0.099	0.108	0.073	0.180	0.167	0.123	0.511	0.351	0.310	0.502	0.509	0.358	0.489	0.529	0.360
C.D.(0.05)	NS	NS	NS	0.517	0.480	0.346	1.468	1.008	0.874	1.444	1.464	1.010	1.406	1.521	1.017
Interaction															
AxS	-	-	Sig.	-	-	Sig.	-	-	Sig.	-	-	Sig.	-	-	Sig.
BxS	-	-	NS	-	-	Sig.	-	-	Sig.	-	-	Sig.	-	-	Sig.
AxB	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
AxBxS	-	-	NS	-	-	NS	-	-	NS	-	-	NS	-	-	NS
C.V.(%)	5.89	5.21	5.52	4.85	4.27	4.55	9.34	5.96	7.72	8.06	7.99	8.02	7.12	7.43	7.28

180 *Sig.-Significant

A3	3.23	2.77	3.00	4.41	3.03	3.72	4.62	3.67	4.15	5.36	5.20	5.28	6.87	6.33	6.60	
A4	3.00	2.77	2.88	4.34	3.13	3.73	6.09	3.25	4.67	7.38	6.91	7.14	7.84	6.36	7.10	
A5	3.30	2.90	3.10	4.37	2.89	3.63	4.59	3.25	3.92	4.90	4.35	4.62	5.97	5.53	5.75	
A6	3.32	2.63	2.97	4.42	3.18	3.80	4.50	3.89	4.20	6.09	4.95	5.52	7.19	6.97	7.08	
A7	3.70	3.03	3.39	4.57	3.23	3.90	6.32	4.80	5.56	8.08	6.48	7.28	8.85	6.67	7.36	
A8	2.90	3.00	2.95	4.39	2.86	3.63	4.85	4.03	3.94	5.25	4.15	4.70	6.36	5.24	5.80	
A9	2.90	2.83	2.87	3.19	3.12	3.15	3.41	4.04	3.72	3.94	4.27	4.11	5.14	4.60	4.87	
S.Em.(±)	0.069	0.089	0.056	0.066	0.059	0.044	0.067	0.125	0.071	0.174	0.148	0.114	0.234	0.191	0.151	
C.D._(0.05)	0.199	0.255	0.159	0.189	0.169	0.177	0.193	0.361	0.201	0.501	0.425	0.323	0.673	0.548	0.426	
Foliar Spray (B)																
B1	3.17	2.97	2.97	4.62	3.16	3.89	4.70	3.92	4.31	6.04	5.69	5.86	7.51	6.37	6.74	
B2	3.17	2.91	3.04	3.95	3.01	3.48	4.40	3.78	4.09	5.60	4.62	5.11	6.32	5.56	6.14	
Mean	3.17	2.84	3.01	4.28	3.08	3.68	4.55	3.85	4.20	5.82	5.15	5.49	6.91	5.97	6.44	
S.Em.(±)	0.033	0.042	0.027	0.031	0.028	0.021	0.032	0.059	0.034	0.082	0.070	0.054	0.110	0.090	0.071	
C.D._(0.05)	NS	NS	NS	0.089	0.080	0.059	0.091	0.170	0.095	0.236	0.201	0.152	0.317	0.259	0.201	
Interaction																
AxS	-	-	Sig.	-	-	Sig.	-	-	Sig.	-	-	Sig.	-	-	Sig.	
BxS	-	-	NS	-	-	Sig.	-	-	Sig.	-	-	Sig.	-	-	Sig.	
AxB	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	
AxBxS	-	-	NS	-	-	NS	-	-	NS	-	-	NS	-	-	NS	
C.V.(%)	5.35	7.66	6.49	3.68	4.68	4.10	3.61	7.98	5.87	7.33	7.04	7.22	8.30	7.84	8.13	

181 **Table 3 Effect of seed hardening and foliar spray on number of primary branches/plant in groundnut during summer and *kharif*,**
182 **2022 as well as in pooled analysis**

183 higher with A1 (21.13 cm) in pooled result during 50 DAS. The significant higher plant
184 height (31.15 cm) was recorded in seed hardening with GA₃-150 mg/L (A7) during
185 summer, while 35.75 cm was noted in GA₃-100 mg/L (A6) during *kharif* seasons which
186 was statistically at par with GA₃-150 mg/L seed hardening (35.33 cm). A7, GA₃-150 mg/L
187 seed hardening recorded higher plant height (33.24 cm) during pooled analysis at 70
188 DAS. Significantly higher plant height at 90 DAS (37.22, 38.37 and 37.24 cm) was
189 recorded with GA₃-150 mg/L (A7), ethrel-150 mg/L (A4) and GA₃-150 mg/L (A7) in both
190 the seasons 2022, and pooled data, respectively which was at par with GA₃-150 mg/L
191 seed hardening (37.26 cm) during *kharif* season, 2022. Data (Table 3.2) clearly indicated
192 that the significantly higher plant height was observed in GA₃-150 mg/L seed hardening
193 (A7) (38.43, 49.64 and 44.03 cm) during summer and *kharif*-2022 and pooled analysis
194 during at harvest, respectively.

195 The beneficial effect on growth regulators at particular concentration can be
196 attributed to the cell elongation and quick cell multiplication. This appears the most
197 probable reason of increase in the plant height in GA₃ treatments in present investigation
198 as well. The present findings are in close agreement with reported by Hasan and Ismail
199 [9], Agwane and Parhe [10] in soyabean, Keykhaet *al.* [11] in Mungbam, Narayanreddy
200 and Biradapatil [12] in sunflower.

201 **3.1.4 Effect of seed hardening on number of primary branches per plant**

202 The number of primary branches/plant at 30 DAS (Table 3) showed statistically
203 significant due to seed hardening treatments. Highest number of primary branches/plant
204 were observed in GA₃-150 mg/L seed hardening (3.70, 3.03, and 3.39) at 30 DAS, (4.57,
205 3.23 and 3.90) at 50 DAS, (6.32, 4.80, 5.56) at 70 DAS during the summer and *kharif*
206 2022, as well as in the pooled results respectively. Maximum number of primary
207 branches/plant was observed in GA₃-150 mg/L seed hardening, A7 (8.08) during
208 summer-2022 while ethrel-150 mg/L, A4 (6.91) showed highest number of primary
209 branches/plant in *kharif*-2022 which was at par with GA₃-150 mg/L seed hardening (A7,
210 6.48). In pooled result, A7 (7.28) presented higher number of primary branches/plant for
211 90 DAS. Meanwhile, higher number of primary branches/plant was observed in GA₃-150
212 mg/L (8.85), GA₃-100 mg/L (6.97) and GA₃-150 mg/L (7.36) seed hardening during the
213 summer, *kharif* -2022, as well as in the pooled basis which was statistically at par with
214 GA₃-150 mg/L (A7) ranged 6.67 during *kharif*, 2022.

215 The improvement in plant growth by increasing number of primary
216 branches/plant because of the use of GA₃ might be attributed to cell elongation and cell
217 division. GA₃ influences the action of various enzymes, particularly amylase and
218 enhances the movement of starch particles in the cotyledons, consequently triggering
219 growth. The result was in accordance with Hasan and Ismail [9] in groundnut.

220 **3.1.5 Effect of seed hardening on number of leaves per plant**

221 The number of leaves/plant at 30, 50, 70, 90 DAS and at harvest (Table 4) was
222 significantly affected by seed hardening treatments. The higher number of leaves/plant
223 were found with seed hardening with GA₃-150 mg/L (A7, 46.97 and 66.42) during
224 summer, 2022 and pooled result while seed hardening with GA₃-100 mg/L (A6, 86.03)
225 showed maximum number of leaves/plant which was at par with A4 (82.82), A5 (85.30)
226 and A7 (85.87) during *kharif*-2022 at 30 DAS. Number of leaves/plant at 50 DAS
227 (126.86, 176.76 and 150.65) showed that significantly higher with seed hardening with

228 ethrel-50 mg/L (A2), GA₃-150 mg/L (A7) and ethrel-50 mg/L (A2) during summer, *kharif*
229 and in pooled results,

230 Table 4 Effect of seed hardening and foliar spray on number of primary branches/Plant in groundnut during summer and *kharif*, 2022 as
231 well as in pooled analysis

	Number of Leaves/Plant														
	30 DAS			50 DAS			70 DAS			90 DAS			At Harvest		
	Summer	<i>kharif</i>	Pooled	Summer	<i>kharif</i>	Pooled	Summer	<i>kharif</i>	Pooled	Summer	<i>kharif</i>	Pooled	Summer	<i>kharif</i>	Pooled
Seed Hardening (A)															
A1	30.80	74.90	52.85	121.23	149.24	135.23	475.85	452.29	464.07	485.14	216.58	350.86	473.07	213.42	343.25
A2	31.91	76.57	54.24	126.86	174.43	150.65	394.57	418.07	406.32	499.02	207.08	353.05	503.88	222.20	363.04
A3	43.53	79.23	61.38	122.46	174.02	148.24	392.07	417.87	404.97	516.87	223.68	370.27	511.97	222.80	367.39
A4	44.41	82.82	63.61	121.02	174.57	147.80	462.67	408.13	435.40	485.59	282.72	384.16	583.57	261.54	422.56
A5	43.67	85.30	64.48	120.87	174.22	147.55	460.09	414.85	437.47	534.98	263.99	399.49	565.56	256.36	410.96
A6	46.23	86.03	66.13	116.92	173.95	145.43	395.22	443.97	419.59	598.28	206.19	402.23	562.82	184.03	373.42
A7	46.97	85.87	66.42	123.04	176.76	149.90	488.07	497.27	492.67	542.72	310.37	426.55	667.41	264.10	465.76
A8	43.74	78.57	61.16	118.99	171.34	145.17	462.67	422.98	442.82	531.99	257.38	394.68	525.38	269.43	397.41
A9	31.64	74.48	53.06	115.35	160.04	137.69	390.30	438.80	414.55	477.74	230.18	353.96	460.33	181.27	320.80
S.Em.(±)	1.074	1.826	1.059	2.107	4.168	2.335	18.025	11.840	10.783	20.760	4.652	10.637	22.105	4.270	11.257
C.D._(0.05)	3.088	5.249	2.990	6.055	11.980	6.590	51.805	34.030	30.430	59.664	13.371	30.019	63.530	12.272	31.767
Foliar application (B)															
B1	40.99	80.47	60.73	139.03	202.38	170.71	512.79	475.89	494.34	608.59	316.24	462.41	650.19	291.84	471.01
B2	39.65	80.37	60.01	102.47	137.30	119.88	358.66	393.94	376.30	429.71	172.24	300.97	428.48	169.31	298.89
S.Em.(±)	0.506	0.861	0.499	0.993	1.965	1.101	8.497	5.582	5.083	9.786	2.193	5.015	10.420	2.013	5.306
C.D._(0.05)	NS	NS	NS	2.854	5.647	3.107	24.421	16.042	14.345	28.126	6.303	14.151	29.948	5.785	14.975
Interaction															
AxS	-	-	Sig.	-	-	Sig.	-	-	Sig.	-	-	Sig.	-	-	Sig.
BxS	-	-	NS	-	-	Sig.	-	-	Sig.	-	-	Sig.	-	-	Sig.
AxB	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
AxBxS	-	-	NS	-	-	NS	-	-	NS	-	-	NS	-	-	NS
C.V.(%)	6.53	5.56	6.08	4.27	6.01	5.57	10.13	6.67	8.58	9.80	4.67	9.80	10.04	4.54	10.13

232 respectively. Maximum number of leaves were found with seed hardening with GA₃-150
233 mg/L (488.07497.27, 492.66) during summer, *kharif*-2022 and pooled analysis. An
234 examination of data given in Table 3.4 indicated that the significantly higher number of
235 leaves/plant at 90 DAS was recorded with seed hardening with GA₃ 100 mg/L (A6,
236 598.28) which was statistically at par with A7 (542.72) in summer, 2022. Whereas, seed
237 hardening with GA₃-150 mg/L (310.37, 426.54) recorded higher leaves/plant during
238 *kharif*, 2022 and pooled result. According to Data (Table 4.4) higher number of
239 leaves/plant were significantly found in A7 (667.41, 465.76) during summer, 2022 and
240 pooled analysis, whereas seed hardening with water soaked treatment, A8 (269.43)
241 during *kharif*, 2022 at harvest.

242 Improvement of vegetative growth represented by enhancement of number of
243 leaves/plant in groundnut showed the positive effect with seed hardening treatment.
244 These result may be attributed to healthy germination of seeds, which in turn gave the
245 plant better start and induced further growth of groundnut seedling. The number of
246 leaves per plant, an additional parameter of growth is greatly influenced by growth
247 regulators. The effect of growth regulators on number of leaves was observed, at peak
248 stage of plant. It is well established fact that GA₃ acts in cell elongation or enlargement
249 resulting in increased number of functional leaves. Elongation of cell may also have
250 resulted in larger blade size of the leaves.

251 **3.2 Effect of Foliar Spraying of CCC on Morphological Parameters**

252

253 **3.2.1 Effect of foliar spray of CCC on days to initiation of flowering, days to 50% 254 flowering and days to maturity**

255 As per result (Table1) the effect of foliar spray of CCC @500 mg/L on days to
256 initiation of flowering, days to 50% flowering and days to maturity during summer and
257 *kharif*, 2022 and in pooled results was found to be non-significant.

258

259 **3.2.2 Effect of foliar spray of CCC on plant height (cm)**

260 First foliar spray of CCC was done at 35 DAS, so there is no significant result
261 found for foliar spraying of CCC at 30 DAS.

262 According to Table 2 plant height showed significantly lower after foliar
263 application of CCC @500 mg/L at 50 DAS (38.43, 49.64 and 44.03 cm), 70 DAS (27.03,
264 29.00 and 28.02 cm), 90 DAS (29.79, 31.56 and 30.67 cm) and at harvest (29.79, 31.56
265 and 30.67 cm) as compared to control (B1) during summer and *kharif*, 2022 and in
266 pooled analysis, respectively.

267 Chlorocholine chloride (CCC) reduces plant height by inhibiting gibberellin
268 biosynthesis, a plant growth hormone, responsible for the stem elongation. As gibberellin
269 promote cell division and elongation, their inhibition stunted growth and reduced plant
270 height. The findings are in conformity with result of Singh and Jambukiya [13] in green
271 gram, Bhadaneet *et al.* [14] in green gram and Prajapati *et al.* [15] in blackgram.

272 **3.2.3 Effect of foliar spray of CCC on number of primary branches per plant**

273 Number of primary branches/plant was significantly minimum after foliar
274 application of CCC @500 mg/L, B2 (3.95, 3.01, 3.48) at 50 DAS, (4.40, 3.78, 4.09) at 70
275 DAS, (5.60, 4.62, 5.11) at 90 DAS and (6.32, 5.56, 6.14) at harvest as compared to
276 control (B1) during summer and *kharif*, 2022 and pooled analysis (Table 3).

277 As, chlorocholine chloride (CCC) plant growth retardants, particularly onium
278 compounds are able to increase the partitioning of assimilates to roots and thereby

279 improve yield through the inhibition of gibberellin biosynthesis or action. According to
280 result, number of primary branches per plant decrease after foliar application of CCC
281 indicated indirectly growth of below ground part.

282 **3.2.4 Effect of foliar spray of CCC on number of leaves per plant**

283 Number of leaves/plant at 50, 70, 90 DAS and at harvest (Table 4) was recorded
284 significantly decrease with foliar spraying of CCC @500 mg/L (B2) during summer, *khari*
285 as well as in pooled results, respectively. The higher number of leaves/plant at 50 DAS
286 were recorded with control (B1). Lower number of leaves/plant was observed (102.47,
287 137.30 and 119.85) at 50 DAS, (358.66, 393.94 and 376.30) 70 DAS, (429.71, 172.24,
288 300.97) 90 DAS and (428.48, 169.31, 298.89) at harvest was recorded with foliar
289 spraying of CCC @500 mg/L (B2) during both the seasons and in pooled analysis,
290 respectively.

291 The result might be due to the response of CCC was antagonistic to GA₃. Similar
292 result was found with Mohammed [16] in *Chrysanthemum* and El- Kheir *et al.* [17] in
293 groundnut.

294 **3.3 Interaction Effect of Seed Hardening And Foliar Spray of CCC on** 295 **Morphological Parameters**

296 The interaction effect between different seed hardening treatments and foliar
297 spraying of CCC @500 mg/L for all morphological parameters like days to initiation of
298 flowering, days to 50% flowering, days to maturity, plant height, number of primary
299 branches per plant and number of leaves per plant were found non-significant at 30, 50,
300 70, 90 DAS and at harvest.

301

302 **4. CONCLUSION**

303

304 The study's findings indicated that treating groundnuts with GA₃-150 mg/L seed
305 hardening treatment and applying CCC as foliar application at a dose of 500 mg/L were
306 effective in positive manner of morphological characteristics such as the days to initiation
307 of flowering, days to 50% flowering, days to maturity, plant height, number of primary
308 branches per plant and number of leaves per plant. To put it succinctly, growers aiming
309 to get a higher morphological produce were advised to use GA₃ at 150 mg/L and
310 Chlorocholine Chloride (CCC) at 500 mg/L by foliar spraying.

311

312 **AUTHORS' CONTRIBUTIONS**

313

314 All writers worked together to complete this work. This study is a component of
315 author KUP's doctoral thesis research project. The first three authors, KUP, SJM, and
316 ASB, collaborated on the work, while the fourth author, JJG, helped with the research
317 analysis and paper writing. KUP collected and analyzed data periodically and according
318 to stages. Analysis of the data and Script writing was done by the author KUP. The
319 experiment's design and oversight were completed by author SJM. Author ASB provided
320 guidance during the field test procedures. The final manuscript was reviewed and
321 approved by all writers.

322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369

REFERENCES

1. Gadhiya HA, Borad PK, Bhut JB. Effectiveness of synthetic insecticides against *Helicoverpa armigera* (Hubner) Hardwick and *Spodopteralitura* (Fabricius) infesting groundnut. *The Bioscan*. 2014 Sep 10;9(1):23-6.
2. Jambunathan R. Uses of tropical grain legumes. 1991.
3. Barman M, Gunri SK, Puste AM, Paul S. Effect of paclobutrazol on growth and yield of kharif groundnut (*Arachis hypogaea* L.). *International Journal of Agriculture, Environment and Biotechnology*. 2017;10(4):513-8.
4. Krishnotar B, Srivastava AK, Shahi JP. Response of rabi maize crop to seed invigoration with magnesium nitrate and distilled water. *seeds*. 2009; 900:119.
5. Solaimalai A, Subburamu K. Seed hardening for field crops—A review. *Agricultural Reviews*. 2004;25(2):129-40.
6. Jain LK, Verma MP, Ram N, Choudhary A, Parewa HP. Seed hardening: A way to tolerate against abiotic stress in rainfed areas. *International Journal of Economic Plants*. 2022 Feb 21;9(Feb, 1):018-21.
7. Bozcuk S. Effects of kinetin and salinity on germination of tomato, barley and cotton seeds. *Annals of Botany*. 1981 Jul 1;48(1):81-4.
8. Lone NA, Khan NA, Bhat MA, Mir MR, Razvi SM, Baht KA, Rather GH, Nawsheeba W, Sabina A, Bukhari SA. Effect of chlorocholine chloride (CCC) on plant growth and development. *Int. J. Curr. Res*. 2010 May 17;6(2):1-7.
9. Hasan M, Ismail BS. Effect of gibberellic acid on the growth and yield of groundnut (*Arachis hypogaea* L.). *SainsMalaysiana*. 2018 Feb 1;47(2):221-5.
10. Agawane RB, Parhe SD. Effect of seed priming on crop growth and seed yield of soybean [*Glycine max* (L.) Merrill]. *The Bioscan*. 2015 Mar 26;10(1):265-70.
11. Keykha, Mojtaba, Hamid Reza Ganjali, and Hamid Reza Mobasser. Effect of salicylic acid and gibberellic acid on some characteristics in mungbean (*Vignaradiata*). 2014; 70-75.
12. Narayanareddy AB, Biradarpatil NK. Effect of pre-sowing invigouration seed treatments on seed quality and crop establishment in sunflower hybrid KBSH-1. 2012;43-46.
13. Singh, C., and H. Jambukiya. Effect of foliar application of plant growth regulators on growth and yield attributing characters of green gram (*Vigna radiata* L. Wilczek). 2020: 258-264.
14. Bhadane RS, Prajapati KR, Patel DB. Effect of seed hardening on morpho-physiological characters in Mung bean (*Vigna radiata* L.). *International Journal of Chemical Studies*. 2019;7(4):1760-3.
15. Prajapati KR, Patel DB, Patil K, Bhadane RS. Effect of seed hardening on morpho-physiological and yield parameters in black gram (*Vigna mungo* L.). *International Journal of Chemical Studies*. 2017;5(4):439-41.
16. Mohammed AR. Effects of Foliar Application of Plant Growth Regulators on Growth and Flowering Characteristics of *Chrysanthemum* CV. Paintball. *Pakistan Journal of Life & Social Sciences*. 2017 May 1;15(2).
17. El-Kheir MS, Kandil SA, Mekki BB. Physiological response of two soybean cultivars grown under water stress conditions as affected by CCC treatment. 1991; 179-200.

