

Antimicrobial Effects of *Aloe vera*, *Psidium guajava* and *Hibiscus sabdariffa* Extracts on Selected Pathogens

Abstract

Plant materials are repositories of bioactive compounds of medicinal values. Traditional medicines rely on the therapeutic potential of herbal plants. This study aimed to assess the antimicrobial effects of *Aloe vera*, *Psidium guajava* and *Hibiscus sabdariffa* extracts on *Salmonella typhi*, *Staphylococcus aureus* and *Escherichia coli* and *Candida albicans*. Plant extracts were obtained by Soxhlet extraction, using ethanol as solvent. Extracts were qualitatively screened for the presence of alkaloid, saponins, flavonoid and tannins. Saponins and flavonoids were detected in all plant extracts; alkaloid in *P. guajava* and *H. sabdariffa* extracts, and tannins in *A. vera* and *P. guajava* extracts. The concentrations of the residual plant extracts were 2.0 mg/ml for *A. vera*, 2.1 mg/ml for *P. Guajava* and 1.83 mg/ml for *H. sabdariffa*. *Staphylococcus aureus* was only susceptible to *A. vera*. *Salmonella typhi* was not sensitive to all extracts. *Escherichia coli* was sensitive to all extracts in the order *H. sabdariffa* > *P. guajava* > *A. vera*. *Candida albicans* was sensitive to *A. vera* and *P. guajava* extracts. The MIC of *A. vera* was 2.0 mg/ml for *S. aureus*, *E. coli* and *C. albicans* respectively. The MIC of *P. guajava* was 0.021 mg/ml and 2.1 mg/ml for *E. coli* and *C. albicans* respectively. The MIC of *H. sabdariffa* was 1.83 mg/ml for *E. coli*. The results show that *A. vera*, *H. sabdariffa* and *P. guajava* extracts contain phytochemicals with inhibitory potencies against *S. aureus*, *E. coli* and *C. albicans*. These plant materials may prove useful as natural remedies for managing infections caused by susceptible microorganisms.

Keywords: *Aloe vera*, Antimicrobial, *Hibiscus sabdariffa*, Pathogens, *Psidium guajava*

Introduction

Plant-based materials offer several potential benefits that can complement the use of synthetic drugs in the management of infectious diseases [1-3]. This has become very important in the face of the growing challenge of antibiotic resistance. Plants have evolved a vast array of secondary metabolites, many of which exhibit antimicrobial properties against a wide range of pathogens, including bacteria, fungi, and viruses. These natural compounds commonly referred to as phytochemicals include compounds such as glycosides, tannins, saponins, terpenoids, alkaloids, phenolics, terpenes, anthraquinones, salicylic acid, lupeol and flavonoids, which have been severally reported to interfere with several cellular mechanisms in microorganisms [4-6].

The clamour for the use of natural compounds with antimicrobial properties has been because of the issue of antimicrobial resistance [4]. It has been inferred that the complex and often synergistic interactions between the various bioactive compounds found in plant extracts make it more challenging for pathogens to develop resistance, as they would need to overcome multiple defence mechanisms simultaneously [7-9]. Again, plant materials can be used at lower doses of individual compounds, which is a useful consideration in drug safety [10].

Aloe vera is a succulent plant that has been used for a long time in traditional medicine due to its various therapeutic properties, believed to be suitable for hastening wound healing, treating anti-inflammatory, arthritis, diabetes sunburns, eczema, inflammatory bowel diseases and psoriasis ulcers, diverse other therapeutic applications [11-13]. *Psidium guajava*, commonly known as guava, is a tropical fruit tree that has been used in traditional medicine for various health

benefits. This versatile medicinal plant is rich in diverse bioactive compounds accountable for their pharmacological activities in the treatment of diarrhea, wounds, arthritis, cardiovascular diseases, diabetes and neurodegenerative disorders [14,15]. *Hibiscus sabdariffa*, also known as sour tea, is used for blood pressure control, to relieve constipation and stomach discomfort, lower cholesterol and serves as good source for antioxidants [16,17].

The utilization of plant material in traditional medicine in Nigeria has made treatment of common ailments possible for the poor and many persons in rural areas, who have no access to finance or orthodox medicine [7,18,19]. Herbal medicine will continue to make great contributions to the healthcare of Nigerians. Therefore, it is important to verify the purported therapeutic values of common medicinal plants which form components of herbal remedies sold in Nigeria. The exploration of natural antibiotics derived from plant materials has become an increasingly important area of research today, as it has the potential to contribute to more effective and sustainable approaches to infection management.

This study aimed to assess the antimicrobial effects of *Aloe vera*, *Psidium guajava* and *Hibiscus sabdariffa* extracts on *Salmonella typhi*, *Staphylococcus aureus* and *Escherichia coli* and *Candida albicans*.

2.0 Materials and Methods

2.1 Samples Collection

Fresh leaves of *Aloe vera* stem, *Psidium guajava* and *Hibiscus sabdariffa* were obtained from gardens within delta State University, Abraka Campus II. Leaves were packed in zipper lock bags and taken to the laboratory for processing.

Bacterial pathogens, *Salmonella typhi*, *Staphylococcus aureus* and *Escherichia coli*, and fungal pathogen, *Candida albicans* were obtained from the Eku General hospital, Delta State. The isolates were collected in sterile McCartney bottles and transported in ice box to the laboratory.

2.2 Extraction of Phytochemicals

The collected plant leaves were washed with distilled water, after carefully separating them from the stalk. *P. guajava* and *H. sabdariffa* were dried in a hot air oven for 15 minutes at 40°C, while *Aloe vera* was dried at 67°C for 3 days. After drying, the leaves were ground to fine powder using SCD-P2030D industrial blender.

Extraction was done using soxhlet extractor to which fifty grams (50g) of powdered leaves was added to 200ml ethanol, and allowed to drain into a clean conical flask. Extract was condensed in a rotatory extractor for 10 minutes, to get rid of the ethanol. 10ml of each extract was pipetted into a weight watch glass and evaporated to dryness in a hot water bath until a constant weight was achieved. Dried extracts were enclosed with aluminium foil and stored in a desiccator for 7 days. Extracts were screened for the presence of saponins, tannins, flavonoids and alkaloid, glycosides and phenol as per the procedures highlighted by Shah et al. [20].

2.3 Determination of Minimum Inhibitory Concentration (MIC)

The plant extracts were diluted to obtain the following concentrations: 1:10, 1:100, 1:1000, 1:10,000 and 1:100,000. 1 ml of each dilution was added to 10ml of molten nutrient agar for bacteria and potato dextrose agar for fungi, and then allowed to solidify. Each test isolate was

streaked on the surface of the plates and incubated at 37°C for 24 hours. The least concentration of extract that inhibited microbial growth, was taken as the MBC.

2.4 Susceptibility Test

Serial dilution of each extract was performed to obtain 1:2, 1:3 and 1:4, which were then used to impregnate sterile paper discs. The discs were allowed to soak in the extract for 20 minutes. Sterile swab sticks were used to prepare lawns of test isolates on Muller Hinton Agar plates. Prepared discs were placed on the plates, 2 cm apart, and the plates were then incubated at 37°C for 24 hours. Sensitivity of test organisms was determined by the zone of inhibition.

3.0 Results

3.1 Phytochemicals

Table 1 shows the phytochemical profile of the plants extracts. *Aloe vera* contains saponins, flavonoids and tannins; *P. guajava* contains saponins, alkaloids, flavonoids and tannins, while *H. sabdariffa* contains saponins, alkaloids and flavonoids.

Table 1: Phytochemicals Detected in the Plants Extracts

Plant	Saponins	Alkaloids	Flavonoids	Tannins
<i>Aloe vera</i>	+	-	+	+
<i>Psidium guajava</i>	+	+	+	+
<i>Hibiscus sabdariffa</i>	+	+	+	-

Keys: +=Present; -=Absent

Table 2 shows the concentrations of plant extracts after evaporation, which was 2.0mg/ml for *A. vera* was, 2.1mg/ml for *P. guajava* and 1.83mg/ml for *H. sabdariffa*.

Table 2: Concentration of Plant Extract

Samples	Concentration (mg/ml)
<i>Aloe vera</i>	2.00
<i>Psidium guajava</i>	2.10
<i>Hibiscus sabdariffa</i>	1.83

3.2 Sensitivity of the test organisms

Table 3 shows the sensitivity pattern of the test organisms to the ethanol extracts of *A. vera*, *P. guajava* and *H. sabdariffa*. *Staphylococcus aureus* was only susceptible to *A. vera*. *Salmonella typhi* was not sensitive to all extract. *Escherichia coli* was sensitive to all extracts in the order *H. sabdariffa* > *P. guajava* > *A. vera*. *Candida albicans* was sensitive to *A. vera* and *P. guajava* extracts.

Table 3: Sensitivity Test Clinical Isolates to Extracts

Isolate	<i>A. vera</i>	<i>P. guajava</i>	<i>H. sabdariffa</i>
<i>Staphylococcus aureus</i>	++	-	-
<i>Salmonella typhi</i>	-	-	-
<i>Escherichia coli</i>	+	+++	++

<i>Candida albicans</i>	+	+	-
-------------------------	---	---	---

Keys: - = Not sensitive; +=Sensitive

3.3 Minimum Inhibitory Concentration of Extracts

Table 4 shows the MIC of *A. vera*, *P. guajava* and *H. sabdariffa* extracts against test organisms. The MIC of *A. vera* was 2.0 mg/ml for *S. aureus*, *E. coli* and *C. albicans* respectively. The MIC of *P. guajava* was 0.021 mg/ml and 2.1 mg/ml for *E. coli* and *C. albicans* respectively. The MIC of *H. sabdariffa* was 1.83 mg/ml for *E. coli*.

Table 4: Minimum Inhibitory Concentration of Extracts to Clinical Isolates

Isolate	<i>A. vera</i>	<i>P. guajava</i>	<i>H. sabdariffa</i>
<i>Staphylococcus aureus</i>	2.00	R	R
<i>Salmonella typhi</i>	R	R	R
<i>Escherichia coli</i>	2.00	0.021	1.83
<i>Candida albicans</i>	2.00	2.10	R

Key: R – Resistant

4.0 Discussion

This study assessed the antimicrobial properties of three medicinal plants (*Aloe vera*, *Psidium Guajava* and *Hibiscus sabdariffa*). Medical plants have medicinal values due to the presence of chemical substances normally found on leaves, stems, fruits and seeds. Phytochemical screening of the plant extracts confirmed the presence of flavonoids, tannins, alkaloids and saponins. Plants are used in the treatment of diseases either alone or in combination depending on their phytochemical content [4-6,10]. *Aloe vera* contains saponins, flavonoids and tannins; *P. guajava* contains saponins, alkaloids, flavonoids and tannins, while *H. sabdariffa* contains saponins, alkaloids and flavonoids. This is consistent with reports on the phytochemical screening of these plants [4,12]

The results of the present investigations showed that the leaves of *A. vera*, *P. guajava* and *H. sabdariffa* inhibited some of the test organisms (*Staphylococcus aureus*, *Escherichia coli* and *Candida albicans*) but not *Salmonella typhi*. The antimicrobial activities of the extracts are attributable to the presence of flavonoids, alkaloids, tannins and saponins that were found on these leaves, which had been reported to inhibit pathogenic microorganisms [4,7,10].

Staphylococcus aureus was only susceptible to *A. vera*. Okaba et al. [4] similarly reported susceptibility of *S. aureus* to *A. vera*. *Salmonella typhi* was not sensitive to all extract. The findings of this study agree with Sule et al. [15] on resistance of *Salmonella* to ethanol extract of *P. guajava*. *Escherichia coli* was sensitive to all extracts in the order *H. sabdariffa*>*P. guajava*>*A. vera*. Okaba et al. [4], Lade et al. [21] and Biswas et al. [22] likewise reported sensitivity of *E. coli* to *A. vera*, *H. sabdariffa* and *P. guajava* respectively. *Candida albicans* was sensitive to *A. vera* and *P. guajava* extracts. Okaba et al. [4] similarly reported susceptibility of *C. albicans* to *A. vera*.

The MIC of *A. vera* was 2.0 mg/ml for *S. aureus*, *E. coli* and *C. albicans* respectively. The MIC of *P. guajava* was 0.021 mg/ml and 2.1 mg/ml for *E. coli* and *C. albicans* respectively. The MIC of *H. sabdariffa* was 1.83 mg/ml for *E. coli*. The MICs of the extract are much lower than 25.0 mg/ml reported by Okaba et al. [4] for *S. aureus*, *E. coli* and *C. albicans*.

Escherichia coli, *Salmonella typhi*, *Staphylococcus aureus* and *Candida albicans* are all significant human pathogens that have shown increasing resistance to various antibiotics, posing a significant challenge in clinical management and public health. Resistance has been reported against commonly used antibiotics, such as ampicillin, beta-lactams, carbapenems, chloramphenicol, fluoroquinolones, trimethoprim-sulfamethoxazole, fluconazole, clotrimazole, ketoconazole, which have been previously effective treatments for infectious diseases caused by *Escherichia coli*, *Salmonella typhi*, *Staphylococcus aureus* and *Candida albicans* [23-25]. Susceptibility of these pathogens to plant extracts is encouraging.

Conclusion

The ethanol extracts of *Aloe vera*, *Psidium guajava* and *Hibiscus sabdariffa* were found to contain alkaloids, tannins, flavonoids and saponins. The extracts exhibited antimicrobial activity against *Staphylococcus aureus*, *Escherichia coli* and *Candida albicans*. New antimicrobial agents to combat infectious diseases caused by susceptible microorganisms can be developed using *Aloe vera*, *Psidium guajava* and *Hibiscus sabdariffa* extracts.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

References

1. Albertyn R, Berg A, Numanoglu A, Rode H. Traditional burn care in sub-Saharan Africa: a long history with wide acceptance. *Burns*. 2015; 41:203–211. <https://doi.org/10.1016/j.burns.2014.06.005>
2. Hosseinzadeh S, Jafarikukhdan A, Hosseini A, Armand R. The application of medicinal plants in traditional and modern medicine: a review of *Thymus vulgaris*. *Int J Clin Med*. 2015; 6:635–642.
3. Jaiswal Y, Liang Z, Zhao Z. Botanical drugs in ayurveda and traditional Chinese medicine. *J Ethnopharmacol*. 2016; 194:245–259.
4. Okaba AE, Immanuel OM, Stow, KM, Oku IY. Evaluation of the antimicrobial activities of selected plant extracts and honey against clinical isolates. *African Scientist*. 2022; 23:(3) 30: 175-179.
5. Rojas JJ, Ochoa VJ, Monoz JF. Screening for antimicrobial activity of ten medicinal plants used in Colombian folkloric medicine: a possible alternative in treatment of nonnosocomial infection. *BMC Complement Altern Med*. 2006; 6:1–6.
6. Immanuel OM, Abu GO, Stanley HO. Inhibition of biogenic sulphide production and biocorrosion of carbon steel by sulphate reducing bacteria using *Ocimum gratissimum* essential oil. *Journal of Biology & Biotechnology*. 2016; 10(2): 1-12.
7. Ugboko HU, Nwinyi OC, Oranusi SU, Fatoki TH, Omonhinmin CA. Antimicrobial importance of medicinal plants in Nigeria. *Sci. World J*. 2020; 1 (1): 1-10, 10.1155/2020/7059323
8. Verpoorte DR. New times for traditional medicine research. *J Ethnopharmacol*. 2017; 197:1. <https://doi.org/10.1016/j.jep.2017.01.018>
9. Zhang J, Onakpoya IJ, Posadzki P, Eddouks M. The safety of herbal medicine: from prejudice to evidence. *Evid Based Complement Altern Med*. 2015; 316706, <https://doi.org/10.1155/2015/316706>

10. Iwu M, Duncan A, Okunji C. New antimicrobials of plant origin in. Perspectives on new crops and new uses, In: Plant Breeding Reviews, J. Janick, Ed, ASHS Press, Alexandria, Virginia. 1999.
11. Shakib Z, Shahraki N, Razavi BM, Hosseinzadeh H. *Aloe vera* as an herbal medicine in the treatment of metabolic syndrome: a review. *Phytother Res.* 2019; 33:2649–2660. <https://doi.org/10.1002/ptr.6465>
12. Agarry OO, Olaleye MT, Bello-Michael, CO. Comparative antimicrobial activities of *Aloe vera* gel and leaf. *Afr J Biotechnol.* 2005; 4 (12): 1413-1414.
13. Oghenemaro EF, Johnson J, Itohan IM, Richard SO, Michael O. Antimicrobial activity of *Aloe vera* gel and honey against bacteria isolates from wound aspirates. *Int J Pharm Sci & Res.* 2018; 9(11): 4890-4893.
14. Nwinyi OC, Chinedu NS, Ajani OO. Evaluation of antibacterial activity of *Psidium guajava* and *Gongronema Latifolium*. *J Med Plant Res.* 2008; 2 (8): 189-192.
15. Sule AO, Olalemi AO, Ogundare AO. Efficacy of leaf extracts of *Psidium guajava* (L.) on enteric bacterial isolates from faecally impacted groundwater. *Water Practice & Technology* 2022; 17(1): 167. doi: 10.2166/wpt.2021.113
16. Puro K, Sanjukta R, Samir S. Medicinal Uses of Roselle Plant (*Hibiscus sabdariffa* L.): A Mini Review. *Indian Journal of Hill Farming*, 2014; 27(1):81-90.
17. Pragya S, Mahejibin K, Hailu H. Nutritional and Health Importance of *Hibiscus Sabdariffa*: A Review and Indication for Research Needs. *Journal of Nutritional Health & Food Engineering.* 2017; 6(5):00212
18. Amaeze OU; Aderemi-Williams RI, Ayo-Vaughan MA, Ogundemuren DA, Ogunmola DS, Anyika EN. Herbal medicine use among Type 2 diabetes mellitus patients in Nigeria: understanding the magnitude and predictors of use. *Int J Clin Pharm.* 2018; 40:580–588. <https://doi.org/10.1007/s11096-018-0648-2>
19. Li S, Odedina S, Agwai I, Ojengbede O, Huo D, Olopade OI. Traditional medicine usage among adult women in Ibadan, Nigeria: a cross-sectional study. *BMC Complement Med Ther.* 2020; 20: 93. doi: 10.1186/s12906-020-02881-z
20. Shah P, Modi HA, Shukla MD, Lahiri SM. Preliminary phytochemical analysis and antibacterial activity of *Ganoderma lucidum* collected from Dang District of Gujarat, India. *Int. J. Curr. Microbiol. App. Sci.* 2014; 3(3): 246-255.
21. Lade SN, Burle SS, Kosalge SB, Bansode MN. Antimicrobial and Antioxidant activity of *Hibiscus Sabdariffa*. linn (Roselle) *International Journal of Pharmacy Research & Technology.* 2022; 12(1): doi: 10.31838/ijprt/12.01.04
22. Biswas B, Rogers K, McLaughlin F, Daniels D, Yad A. Antimicrobial Activities of Leaf Extracts of Guava (*Psidium guajava* L.) on Two Gram-Negative and Gram-Positive Bacteria. *International Journal of Microbiology.* 2013; 1-7 pages <http://dx.doi.org/10.1155/2013/746165av1>
23. Ocean HO, Eze EA, Eze CN, Enebe MC. Drug resistance pattern of *Salmonella* and *Shigella* species isolated from selected hospitals in Anyigba, Kogi State, Nigeria. *African Journal of Microbiological Research.* 2015; 9(37), 2023–2036.
24. Gurung RR, Maharjan P, Chhetri GG. Antibiotic resistance pattern of *Staphylococcus aureus* with reference to MRSA isolates from pediatric patients. *Future Sci OA.* 2020 Apr; 6(4): FSO464. doi: 10.2144/foa-2019-0122
25. White TC, Holleman S, Dy F, Mirels LF, Stevens DA. Resistance Mechanisms in Clinical Isolates of *Candida albicans*. *Antimicrob Agents Chemother.* 2002; 46(6): 1704–1713. doi: 10.1128/AAC.46.6.1704-1713.2002