

Effect of integrated nutrient management on meadow orcharding of guava for better growth, yield

Abstract

The present experiment was conducted at Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj. Prayagraj during the session 2023 - 2024. The experiment was laid out in randomized block design with three replications, and the study consists of ten treatment combinations including control by using different Effect of integrated nutrient management on growth yield and quality of guava under meadow orcharding. The best treatment was T₉ (50% RDF + 10 Kg FYM + Azotobacter + VAM) & T₈ (50% RDF + 10 Kg FYM + VAM) which shows highest values in all the parameters viz., fruit weight (145.98 g), fruit diameter (8.00 cm), number fruit/tree (41.81), fruit yield/tree (6.02 kg), yield (t/ha) (30.09). All the treatments were significantly superior in their fruit yield and fruit growth of guava over control (T₀) and (T₉).

Key words: Guava, INM, *Psidium guajava*,

Introduction

One of the most widely grown fruit crops in tropical and subtropical regions is the guava (*Psidium guajava* L.), sometimes referred to as the "apple of the tropics" or the "poor man's apple." It is indigenous to Tropical America, which stretches from Mexico to Peru, and is a member of the Myrtaceae family with chromosome number $2n = 22$. (**Radha and Mathew, 2007**). Early in the 17th century, the Portuguese brought the plant to the Indian subcontinent (**Singh, 1995**). However, at the moment, the main nations that produce guavas are Bangladesh, China, Thailand, India, Mexico, Indonesia, and Pakistan [21-23]. However, it appears to be an Indian fruit since it is readily available, affordable for the average person, and a good source of nutrients (**Dinesh and Vasugi, 2010**).

After mango, banana, citrus, and apple, guava is India's fifth most significant fruit in terms of acreage and output. Approximately 0.27 million hectares are planted with guava in India, yielding 4.10 million tonnes and a productivity of 13.7 million tonnes per hectare. The state

of Uttar Pradesh in India has the greatest area and highest output of guava fruit, whereas Andhra Pradesh has the highest productivity. “In homestead gardens across India, it grows without much attention, but it's also grown commercially in Uttar Pradesh, Bihar, Madhya Pradesh, West Bengal, Punjab, Gujarat, Maharashtra, Karnataka, and Andhra Pradesh”(Anon. 2018-2019).

“In India, guava is the perfect fruit crop for ensuring nutritional security. With 2 to 5 times more fibre (6.9%) than fruit next to fig crop, it is the third-richest source of Vitamin C (299 mg/100g) after Barbados cherry (1000–4000 mg/100g pulp) and Aonla (600 mg/100 g of pulp)”(Gupta, 2014). In addition to being high in vitamin C, it also contains significant amounts of vitamin A, calcium, thiamine (B1), riboflavin (B2), and iron. The fruit has a pleasant scent, and both the peel and the fruit are edible. It is often consumed raw, green, and ripe (when it starts to smell good). Three to thirteen percent of the seed is oil, which is rich in important fatty acids and may be dressed salads. A variety of goods, including jam, jelly, syrup, drinks, sauces, ice cream, butter, marmalade, chutney, and other items, are made from guava. The leaves are used for tanning, drying, and curing diarrhoea in some regions.

Integrated nutrient supply/management, or INM, optimises the benefits from all available plant nutrient sources in an integrated way with the goal of maintaining or adjusting soil fertility and plant nutrient supply to an optimal level for maintaining the targeted crop productivity (Roy and Ange, 1991). Because there are nutrients in the soil for crops in the next season, it increases production levels in a long-term sustainable way. Additionally, by slowly releasing fertilisers, it reduces the current discrepancy between nutrient supply and removal caused by the constant use of chemical fertilisers (Lataet al., 2013). Using organic fertilisers is a standard procedure that helps to reduce the use of chemical treatments and their detrimental effects on the ecosystem as a whole. The most important organic matter added to the soil is high-quality farm yard manure, chicken manure, vermicompost, biofertilizers and green manures.

2. MATERIALS AND METHODS

This experiment was laid out during the August 2023 to March 2024 at Horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.). The horticulture research farm is situated at 25⁰ 39' 42" N latitude, 81⁰ 67' 56" E longitude and at an altitude of 98 m above mean sea level. Six to seven years old uniform trees of guava (*Psidium guajava* L.) cv. Allahabad Safeda under

meadow orcharding were selected for this study at Guava Orchard. The treatment consisted of T₀- 100% RDF (NPK-180,90,90g), T₁- 75% RDF + 2.5 kg vermicompost, T₂- 50% RDF + 2.5 kg vermicompost, T₃ - 75% RDF + 10 kg FYM, T₄- 50% RDF + 10 kg FYM, T₅ (75% RDF + 10 kg FYM) + Micronutrient, T₆- 50% RDF + 10 kg FYM) + Micronutrient, T₇ - 50% RDF + 10 kg FYM + *Azotobacter*, T₈ -50% RDF + 10 Kg FYM + VAM, T₉ -50% RDF + 10 Kg FYM + *Azotobacter*+ VAM. The experiment was laid out in a Randomized Block Design with 10 treatments and replicated thrice. Data recorded on different aspects of fruit crop, viz., growth, yield were subjected to statistical analysis by analysis of variance method. (Gomez and Gomez, 1976) and economic data analysis mathematical method.

RESULT AND DISCUSSION:

Fruit physical parameters

Fruit weight (g)

In data on fruit weight of guava as influenced by integrated nutrient management are summarized in Table 1.

The data reveals that the fruit weight of guava increased significantly by the application of integrated nutrient management under experimentation over the control. The maximum fruit weight (145.98 g) was recorded with treatments 9 (50% RDF + 10 Kg FYM + *Azotobacter* + VAM) respectively, while the minimum fruit weight (118.63 g) was recorded under treatment 0 (control), respectively. Further, the interaction effect of integrated nutrient management significantly influenced the fruit weight in guava.

It may be due to increased the rhizosphere microbial activity and larger quantity of nutrients of the soil. **Ramana et al. (2014)** found that “application of different fertilizers, organic manures and biofertilizer improve the vegetative growth, number of fruits and yield of guava cv. Sardar”. The similar effect were found by **Rao et al. (2014)** and **Ray et al. (2014)**

Fruit Diameter (cm)

In data on fruit girth of guava as influenced by integrated nutrient management are summarized in Table 1.

The data reveals that the fruit diameter of guava increased significantly by the application of integrated nutrient management under experimentation over the control. The

maximum fruit diameter (6.60 cm) was recorded with treatments 9 (50% RDF + 10 Kg FYM + *Azotobacter* + VAM) respectively, while the minimum fruit diameter (5.30) was recorded under treatment 0 (control), respectively. Further, the interaction effect of integrated nutrient management significantly influenced the fruit diameter in guava.

It may be due to increased the rhizosphere microbial activity and larger quantity of nutrients of the soil. **Kumrawat et al., (2018)** found that application of different fertilizers, organic manures and biofertilizer improve the vegetative growth, number of fruits and yield of guava cv. Sardar. The similar effect were found by **Jamwalet al. (2018) and Prabhuet al. (2018)**.

Fruit shape

The data on fruit shape of guava as influenced by integrated nutrient management are summarized in table number 1

The best fruit shape was best recorded with treatment 9(50% RDF + 10 Kg FYM + *Azotobacter* + VAM).

Fruit colour

The data on fruit colour of guava as influenced by integrated nutrient management are summarized in table number 1.

The best fruit colour was best recorded with treatment 9(50% RDF + 10 Kg FYM + *Azotobacter* + VAM).

Yield Parameter

Number of fruits/plant

In data on number of fruits/plant of guava as influenced by integrated nutrient management are summarized in Table 2

The data reveals that the number of fruits/plant of guava increased significantly by the application of integrated nutrient management under experimentation over the control. The maximum number of fruits/plant (41.81) was recorded with treatments 9 (50% RDF + 10 Kg FYM + *Azotobacter* + VAM) respectively, while the minimum number of fruits/plant (20.92)

was recorded under treatment 0 (control), respectively. Further, the interaction effect of integrated nutrient management significantly influenced the number of fruits/plant in guava.

Tiwari *et al.*, (2018) reported that “the application of different fertilizers, organic manures and biofertilizers improve the vegetative growth, number of fruits and yield of guava cv. Sardar”. Similar findings have been reported by **Porismita *et al.*, (2018)**.

Yield/plant (kg)

In data on yield/plant of guava as influenced by integrated nutrient management are summarized in Table 2

The data reveals that the yield/plant of guava increased significantly by the application of integrated nutrient management under experimentation over the control. The maximum yield/plant (41.81) was recorded with treatments 9 (50% RDF + 10 Kg FYM + *Azotobacter* + VAM respectively, while the minimum yield/plant (20.92) was recorded under treatment 0 (control), respectively. Further, the interaction effect of integrated nutrient management significantly influenced the yield/plant in guava.

Sandhyarani *et al.*, (2022) found that “application of different fertilizers, organic manures and biofertilizer improve the vegetative growth, number of fruits and yield of guava cv. Sardar”. The similar effect were found by **Gupta *et al.*, (2019)**.

Yield (t/ha)

In data on yield (t/ha) of guava as influenced by integrated nutrient management are summarized in Table 2.

The data reveals that the yield (t/ha) of guava increased significantly by the application of integrated nutrient management under experimentation over the control. The maximum yield (t/ha) (41.81) was recorded with treatments 9 (50% RDF + 10 Kg FYM + *Azotobacter* + VAM respectively, while the minimum yield (t/ha) (20.92) was recorded under treatment 0 (control), respectively. Further, the interaction effect of integrated nutrient management significantly influenced the yield (t/ha) in guava.

Shubham *et al.*, (2020) found that “application of different fertilizers, organic manures and biofertilizer improve the vegetative growth, number of fruits and yield of guava cv. Sardar”. The similar effect were found by **Jaiprakash *et al.*, (2021)** and **Tyagi *et al.*, (2021)**.

CONCLUSION

In the present investigation concluded that among the different treatment combinations the treatment T9 [50% RDF + 10 Kg FYM + *Azotobacter* + VAM) was superior in respect to fruit growth and yield parameters.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

Table 1. Effect of different levels Integrated Nutrient Management on fruit growth of Guava

Treatment Symbol	Treatment combinations				
		Fruit weight (g)	Fruit diameter (cm)	Fruit shape	Fruit colour
T ₀	100% RDF (NPK-180,90,90g)	118.63	5.91	Oval	Dark green
T ₁	75% RDF + 2.5 kg vermicompost	125.40	6.50	Oval	Dark green
T ₂	50% RDF + 2.5 kg vermicompost	120.10	6.21	Oval	Dark green
T ₃	75% RDF + 10 kg FYM	138.61	7.60	Round	Green
T ₄	50% RDF + 10 kg FYM	135.30	7.55	Round	Green
T ₅	(75% RDF + 10 kg FYM) + Micronutrient	129.81	6.88	Round	Green
T ₆	(50% RDF + 10 kg FYM) + Micronutrient	131.90	6.95	Oval	Dark green
T ₇	50% RDF + 10 kg FYM + <i>Azotobacter</i>	140.52	7.81	Round	Yellowish
T ₈	50% RDF + 10 Kg FYM + VAM	142.85	7.90	Round	Yellowish
T ₉	50% RDF + 10 Kg FYM + <i>Azotobacter</i> + VAM	145.98	8.00	Round	Yellowish
F-test		S	S	-	-
SEm(±)		3.94	0.24	-	-
CD (p=0.05)		11.70	0.73	-	-

Table 2. Effect of different levels Integrated Nutrient Management on fruit yield of Guava

Treatment Symbol	Treatment combinations			
		Number of fruits/plant	Yield/plant (kg)	Yield (t/ha)
T ₀	100% RDF (NPK-180,90,90g)	20.92	2.49	12.47
T ₁	75% RDF + 2.5 kg vermicompost	25.71	3.27	16.37
T ₂	50% RDF + 2.5 kg vermicompost	22.60	2.71	13.56
T ₃	75% RDF + 10 kg FYM	35.31	4.89	24.47
T ₄	50% RDF + 10 kg FYM	32.51	4.35	21.73
T ₅	(75% RDF + 10 kg FYM) + Micronutrient	28.60	3.72	18.60
T ₆	(50% RDF + 10 kg FYM) + Micronutrient	30.48	4.06	20.31
T ₇	50% RDF + 10 kg FYM + <i>Azotobacter</i>	38.98	5.28	26.38
T ₈	50% RDF + 10 Kg FYM + VAM	39.05	5.55	27.76
T ₉	50% RDF + 10 Kg FYM + <i>Azotobacter</i> + VAM	41.81	6.02	30.09
F-test		S	S	S
SEm(±)		0.92	0.13	0.66
CD (p=0.05)		2.72	0.39	1.95

REFERENCES

1. Anonymous, (2018-19). Indian Horticulture Database, National Horticulture Board, Gurgaon, Govt. of India.
2. Dinesh, M.R. and Vasugi, C. (2010). Guava improvement in India and future needs. *J. Hort. Sci.*, **5**(2): 94-108.
3. Gupta, S.N. (2014). Instant Horticulture. Jain Brothers, New Delhi, pp. 206.
4. Jaiprakash, Prasad, V.M., Bahadur, V. and Dinesh (2021). Studies on Impact of Organic and Inorganic Fertilizers on Growth, Yield and Quality of Guava (*Psidiumguajava*) Allahabad Safeda Under Sub Tropical Climatic Conditions. *Biological Forum – An International Journal*, **13**(3a): 285-288.
5. Jamwal, S., Mishra, S. and Singh, S. 2018. Effects of integrated nutrient management on physical characteristics of guava under meadow orcharding cv. Allahabad Safeda. *J. Pharm. and Phytochemi.*, **1**: 2076-2079.
6. Kumrawat D, Kanpure RN, Singh OP, Bhandari J, and Kachouli B (2018) Effect of integrated nutrient management on quality and yield parameters of guava (*Psidiumguajava* L.) cv. I-49. *Journal of Pharmacognosy and Phytochemistry* **7**(5): 1668-1670
7. Lata, R., Dwivedi, D.H., Ram, R.B, Meena, M.L. and Babu, M. 2013. Impact of integrated nutrient management on growth parameters of strawberry cv. Chandler under sub-tropical conditions of lucknow. *Int. J. Adv. Biol. Res.*, **3**(3): 418-421.
8. Muthukumar, P. and Selvakumar, R. (2013). *Glaustas Horticulture*. New Delhi, New vishal publications, p. 120.
9. Porismita Dutta, Utpalkotoky and Kaushik Das. Effect of Integrated Plant Nutrient Management On Growth and Yield Of Guava (*Psidiumguajava* L.) cv. Allahabad Safeda under meadow orcharding . *Bull. Env. Pharmacol. Life Sci.*, Vol 7 [11] October 2018: 01-06
10. Prabhu, M., Parthiban, S., Kumar, A.R., Rani, B.U. and Vijayasamundeeswari. 2018. Effect of integrated nutrient management on acidlime (*Citrus aurantifolia*. Swingle. (L.)). *Indian Journal of Agricultural Research*, **52**(3): 290-294.
11. Radha, T. and Mathew, L. (2007). *Fruit crops*. New Delhi, New India publishing agency, p. 59.
12. Ramana, K. T. V., Lakshmi, L. M., Gopal, K., Krishna, V. N. P. S., Lakshmi, T. N., Sarada, G., Gopi, V. and Sankar, T. G. 2014. Nitrogen and potassium based fertigation response on plant growth, yield and quality of sweet orange (*Citrus sinensis* Linn. Osbeck) cv. Sathgudi. *Research and Reviews: J. Agril. and Allied Sci.*, **3**(3): 7- 10.

13. Rao, B. N., Roy, S. S., Jha A. K., Deo, C. and Kumar, S. 2014. Influence of nitrogen and irrigation on growth, yield and quality of passion fruit (*Passiflora edulis* Sims.) under foothill condition of Manipur. *Progressive Hort.*, **46**(2): 255-259.
14. Ray, S. K. D.; Takawale, P.; Chatterjee, V. R. and Hanamte, E. V. 2014. Yield and quality of pomegranate as influenced by organic and inorganic nutrients. *An International Quarterly J. Life Sci.*, **9**(2): 617-620.
15. Roy, R. N. and Ange, A. L. 1991. In: Integrated plant nutrient systems (IPNS) and sustainable agriculture. Proceedings FAI Annual Seminar, FAI, New Delhi, SV 1-1:1-12.
16. Sandhyarani M., A. Bhagwan, A. Kiran Kumar and M. Sreedhar (2022). Studies on effect of Biofertilizers and Biostimulant on Post Harvest Quality Parameters and Shelf Life of Guava (*Psidium guajava* L.) cv. Allahabad Safeda under meadow orcharding under Meadow Planting System. *Biological Forum – An International Journal*, **14**(2): 1166-1169.
17. ShubhamMangal, SS Singh, Sant Kumar, NR Rangare. Impact of inorganic, organic and bio-fertilizers on growth and yield of guava (*Psidium guajava* L.). *Int J Chem Stud* 2020;**8**(6):101-107.
18. Singh, S. P. (1995). Commercial fruits. New Delhi, Kalyani publisher, p. 148.
19. Tiwari Neelam, SINGH S. S. Singh Rajesh and Charmkar Neeraj K. (2018) Effect of organic-cum-inorganic sources of nutrients on growth, yield and economical gain from guava *International Journal of Agricultural Science and Research***8**(1) 79-84.
20. Tyagi S K, G S Kulmi and A R Khire (2021). Effect of Integrated Nutrient Management on Growth, Yield and Economics of Guava (*Psidium guajava* L.). *J Krishi Vigyan*, **10** (1) :69-72.
21. Anmol, Devi Singh, Vaibhav Kapadnis, and Sweekritilsha Ekka. 2023. "Effect of Integrated Nutrient Management on Physical Characteristics of Guava (*Psidium Guajava*) Cv. Allahabad Safeda under Meadow Orchardling". *International Journal of Environment and Climate Change* 13 (8):1615-23. <https://doi.org/10.9734/ijecc/2023/v13i82111>.
22. Mohapatra ,Sheetal, Sarat Chandra Sahoo, DeepaSamant, and Subash Chandra Swain. 2023. "Influence of Integrated Nutrient Management on Physico-Chemical Properties of Soil, Growth, Flowering and Yield of Guava Grown As a Component Crop in Coconut-Based Cropping System". *International Journal of Plant & Soil Science* 35 (22):102-11. <https://doi.org/10.9734/ijpss/2023/v35i224117>.
23. Sun, G., Hu, T., Liu, X., Peng, Y., Leng, X., Li, Y., & Yang, Q. (2022). Optimizing irrigation and fertilization at various growth stages to improve mango yield, fruit quality and water-fertilizer use efficiency in xerothermic regions. *Agricultural Water Management*, 260, 107296.
24. Gupta P, Singh D, Prasad VM, Kumar V. Effect of integrated nutrient management on growth and yield of guava (*Psidium guajava* L.) cv. Allahabad safeda under high density planting. *Journal of Pharmacognosy and Phytochemistry*. 2019;**8**(1):1233-6.

