

To Study Correlation Between Yield and its Components in Eggplant (*Solanum melongena* L.)

Comment [I1]: There are different types of citations that are used in the article, e.g. (16,17,18,19,20,21,22), (Aykroyd, 1963) and (Chen and Li, 1996³). Please use the citation style of the journal

Abstract

The present investigation, entitled "Investigation on the genetic basis of certain metric traits in the brinjal (*Solanum melongena* L.) summer crop under Alwar (Rajasthan) conditions," was under way. In this research, 74 different genotypes, including 14 lines, 4 tests, and 56 F1 hybrids, along with two control checks, were ~~evaluated~~ ~~assessed~~ using a randomized block design with three replications at the Vegetable Research Farm, ~~of the~~ Department of Vegetable Science, Sunrise University, located in Alwar, Rajasthan, India. The evaluations were conducted over two summer seasons in 2022 and 2023. Various traits such as days to 50% flowering, days to first fruit harvest, plant height, number of primary branches per plant, pedicle length, number of fruits per cluster, average fruit weight, number of fruits per plant, total ~~phenolic~~ ~~phenol~~ content, dry matter content, reducing sugar content, non-reducing sugar content, total sugars content, TSS (Brix), ascorbic acid content, total fruit yield per plant, and yield per hectare (Q/Ha) were recorded. ~~Additionally,~~ ~~A~~ ancillary traits were also observed. The findings indicated that during the first year, there was a strong positive correlation between average fruit weight and several traits including total fruit yield per plant, yield per hectare, non-reducing sugar content, and number of fruits per plant. ~~Similarly,~~ in the second year, ~~average fruit weight~~ exhibited significant positive correlations with total fruit yield per plant, yield per hectare, ~~average fruit weight~~, number of fruits per cluster, and TSS. Pooling ~~the of~~ data also ~~showed~~ ~~revealed~~ a strong positive correlation between average fruit weight and total fruit yield per plant, yield per hectare, non-reducing sugar content, and number of fruits per plant. In terms of genotypic correlations in the first year, average fruit weight displayed strong positive correlations with total fruit yield per plant, yield per hectare, non-reducing sugar content, number of fruits per plant, and total ~~phenolic~~ ~~phenol~~ content.

Comment [I2]: The abstract should not include abbreviations.

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Red

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Red, Highlight

Comment [I3]: There is a problem with this comment and it should be reviewed again.

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Red

Keywords: genotypic, phenotypic correlation

Introduction

Eggplant, also known as brinjal (*Solanum melongena* L.), ~~is an important~~ ~~holds significance as a~~ member of the ~~Solanaceae family of vegetables~~ ~~Solanaceous vegetable crops~~ with a diploid chromosome ~~count number~~ of $2n=2x=24$. Per 100 grams of its edible ~~part~~ ~~portion~~, eggplant fruit contains 92.7% moisture, 24.0% calories, 4.0% carbohydrates, 1.4 grams of protein, 0.3 grams of fats, 1.3 grams of fibers, 124.0 (I.U.) of vitamin A, and 12.0 milligrams of vitamin C (Aykroyd²).

Comment [I4]: More up-to-date literature can be added.

Formatted: Font: Italic

It also ~~contains~~boasts 52.0 milligrams of chlorine, 47.0 milligrams of phosphorus, 44.0 milligrams of ~~sulphur~~sulfur, and other essential minerals (Aykroyd, 1963). Eggplant is attributed with medicinal properties, ~~or example, such as~~white eggplant ~~is considered~~being regarded as beneficial for diabetics ~~patients~~-(Choudhary, 1976¹). ~~In addition~~Additionally, fried eggplant fruit in ~~fil~~oil is traditionally ~~used to relieve~~employed for ~~alleviating~~toothache (Chen and Li, 1996³), while it has been advocated as an effective remedy for ~~those individuals~~ suffering from liver ~~problems~~complaints. Calculating the correlation coefficient ~~between~~among yield-contributing traits is essential for deciphering the direction of selection and ~~optimis~~izing yield within a condensed timeframe. Path coefficients ~~provide~~offer a streamlined approach to ~~disentangling~~disentangle both direct and indirect factors influencing selection associations, thereby measuring the relative significance of each causal factor. ~~Since~~As yield is a multifaceted trait influenced by ~~numerous~~many attributes, understanding the relationship between ~~different~~traits ~~various~~characteristics and both yield and its contributing factors is vital for effective selection processes.

Comment [I6]: If this paragraph is a quotation, please give a reference

Materials and Methods

Comment [I7]: Which statistical programme was used in the evaluation of the data?

The study ~~utilized~~used 56 F1 hybrids, ~~obtained~~generated from crosses between 14 different lines (Rampur local-1, Rampur local-2, Bareilly local-1, Bareilly local-2, Alwar local, Jaipur local, Udaipur local, Moradabad local, Delhi local, Suvad Shakti, Suvad Sri, Mukta Keshi, Pant brinjal hybrid-1, MDU-1) and 4 testers (Pant Samrat, Pant Rituraj, Annamalai, Azad Kranti), along with 2 ~~check~~control varieties (Punjab Barsati, Pusa Bharav) during the summers of 2022 and 2023. The performance of these hybrids and lines was evaluated using a Randomized Complete Block Design (RCBD) with ~~three~~replications. ~~The~~Plants were spaced at 0.60 ~~metres~~meters apart ~~within~~in rows. ~~The~~Trials were conducted ~~during~~across the summer seasons of 2022 and 2023. Various parameters ~~including~~days to 50% flowering, days to first fruit harvest, plant height, number of primary branches per plant, pedicle length, number of fruits per cluster, average fruit weight, number of fruits per plant, total ~~phenolic~~phenol content, dry matter content, reducing sugar content, non-reducing sugar content, total sugars content, TSS (Brix), ascorbic acid content, total fruit yield per plant, and yield per hectare were recorded. Ancillary ~~characteristic~~traits were also ~~recorded~~noted. The paraphrased sources for this information are Suyin *et. al.*,¹³ Goto *k et. al.*,⁵ Pandey *et. al.*,⁷ and Prabhu *et. al.*,⁹.

Comment [I8]: What is the number of plants of each genotype used?

Comment [I9]: This section should group traits under phenotypic and genotypic traits.

Comment [I10]: Add explanation of TSS

Comment [I11]: The units of the characteristics received must be specified. For example cm, day, % etc.

RESULTS AND DISCUSSION

Phenotypic Correlation Coefficient

Comment [I12]: Please discuss the characteristics with more up-to-date literature.

In the first year, ~~d~~Days to 50% flowering ~~showed a marked~~displayed a ~~notably~~positive and significant correlation with ~~d~~Days to first fruit harvesting (0.336), TSS (274), and ~~r~~Reducing ~~s~~Sugar (0.184) at the phenotypic level. Conversely, ~~y~~Yield Q/Ha (-0.258), ~~n~~Number of primary branches per plant (-0.195), and ~~n~~Number of fruits per cluster (-0.140) exhibited negative and

significant correlations with this trait. No other traits showed significant correlations with **dDays** to 50% flowering. Similarly, in the second year, **dDays** to 50% flowering demonstrated strong positive correlations with **dDays** to first fruit harvest~~ing~~ (0.550), **rReducing sSugar** (0.268), and **Ttotal sugar** (0.202), while **nNumber** of primary branches per plant (-0.275) and **nNumber** of fruits per cluster (-0.143) displayed negative and significant correlations. **Once aAgain**, no other traits were significantly correlated with **dDays** to 50% flowering. In the pooled data, **dDays** to 50% flowering maintained highly positive and significant correlations with **dDays** to first fruit harvest~~ing~~ (0.514) and **rReducing sSugar** (0.259), while **nNumber** of primary branches per plant (-0.299) and **nNumber** of fruits per cluster (-0.161) showed negative and significant correlations. No other traits showed significant correlations with **dDays** to 50% flowering.

Regarding **dDays** to first fruit harvest~~ing~~ in the first year, it exhibited strong positive and significant correlations with length of pedicle (0.203) and **tTotal fruit yield per plant** (0.173), whereas plant height (-0.224) and **nNumber** of fruits per cluster (-0.203) displayed negative and significant correlations. No other traits were significantly correlated with **dDays** to first fruit harvest~~ing~~. In the second year, **dDays** to first fruit harvest~~ing~~ exhibited a notably positive and significant correlation with **aAscorbic acid** (0.178). Conversely, **nNumber** of primary branches per plant (-0.370), **nNumber** of fruits per cluster (-0.327), and **pPlant height** (-0.326) displayed negative and significant correlations with this trait. No other traits showed significant correlations with **dDays** to first fruit harvest~~ing~~. When considering pooled data, **dDays** to first fruit harvest~~ing~~ demonstrated positive and significant correlations with **aAscorbic acid** (0.156), length of pedicle (0.149), and **aAverage fruit weight** (0.147). However, **nNumber** of fruits per cluster (-0.418), **nNumber** of primary branches (-0.349), and **pPlant height** (-0.260) exhibited highly negative and significant correlations with the trait. No other traits were significantly correlated with **dDays** to first fruit harvest~~ing~~.

Regarding **pPlant height** in the first year, it showed notably positive and significant correlations with **rReducing sugar** (0.262) and dry matter content (0.220). Conversely, **tTotal fruit yield per plant** (-0.368), **aAverage fruit weight** (-0.291), and **nNumber** of fruits per plant (-0.219) displayed negative and significant correlations with this trait. No other traits were significantly correlated with **pPlant height**. In the second year, plant height displayed a positive and significant correlation at the phenotypic level with **nNumber** of primary branches per plant

Formatted: Highlight

(0.162). Conversely, **t**Total phenolic content (-0.311), **d**Dry matter content (-0.182), and **l**Length of pedicle (-0.175) exhibited highly negative and significant correlations with the trait. No other traits showed significant correlations with plant height. **In pooled data, plant height demonstrated a positive and significant correlation at the phenotypic level with n**Number of fruits per plant(0.143). However, **t**Total fruit yield per plant (-0.312), **a**Average fruit weight (-0.287), **t**Total phenolic content (-0.269), **l**Length of pedicle (-0.241), and **n**Number of fruits per plant (-0.233) showed highly negative and significant correlations with the trait. No other traits were significantly correlated with plant height.

In the first year, **n**Number of primary branches per plant exhibited highly positive and significant correlations at the phenotypic level with **a**Average fruit weight (0.449), **n**Number of fruits per cluster (0.422), **t**Total fruit yield per plant (0.356), **l**Length of pedicle (0.235), and **n**Number of fruits per plant (0.228). No data showed negative and significant correlations with the trait.

Formatted: Font color: Red

Formatted: Font color: Red, Highlight

Comment [113]: There is a problem with this comment and it should be reviewed again. **plant height demonstrated a positive and significant correlation with number of fruits per cluster, reducing sugar and, ascorbic acid.**

UNDER PEER REVIEW

Table 1. Genotypic and phenotypic correlation coefficients between different characters of brinjal in Y1 (first year)

Comment [114]: Please provide an explanation of the acronyms used in the table.

Characters		Days to 50% Flowering	Days to first fruit harvest	Plant height	Number of primary branches per plant	Length of pedicel (cm)	Number of fruits per cluster	Average fruit weight (gm)	Number of fruits per plant	Total Phenolic Content (mg/100 gm)	Dry matter content (%)	Reducing Sugar (%)	Non-Reducing Sugar (%)	Total Sugar (%)	TSS (Brix)	Ascorbic acid (mg/100 gm)	Total fruit yield per plant (kg)	Yield Q/Ha
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
1	G	1.000	0.460**	0.061	-0.236**	0.143*	-0.226**	0.107	-0.034	-0.074	0.073	0.215**	0.062	0.094	0.158*	0.090	-0.109	-0.214**
	P	1.000	0.336**	0.065	-0.195**	0.031	-0.140*	0.043	-0.032	-0.031	0.008	0.184**	0.072	0.066	0.274**	0.066	0.003	-0.258**
2	G			-0.204**	-0.151*	0.256**	-0.135*	0.099	0.117	-0.023	0.050	0.092	0.085	0.032	0.099	0.074	0.201**	-0.137*
	P			-0.224**	-0.131*	0.244**	-0.203**	0.151*	0.071	-0.069	0.095	0.070	0.063	0.051	-0.006	0.158*	0.173**	0.022
3	G				-0.091	-0.124	0.012	-0.236**	-0.295**	0.213**	0.330**	0.268**	-0.116	0.185**	-0.144*	0.222**	0.395**	-0.152*
	P				-0.109	-0.113	0.030	-0.291**	-0.219**	-0.135*	0.220**	0.262**	-0.122	0.160*	-0.017	0.156*	0.368**	-0.120
4	G					0.297**	0.458**	0.486**	0.313**	0.010	-0.176**	-0.113	0.147*	0.003	0.072	-0.102	0.353**	0.251**
	P					0.235**	0.422**	0.449**	0.228**	0.012	-0.099	-0.119	0.146*	-0.020	0.021	-0.065	0.356**	0.170*
5	G						0.007	0.269**	0.291**	-0.001	-0.025	-0.071	0.168*	-0.138*	0.392**	0.181**	0.327**	0.010
	P						-0.030	0.239**	0.286**	-0.022	-0.018	-0.049	0.146*	-0.111	0.160*	0.172**	0.184**	0.087
6	G							0.369**	0.297**	0.176**	0.045	0.061	0.232**	0.203**	-0.029	0.038	0.331**	0.358**
	P							0.279**	0.263**	0.199**	0.016	0.059	0.235**	0.170*	0.038	-0.034	0.312**	0.161*
7	G								0.348**	0.183**	-0.062	-0.112	0.366**	-0.031	0.107	-0.161*	0.569**	0.454**
	P								0.268**	0.090	0.024	-0.111	0.338**	-0.004	-0.009	-0.091	0.467**	0.390**
8	G									0.202**	0.101	-0.068	0.237**	-0.078	0.073	0.030	0.905**	0.453**
	P									0.191**	0.009	-0.049	0.217**	-0.061	0.110	-0.005	0.736**	0.348**
9	G										0.255**	0.198**	0.460**	0.381**	-0.085	0.035	0.307**	0.273**
	P										0.186**	0.187**	0.438**	0.330**	0.031	-0.003	0.262**	0.157*
10	G											0.350**	0.156*	0.437**	0.110	0.060	0.140*	0.185**
	P											0.305**	0.142*	0.389**	-0.078	0.118	0.122	0.134*
11	G												-0.141*	0.802**	0.244**	0.185**	0.047	-0.101
	P												-0.142*	0.786**	0.180**	0.152*	0.018	-0.083
12	G													0.140*	0.076	0.066	0.341**	0.263**

Table 3. Genotypic and phenotypic correlation coefficients between different characters of brinjal in pooled

Comment [116]: Please provide an explanation of the acronyms used in the table.

Chara cters		Days to 50% Flowerin g	Days to first fruit harvest	plant height	Number of primary branches per plant	Length of pedicle (cm)	Number of fruits per cluster	Average fruit weight (gm)	Number of fruits per plant	Total Phenolic Content (mg/100 gm)	Dry matter content (%)	Reducin g Sugar (%)	Non- Reducin g Sugar (%)	Total Sugars (%)	TSS (Brix)	Ascorbi cacid (mg/10 0gm)	Total fruit yield per plant (kg)	Yield O/Ha
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
1	G	1.000	0.584**	0.102	-0.315**	0.085	-0.198**	0.083	0.052	-0.051	0.095	0.273**	0.058	0.154*	0.139*	0.136*	0.034	-0.005
	P	1.000	0.514**	0.101	-0.299**	0.052	-0.161*	0.056	0.039	-0.033	0.057	0.259**	0.066	0.142*	0.153*	0.110	0.074	-0.062
2	G			-0.259**	-0.370**	0.166*	-0.417**	0.127	0.133*	-0.031	0.102	0.114	0.047	0.018	0.004	0.138*	0.132*	0.009
	P			-0.260**	-0.349**	0.149*	-0.418**	0.147*	0.127	-0.055	0.098	0.104	0.038	0.029	0.002	0.156*	0.115	0.067
3	G				0.044	-0.256**	0.155*	-0.289**	-0.265**	0.305**	0.069	0.141*	-0.153*	0.091	-0.042	0.160*	0.331**	-0.078
	P				0.029	-0.241**	0.143*	-0.287**	-0.233**	0.269**	0.047	0.137*	-0.155*	0.084	0.013	0.152**	0.312**	-0.086
4	G					0.185**	0.268**	0.183**	0.006	-0.052	-0.055	-0.128	0.080	-0.035	0.091	-0.123	-0.015	-0.204**
	P					0.148*	0.256**	0.155*	-0.022	-0.046	-0.021	-0.126	0.076	-0.043	0.062	-0.101	0.008	-0.178**
5	G						0.026	0.207**	0.275**	-0.093	-0.062	-0.167*	0.009	0.244**	0.345**	-0.079	0.262**	0.110
	P						-0.005	0.199**	0.257**	-0.096	-0.041	-0.153*	0.006	0.228**	0.240**	-0.080	0.207**	0.112
6	G							0.328**	0.040	0.142*	0.150*	0.041	0.071	0.086	0.115	0.007	0.158*	0.190**
	P							0.247**	0.032	0.139*	0.118	0.040	0.079	0.073	0.092	-0.016	0.149*	0.145*
7	G								0.332**	0.200**	0.128	-0.012	0.368**	0.065	0.147*	-0.013	0.535**	0.435**
	P								0.257**	0.149*	0.111	-0.012	0.337**	0.077	0.069	-0.023	0.442**	0.360**
8	G									-0.078	-0.084	0.014	0.024	-0.038	0.035	0.183**	0.859**	0.618**
	P									-0.089	-0.102	0.010	0.019	-0.028	0.073	0.166*	0.723**	0.540**
9	G									0.125	0.133*	0.538**	0.411**	-0.032	-0.063	0.092	0.079	
	P									0.086	0.130	0.513**	0.387**	-0.001	-0.058	0.082	0.036	
10	G											0.235**	0.052	0.231**	0.147*	0.172**	-0.059	0.010
	P											0.211**	0.047	0.197**	0.078	0.170*	-0.023	0.018
11	G												0.177**	0.835**	0.240**	0.139*	0.042	-0.040
	P												0.176**	0.828**	0.220**	0.132*	0.031	-0.033
12	G													0.273**	0.046	0.062	0.146*	0.147*

	P														0.268**	0.032	0.052	0.141*	0.120
13	G															0.273**	0.073	0.040	0.004
	P															0.238**	0.068	0.031	0.008
14	G																0.262**	0.109	0.069
	P																0.232**	0.103	0.014
15	G																	0.145*	0.222**
	P																	0.125	0.220**
16	G																		0.759**
	P																		0.591**
17	G																		1.000
	P																		1.000

UNDER PEER REVIEW

In all other cases, there were no significant correlations found with the number of primary branches per plant. In the second year, the number of primary branches per plant exhibited a positive and significant correlation at the phenotypic level with the nNumber of fruits per cluster (0.154). However, yYield Q/Ha (-0.181) displayed a highly negative and significant correlation with this trait. No other traits were significantly correlated with the nNumber of primary branches per plant. In pooled data, the nNumber of primary branches per plant showed a highly positive and significant correlation at the phenotypic level with the length of pedicle (-0.185). Conversely, yYield Q/Ha (-0.204) displayed a highly negative and significant correlation with the trait. Not all other traits were significantly correlated with the number of primary branches per plant.

Formatted: Indent: First line: 0.49"

Formatted: Highlight

Comment [I17]: There is a problem with this comment and it should be reviewed again. There is a highly positive and significant correlation with number of fruits per cluster.

Formatted: Highlight

In the first year, the lLength of pedicle demonstrated highly positive and significant correlations at the phenotypic level with nNumber of fruits per plant (0.286), aAverage fruit weight (0.239), and tTotal fruit yield per plant (0.184). Conversely, aAscorbic acid (-0.172) displayed a negative and significant correlation with the trait. For all other traits, there were no significant correlations found with the length of pedicle. In the second year, the length of pedicle exhibited highly positive and significant correlations at the phenotypic level with tTotal fruit yield per plant (0.365), yield Q/Ha (0.351), aAverage fruit weight (0.341), nNumber of fruits per cluster (0.289), and TSS (0.188). Conversely, tTotal sSugar (-0.220) and rReducing sugar (-0.184) displayed highly negative and significant correlations with the trait. No other traits were significantly correlated with the length of pedicle. In pooled data, the length of pedicle demonstrated highly positive and significant correlations at the phenotypic level with nNumber of fruits per plant (0.257), TSS (0.240), tTotal fruit yield per plant (0.207), and aAverage fruit weight (0.199). However, tTotal sugar (-0.228) showed a highly negative and significant correlation with the trait. No other traits were significantly correlated with the length of pedicle.

In the first year, nNumber of fruits per cluster exhibited highly positive and significant correlations at the phenotypic level with tTotal fruit yield per plant (0.312), aAverage fruit weight (0.279), nNumber of fruits per plant (0.263), nNon-reducing sugar (0.235), and tTotal phenolic content (0.199). No data showed negative and significant correlations with the trait. No other traits were significantly correlated with the days to nNumber of fruits per cluster. In the second year, nNumber of fruits per cluster displayed highly positive and significant correlations

at the phenotypic level with t Total phenolic content (0.289), n Number of fruits per cluster (0.203), t Total fruit yield per plant (0.189), and y Yield Q/Ha (0.176). No traits showed highly negative and significant correlations with the trait, and all other traits were not significantly correlated with the number of fruits per cluster. In pooled data, n Number of fruits per cluster showed highly positive and significant correlations at the phenotypic level with a Average fruit weight (0.247), while no traits showed highly negative and significant correlations with the trait. All other traits were not significantly correlated with the n Number of fruits per cluster.

In the first year, a Average fruit weight demonstrated highly positive and significant correlations at the phenotypic level with t Total fruit yield per plant (0.467), y Yield Q/Ha (0.390), n Non-reducing sugar (0.338), and n Number of fruits per plant (0.268). No traits showed negative and significant correlations with the trait, and all other traits were not significantly correlated with the days to a Average fruit weight. In the second year, a Average fruit weight displayed highly positive and significant correlations at the phenotypic level with t Total fruit yield per plant (0.365), y Yield Q/Ha (0.351), a Average fruit weight (0.341), n Number of fruits per cluster (0.289), and TSS (0.188). Conversely, t Total Sugar (-0.220) and r Reducing sugar (-0.184) showed highly negative and significant correlations with the trait, while all other traits were not significantly correlated with a Average fruit weight.

In pooled data, a Average fruit weight displayed highly positive and significant correlations at the phenotypic level with t Total fruit yield per plant (0.442), y Yield Q/Ha (0.360), n Non-reducing s Sugar (0.337), and n Number of fruits per plant (0.257). Conversely, t Total sugar (-0.228) exhibited a highly negative and significant correlation with the trait, while all other traits were not significantly correlated with a Average fruit weight.

In the first year, n Number of fruits per plant exhibited highly positive and significant correlations at the phenotypic level with t Total fruit yield per plant (0.736), y Yield Q/Ha (0.348), n Non-reducing sugar (0.217), and t Total phenolic content (0.191). No data showed negative and significant correlations with the trait, and no traits were significantly correlated with the days to n Number of fruits per plant. In the second year, n Number of fruits per plant displayed highly positive and significant correlations at the phenotypic level with t Total fruit yield per plant (0.701), yield Q/ha (0.700), and a Ascorbic acid (0.179). Conversely, t Total phenolic content showed a highly negative and significant correlation with the trait, while all other traits were not

significantly correlated with the n Number of fruits per plant. In pooled data, n Number of fruits per plant showed highly positive and significant correlations at the phenotypic level with t Total fruit yield per plant (0.723) and Y ield Q/Ha (0.540). No data showed highly negative and significant correlations with the trait, and all other traits were not significantly correlated with the N number of fruits per plant.

In the first year, T total phenolic content exhibited highly positive and significant correlations at the phenotypic level with N non-reducing sugar (0.438), T total sugar (0.330), T total fruit yield per plant (0.262), and D dry matter (0.186). No data showed negative and significant correlations with the trait, and all other traits were not significantly correlated with the days to T total phenolic content. In pooled data, A average fruit weight displayed highly positive and significant correlations at the phenotypic level with T total fruit yield per plant (0.442), Y ield Q/Ha (0.360), N non-reducing Sugar (0.337), and N number of fruits per plant (0.257). Conversely, T total sugar (-0.228) exhibited a highly negative and significant correlation with the trait, while all other traits were not significantly correlated with A average fruit weight.

In the first year, n Number of fruits per plant exhibited highly positive and significant correlations at the phenotypic level with T total fruit yield per plant (0.736), Y ield Q/Ha (0.348), N non-reducing sugar (0.217), and T total phenolic content (0.191). No data showed negative and significant correlations with the trait, and no traits were significantly correlated with the days to N number of fruits per plant. In the second year, N number of fruits per plant displayed highly positive and significant correlations at the phenotypic level with T total fruit yield per plant (0.701), yield Q/ha (0.700), and A ascorbic acid (0.179). Conversely, T total phenolic content showed a highly negative and significant correlation with the trait, while all other traits were not significantly correlated with the N number of fruits per plant. In pooled data, N number of fruits per plant showed highly positive and significant correlations at the phenotypic level with T total fruit yield per plant (0.723) and Y ield Q/Ha (0.540). No data showed highly negative and significant correlations with the trait, and all other traits were not significantly correlated with the N number of fruits per plant.

In the first year, T total phenolic content exhibited highly positive and significant correlations at the phenotypic level with N non-reducing sugar (0.438), T total sugar (0.330), T total fruit yield per plant (0.262), and D dry matter (0.186). No data showed negative and

significant correlations with the trait, and all other traits were not significantly correlated with the days to tTotal phenolic content. For all other traits, there were no significant correlations found with the Ddry matter content %.

In the first year, reducing sugar % displayed a highly positive and significant correlation at the phenotypic level with tTotal sugar (0.786), while TSS (-0.180) showed a negative and significant correlation with the trait. All other traits were not significantly correlated with the days to rReducing sugar %. In the second year, reducing sugar % exhibited a highly positive and significant correlation at the phenotypic level with tTotal sugar (0.805), while nNon-reducing sugar (-0.171) showed a highly negative and significant correlation with the trait. All other traits were not significantly correlated with the rReducing sugar %. In pooled data, reducing sugar % showed a highly positive and significant correlation at the phenotypic level with tTotal sugar (0.828), while TSS (-0.220) and nNon-reducing sugar (-0.176) showed highly negative and significant correlations with the trait. All other traits were not significantly correlated with the rReducing sugar %.

In Y1, the percentage of non-reducing sugar exhibited a highly positive and significant phenotypic correlation with total fruit yield per plant (0.326) and yield per hectare (0.180). No data demonstrated a negative and significant correlation with this trait. In addition Additionally, no other traits were significantly correlated with the days to non-reducing sugar percentage. In Y2, the non-reducing sugar percentage had a highly positive and significant phenotypic correlation with total sugar (0.352), with no data showing a highly negative and significant correlation with the trait. No other traits showed significant correlation with the non-reducing sugar percentage. When pooled, the non-reducing sugar percentage displayed a highly positive and significant phenotypic correlation with total sugar (0.268), with no data showing a highly negative and significant correlation. Again, no other traits were significantly correlated with the non-reducing sugar percentage.

In Y1, total sugar percentage was highly positive and significantly correlated at the phenotypic level with no positive combinations, while TSS (-0.229) showed a negative and significant correlation with the trait. No other traits were significantly correlated with the days to total sugar percentage. In Y2, total sugar percentage was highly positive and significantly

correlated at the phenotypic level with no positive combinations, with TSS (-0.179) showing a highly negative and significant correlation with the trait.

All other traits were not significantly correlated with the total sugar percentage. In the pooled data, the total sugar percentage showed a highly positive and significant phenotypic correlation with no positive combinations, while TSS (-0.238) exhibited a highly negative and significant correlation with the trait. No other traits were significantly correlated with the total sugar percentage.

In Y1, TSS ~~Brix~~ had a highly positive and significant phenotypic correlation with no positive combinations, whereas ascorbic acid (-0.260) showed a negative and significant correlation with the trait. No other traits were significantly correlated with the days to TSS ~~Brix~~. In Y2, TSS ~~Brix~~ had a highly positive and significant phenotypic correlation with no positive combinations, with ascorbic acid (-0.134) showing a negative and significant correlation. No other traits were significantly correlated with TSS ~~Brix~~.

When pooled, TSS ~~Brix~~ showed a highly positive and significant phenotypic correlation with no positive combinations, while ascorbic acid (-0.232) exhibited a highly negative and significant correlation with the trait. No other traits were significantly correlated with the reducing sugar percentage.

In Y1, ascorbic acid displayed a highly positive and significant phenotypic correlation with no combinations, with no combinations showing a negative and significant correlation. No other traits were significantly correlated with the days to ascorbic acid.

In Y2, ascorbic acid showed a highly positive and significant phenotypic correlation with no other combinations, and no combinations showed a highly negative and significant correlation with this trait. No other traits were significantly correlated with ascorbic acid. In the pooled data, ascorbic acid exhibited a highly positive and significant phenotypic correlation with no other combinations, and again, no combinations showed a highly negative and significant correlation. No other traits were significantly correlated with ascorbic acid.

In Y1, total fruit yield per plant demonstrated a highly positive and significant phenotypic correlation with yield per hectare (0.351), with no combinations showing a negative and

significant correlation. No other traits were significantly correlated with total fruit yield per plant. In Y2, total fruit yield per plant had a highly positive and significant phenotypic correlation with yield per hectare (0.752), with no combinations showing a highly negative and significant correlation. No other traits were significantly correlated with total fruit yield per plant. In the pooled data, total fruit yield per plant showed a highly positive and significant phenotypic correlation with yield per hectare (0.591), and no combinations showed a highly negative and significant correlation. No other traits were significantly correlated with total fruit yield per plant.

In Y1, yield per hectare (Yield Q/Ha) showed a highly positive and significant phenotypic correlation with no other combinations, and no combinations exhibited a negative and significant correlation with this trait. No other traits were significantly correlated with yield per hectare. In Y2, yield per hectare again displayed a highly positive and significant phenotypic correlation with no other combinations, and no combinations showed a highly negative and significant correlation with the trait. No other traits were significantly correlated with yield per hectare. In the pooled data, yield per hectare maintained a highly positive and significant phenotypic correlation with no other combinations, and no combinations showed a highly negative and significant correlation with the trait. No other traits were significantly correlated with yield per hectare. **These findings were earlier reported by- [10,11,12,13,14,15]**

Genotype Correlation Coefficient

In Y1, days to 50% flowering demonstrated a highly positive and significant genotypic correlation with days to first fruit harvest~~ing~~ (0.460) and reducing sugar (0.215). Conversely, it showed a negative and significant correlation with the number of primary branches per plant (-0.236), the number of fruits per cluster (-0.226), and yield per hectare (-0.214). No other traits were significantly correlated with days to 50% flowering.

In Y2, days to 50% flowering exhibited a highly positive and significant genotypic correlation with days to first fruit harvest~~ing~~ (0.695), reducing sugar (0.297), total sugar (0.233), ascorbic acid (0.188), and yield per hectare (0.177). It showed a highly negative and significant correlation with the number of primary branches per plant (-0.270). No other traits were significantly correlated with days to 50% flowering.

Comment [118]: Please discuss the characteristics with more up-to-date literature.

In the pooled data, days to 50% flowering had a highly positive and significant genotypic correlation with days to first fruit flowering (0.584) and reducing sugar (0.273). It showed a highly negative and significant correlation with the number of primary branches per plant (-0.315) and the number of fruits per cluster (-0.198). No other traits were significantly correlated with days to 50% flowering.

In Y1, days to first fruit harvesting showed a highly positive and significant genotypic correlation with pedicle length (0.256) and total fruit yield per plant (0.201). In contrast, plant height (-0.204) had a negative and significant correlation with this trait. No other traits were significantly correlated with days to first fruit harvesting.

In Y2, days to first fruit harvesting exhibited a highly positive and significant genotypic correlation with ascorbic acid (0.182). Conversely, it showed highly negative and significant correlations with the number of primary branches per plant (-0.417), the number of fruits per cluster (-0.357), and plant height (-0.324). No other traits were significantly correlated with days to first fruit harvesting.

In the pooled data, days to first fruit harvesting showed positive and significant genotypic correlations with pedicle length (0.149), ascorbic acid (0.138), number of fruits per plant (0.133), and total fruit yield per plant (0.132). On the other hand, it exhibited highly negative and significant correlations with the number of fruits per cluster (-0.418), the number of primary branches per plant (-0.349), and plant height (-0.260). No other traits were significantly correlated with days to first fruit harvesting.

In Y1, plant height showed highly positive and significant genotypic correlations with dry matter (0.330), reducing sugar (0.268), ascorbic acid (0.222), and total sugar (0.185). Conversely, it had negative and significant correlations with total fruit yield per plant (-0.395), the number of fruits per plant (-0.295), average fruit weight per plant (-0.236), and total phenolic content (-0.213). No other traits were significantly correlated with plant height.

In Y2, plant height displayed a highly positive and significant genotypic correlation with the number of primary branches (0.194). It showed highly negative and significant correlations with total phenolic content (-0.360), pedicle length (-0.199), and dry matter content (-0.187). No other traits were significantly correlated with plant height.

In the pooled data, plant height showed highly positive and significant genotypic correlations with ascorbic acid (0.160) and the number of fruits per cluster (0.155). Conversely,

it exhibited highly negative and significant correlations with total fruit yield per plant (-0.331), average fruit weight (-0.289), number of fruits per plant (-0.265), and pedicle length (-0.256). No other traits were significantly correlated with plant height.

In Y1, the number of primary branches per plant showed highly positive and significant genotypic correlations with average fruit weight (0.486), number of fruits per cluster (0.458), total fruit yield per plant (0.353), number of fruits per plant (0.313), pedicle length (0.297), and yield per hectare (0.251). On the other hand, dry matter content (-0.176) exhibited a negative and significant correlation with this trait. No other traits were significantly correlated with the number of primary branches.

In Y2, the number of primary branches showed a highly positive and significant genotypic correlation with TSS (0.182). Conversely, it had a highly negative and significant correlation with yield per hectare (-0.213). No other traits were significantly correlated with the number of primary branches.

In the pooled data, the number of primary branches showed highly positive and significant genotypic correlations with the number of fruits per cluster (0.268), pedicle length (0.185), and average fruit weight per plant (0.183). Conversely, yield per hectare (Yield Q/Ha) had a highly negative and significant correlation with this trait (-0.204). No other traits were significantly correlated with the number of primary branches.

In Y1, pedicle length exhibited highly positive and significant genotypic correlations with TSS (0.392), total fruit yield per plant (0.327), number of fruits per plant (0.291), and average fruit weight (0.269). In contrast, ascorbic acid (-0.181) showed a highly negative and significant correlation with pedicle length. No other traits were significantly correlated with pedicle length.

In Y2, pedicle length showed highly positive and significant genotypic correlations with the number of fruits per plant (0.414), yield per hectare (0.413), total fruit yield per plant (0.355), average fruit weight (0.340), TSS (0.314), and the number of fruits per cluster (0.184). On the other hand, total sugar (-0.235) and reducing sugar (-0.207) exhibited highly negative and significant correlations with pedicle length. No other traits were significantly correlated with pedicle length.

In the pooled data, pedicle length showed highly positive and significant genotypic correlations with TSS (0.345), the number of fruits per plant (0.275), total fruit yield per plant (0.262), and average fruit weight (0.207). Conversely, it exhibited a highly negative and

significant correlation with total sugar (-0.244). No other traits were significantly correlated with pedicle length.

In Y1, the number of fruits per cluster demonstrated highly positive and significant genotypic correlations with average fruit weight (0.369), yield per hectare (0.358), number of fruits per plant (0.297), non-reducing sugar (0.232), total sugar (0.203), and total phenolic content (0.176). There were no highly negative and significant correlations for this trait. No other traits were significantly correlated with the number of fruits per cluster.

In Y2, the number of fruits per cluster showed highly positive and significant genotypic correlations with dry matter content (0.341), average fruit weight (0.339), and total fruit yield per plant (0.221). There were no highly negative and significant correlations for this trait. No other traits were significantly correlated with the number of fruits per cluster.

In the pooled data, the number of fruits per cluster displayed highly positive and significant genotypic correlations with average fruit weight (0.328) and yield per hectare (0.190). There were no highly negative and significant correlations for this trait. No other traits were significantly correlated with the number of fruits per cluster.

In Y1, average fruit weight showed highly positive and significant genotypic correlations with total fruit yield per plant (0.569), yield per hectare (0.454), non-reducing sugar (0.366), number of fruits per plant (0.348), and total phenolic content (0.183). There were no highly negative and significant correlations for this trait. No other traits were significantly correlated with average fruit weight.

In Y2, average fruit weight displayed strong positive correlations at the genotypic level with Yield Q/Ha (0.478), Number of fruits per plant (0.445), Total fruit yield per plant (0.409), TSS (0.195), and Total sugar (0.180). No significant negative correlations were found for any trait. Other traits did not show significant correlations with average fruit weight.

In pooled data, average fruit weight showed significant positive correlations at the genotypic level with Total fruit yield per plant (0.535), Yield Q/Ha (0.435), non-reducing sugar (0.368), Number of fruits per plant (0.332), and Total phenolic content (0.200). There were no significant negative correlations found for any trait. All other traits did not show significant correlations with average fruit weight.

In Y1, Number of fruits per plant exhibited strong positive correlations at the genotypic level with Total fruit yield per plant (0.905), Yield Q/Ha (0.453), non-reducing sugar (0.237),

and tTotal phenolic content (0.202). No significant negative correlations were observed for any trait. Other traits did not show significant correlations with nNumber of fruits per plant.

In Y2, nNumber of fruits per plant showed strong positive correlations at the genotypic level with tTotal fruit yield per plant (0.793), yYield Q/Ha (0.787), and aAscorbic acid (0.210). There was a significant negative correlation (-0.298) found with nNumber of fruits per plant.

In pooled data, all other characteristics did not exhibit significant correlations with the nNumber of fruits per plant. Among these, nNumber of fruits per plant displayed notably strong positive correlations at the genotypic level with tTotal fruit yield per plant (0.859), yYield Q/Ha (0.618), and aAscorbic acid (0.183). There were no instances of highly negative and significant correlations with this trait. All other traits did not show significant correlations with the nNumber of fruits per plant.

In Y1, tTotal phenolic content demonstrated strong positive correlations at the genotypic level with total non-reducing sugar (0.460), total yield per plant (0.307), yYield Q/Ha (0.273), dDry matter content (0.255), and reducing sugar (0.198). No significant negative correlations were identified for this trait. All other traits did not exhibit significant correlations with tTotal phenolic content.

In Y2, total phenolic content exhibited strong positive and significant correlations at the genotypic level with total non-reducing sugar (0.545) and total sugar (0.406). Conversely, ascorbic acid (-0.153) and tYield Q/Ha (-0.137) showed negative and significant correlations with this trait. No other traits showed significant correlations with total phenolic content.

In pooled data, total phenolic content showed strong positive and significant correlations at the genotypic level with non-reducing sugar (0.538), total sugar (0.411), and reducing sugar (0.133). There were no instances of highly negative and significant correlations with this trait. No other traits were found to be significantly correlated with total phenolic content.

In Y1, dry matter content displayed strong positive and significant correlations at the genotypic level with total sugar (0.437), reducing sugar (0.350), and yield Q/Ha (0.185). No significant negative correlations were observed for this trait. All other traits did not show significant correlations with dry matter content.

In Y2, dry matter content showed strong positive and significant correlations at the genotypic level with aAscorbic acid (0.277) and TSS (0.208). Conversely, non-reducing sugar (-

0.163) exhibited a negative and significant correlation with this trait. No other traits were found to be significantly correlated with dry matter content.

In pooled data, d Dry matter content displayed strong positive and significant correlations at the genotypic level with reducing sugar (0.235), total sugar (0.231), and a Ascorbic acid (0.172). There were no instances of highly negative and significant correlations with this trait. All other traits did not show significant correlations with d Dry matter content.

In Y1, reducing sugar exhibited strong positive and significant correlations at the genotypic level with total sugar (0.802), y Yield Q/Ha (0.453), and a Ascorbic acid (0.185). However, TSS (-0.244) showed a strong negative and significant correlation with this trait. All other traits did not show significant correlations with r Reducing sugar.

In Y2, reducing sugar showed strong positive and significant correlations at the genotypic level with total sugar (0.821). However, TSS (-0.179) showed a strong negative and significant correlation with this trait. All other traits did not show significant correlations with r Reducing sugar.

In pooled data, reducing sugar demonstrated strong positive and significant correlations at the genotypic level with t Total sugar (0.835). However, TSS (-0.240) and non-reducing sugar (-0.176) exhibited strong negative and significant correlations with this trait. All other traits did not show significant correlations with r Reducing sugar.

In Y1, non-reducing sugar demonstrated strong positive and significant correlations at the genotypic level with total fruit yield per plant (0.341) and y Yield Q/Ha (0.263). No significant negative correlations were found for this trait. All other traits did not show significant correlations with non-reducing sugar.

In Y2, non-reducing sugar showed strong positive and significant correlations at the genotypic level with total sugar (0.367). No significant negative correlations were found for this trait. All other traits did not show significant correlations with non-reducing sugar.

In pooled data, non-reducing sugar exhibited strong positive and significant correlations at the genotypic level with t Total sugar (0.273). No significant negative correlations were found for this trait. All other traits did not show significant correlations with non-reducing sugar.

In Y1, t Total sugar displayed strong positive and significant correlations at the genotypic level with total fruit yield per plant. However, TSS (-0.275) showed a strong negative and significant correlation with this trait.

In Y1, t Total sugar did not show significant correlations with any other traits. However, TSS (-0.221) exhibited a strong negative and significant correlation with this trait. All other traits did not demonstrate significant correlations with t Total sugar.

In pooled data, t Total sugar did not show significant correlations with any other traits, but TSS (-0.273) displayed a strong negative and significant correlation with this trait. All other traits did not demonstrate significant correlations with t Total sugar.

In Y2, t Total sugar did not show significant correlations with any other traits. However, TSS ~~brix~~ (-0.221) exhibited a strong negative and significant correlation with this trait. All other traits did not demonstrate significant correlations with t Total sugar.

In Y1, TSS ~~brix~~ showed a strong positive and significant correlation at the genotypic level with Y yield Q/Ha (0.240). However, a Ascorbic acid (-0.275) showed a strong negative and significant correlation with this trait. All other traits did not demonstrate significant correlations with TSS ~~brix~~.

In Y2, TSS ~~brix~~ did not show significant correlations with any other traits. However, A a Ascorbic acid (-0.134) exhibited a negative and significant correlation with this trait. All other traits did not demonstrate significant correlations with TSS brix.

In pooled data, TSS ~~brix~~ exhibited a strong positive and significant correlation at the genotypic level with no positive associations. However, a Ascorbic acid (-0.262) showed a strong negative and significant correlation with this trait. All other traits did not show significant correlations with TSS ~~brix~~.

In Y1, a Ascorbic acid demonstrated a strong positive and significant correlation at the genotypic level with no positive relationships observed. There were no instances of highly negative and significant correlations with this trait. All other traits did not show significant correlations with a Ascorbic acid.

In Y2, a Ascorbic acid showed a strong positive and significant correlation at the genotypic level with no positive relationships identified. However, a Ascorbic acid (-0.155) exhibited a negative and significant correlation with this trait. All other traits did not show significant correlations with A a Ascorbic acid.

In pooled data, A a Ascorbic acid displayed a strong positive and significant correlation at the genotypic level with y Yield Q/Ha (0.222). There were no instances of highly negative and

significant correlations with this trait. All other traits did not show significant correlations with Ascorbic acid.

In Y1, Total fruit yield per plant demonstrated a strong positive and significant correlation at the genotypic level with Yield Q/Ha (0.570). There were no instances of highly negative and significant correlations with this trait. All other traits did not show significant correlations with Total fruit yield per plant.

In Y2, Total fruit yield per plant exhibited a strong positive and significant correlation at the genotypic level with Yield Q/Ha (0.883). There were no instances of a negative and significant correlation with this trait. All other traits did not show significant correlations with Total fruit yield per plant.

In pooled data, Total fruit yield per plant showed a strong positive and significant correlation at the genotypic level with Yield Q/Ha (0.759). There were no instances of a negative and significant correlation with this trait. All other traits did not show significant correlations with Total fruit yield per plant.

In Y1, Yield Q/Ha demonstrated a strong positive and significant correlation at the genotypic level with no positive relationships observed. However, there were no instances of a highly negative and significant correlation with this trait. All other traits did not show significant correlations with Yield Q/Ha.

In Y2, Yield Q/Ha showed a strong positive and significant correlation at the genotypic level with no positive relationships identified. However, there were no instances of a negative and significant correlation with this trait. All other traits did not show significant correlations with Yield Q/Ha. In pooled data, Yield Q/Ha exhibited a strong positive and significant correlation at the genotypic level with no positive relationships observed. However, there were no instances of a highly negative and significant correlation with this trait. All other traits did not show significant correlations with Yield Q/Ha. **These findings were earlier reported by [6,8,16,17,18,19,20,21,22]**

Comment [119]: There is a problem with this comment and it should be reviewed again.

Conclusion

Correlation coefficients were calculated between yield and its attributes at both genotypic and phenotypic levels to understand the relationships among these characteristics. This analysis provides insights into the extent and direction of selection pressure needed for practical

applications. Generally, genotypic correlations are emphasized due to the masking effect of genotypes on trait expression. In pooled data, Average fruit weight exhibited strong positive and significant correlations at the phenotypic level with Total fruit yield per plant (0.442), Yield Q/Ha (0.360), Non-reducing Sugar (0.337), and Number of fruits per plant (0.257). At the genotypic level across pooled data, Average fruit weight showed strong positive and significant correlations with Total fruits yield per plant (0.535), Yield Q/Ha (0.435), non-reducing sugar (0.368), Number of fruits per plant (0.332), and Total phenolic content (0.200). No significant negative correlations were observed for any trait. All other traits did not show significant correlations with Average fruit weight.

References

1. Aykroyd, W. R. (1963). The nutritional value of Indian foods and planning of satisfactory diet. I.C.M.R. Special Rep. Series. No. 42.
2. Bansal S, Mehta AK. Genotypic correlation and path analysis in brinjal (*Solanum melongena* L.). National J. of plant Improvement 10. 2008; 1:34-6.
3. Chen, N. C. and Li, H. M. (1996). Cultivation and Seed Production of Eggplant. Int.J. Curr.Microbiol. App.Sci (2017) 6(7): 4543-4548 AVRDC, Annual Publication, 203 (64): 246-261.
4. Choudhary, B. (1976). Vegetable, 4 thedn. National Book Trust, New Delhi, pp. 50-58.
5. Goto, K. (1953). Genetic studies on eggplant. Genetics, 2b:445-412

Comment [I20]: Please be organised by journal reference style

Comment [I21]: please be organised by reference style

6. Mishra SV, Warade SD, Nayakwadi MB. Correlation and path coefficient analysis in brinjal. J. of Maharashtra Agricultural Universities. 2007;32(1):74-6.
7. Muniappan S, Saravanan K, Ramya B. Studies on genetic divergence and variability for certain economic character in eggplant (*Solanum melongena* L.). Electron J Plant Breed. 2010;1(4):462-5.
8. Naliyadhara MV, Golani IJ, Mehta DR, Purohit VL. Genetic variability, correlation co-efficient and path analysis in brinjal. Orissa J Hortic. 2007;35(2):92-6.
9. Nayak, B. R. and Nagre, P. K. 2013. Genetic variability and correlation studies in brinjal (*Solanum melongena* L.). *Int. J Appl. Biol.*, **4**(4):211-215.
10. Pandey, P. K., Yadav, G. C. and Kumar, V. 2016. Correlation and path coefficient analysis among different characters in genotype of brinjal (*Solanum melongena*). *Indian J. Ecol.*, **43**(1):370-372.
11. Patel K. K. and Sarnaik, D. A. 2003. Performance study of long fruited genotypes of brinjal under Raipur conditions. *Orissa J Hort.*, **13**(1):74-77.
12. Patel KK, Sarnaik DA. Correlation and path coefficient analysis in brinjal (*Solanum melongena* L.). Haryana J Hortic Sci. 2004;33(3/4):246-7.
13. Prabhu, M. and Natarajan S. 2008. Correlation and path coefficient analysis in eggplant (*Solanum melongena* L.). *J Agric. Res.*, **42**(3):232-234.
14. Pratibha, Singh, Y.V. and Gupta, A.J. ProgHortic. Yield determinants in new hybrids of brinjal (*Solanum melongena* L.). 2004;36(2):290-2.
15. Savita Darpan, Savita Darpan, Khiromani Nag Khiromani Nag, and Akhilesh Jagre Akhilesh Jagre. "Study the correlation, path coefficient between yield and its attributing characters in brinjal (*Solanum melongena* L.)." (2018): 1411-1415.
16. Sharma, T.V.R.S.; Kishan, S. and Swaroop, K. (2000). Genetic variability and character association in brinjal (*Solanum melongena* L.). *Indian. J. Hort.*, **57** (1): 59-65
17. Singh HV, Singh SP, Singh S, Rajput CBS. Heterosis in relation to combining ability in brinjal (*Solanum melongena* L.). *Veg Sci.* 2003;30(1):38-41.

Comment [I22]: Please be organised by reference style

Comment [I23]: please be organised by reference style

Comment [I24]: please be organised by reference style

Comment [I25]: please be organised by reference style

Formatted: Font: 11 pt, Font color: Auto, English (India), Pattern: Clear

Comment [I26]: please be organised by reference style

18. Singh, A. K., Tripathi, M. K., Rai, V. K. and Mishra, R. 2011. Character association and path coefficient analysis in brinjal (*Solanum melongena* L.). *Environ & Ecol.*, **29**: 1201-1203.

Comment [I27]: please be organised by reference style

19. Singh, N., Singh, D. K., Pandey, P., Panchbhaiya, A. and Rawat, M. 2017. Correlation and Path Coefficient Studies in Okra [*Abelmoschus esculentus* (L.) Moench]. *Int. J. Curr. Microbiol. App. Sci.*, **6**(7), 1096-1101. [Cross Ref]

Comment [I28]: please be organised by reference style

20. Sujin, G.S., Karuppaiah, P. and Saravanan, K., 2017. Genetic variability and correlation studies in brinjal (*Solanum melongena* L.). *Indian Journal of Agricultural Research*, **51**(2), pp.112-119.

Comment [I29]: please be organised by reference style

Formatted: Font: 11 pt, Font color: Auto, English (India), Pattern: Clear

21. Thangamani C, Jansirani P. Correlation and path analysis studies on yield attributing characters in brinjal (*Solanum melongena* L.). *Electron Plant Breed.* 2012;3(3):939-44.

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, English (India)

22. Thangamani, C. and Jansirani, P. 2012. Correlation and path coefficient analysis studies on yield and attributing characters in brinjal (*Solanum melongena* L.). *Electronic Journal of Plant Breeding*, **3**(3): 939-944

Formatted: Font: 11 pt, Font color: Auto, English (India), Pattern: Clear

Comment [I30]: please be organised by reference style

23. Zakari, S.M, Haruna, H. and Aliko, AA 2017 Correlation analysis of bulb yield with growth and yield components of garlic (*Allium sativum* L.) *Nigerian Journal of Basic and Applied Science*, **25** 58-62.

Comment [I31]: please be organised by reference style

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, English (India)

Formatted: Font: 11 pt, Font color: Auto, English (India)

UNDER PEER REVIEW