

Role of fermented organic manure and mycorrhiza on physio-chemical properties of soil of Prayagraj

ABSTRACT

The integrated use of fermented organic manure and mycorrhiza has been employed to enhance crop productivity and maintain soil health and fertility. The research titled “Role of fermented organic manure and mycorrhiza on physio-chemical properties of soil” was conducted during the Zaid season of 2023 at the Central Research Farm, Department of Soil Science and Agricultural Chemistry, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology, and Sciences, Prayagraj, Uttar Pradesh, India. The plot was designed in 3X3 RBD having 3 levels of fermented organic manure @ 0, 50, 100 %, Mycorrhiza @ 0, 50, 100 % and N, P, K at RDF respectively. The Variety selected was PDM-139 released by SVRC, Uttar Pradesh in the year 1974. The average yield recorded was 6-9 q ha⁻¹. Results indicated non-significant with the highest values observed in the control treatment in case of bulk density (Mg m⁻³) and particle density (Mg m⁻³) of soil was recorded maximum in T₁ (Absolute control) and minimum was recorded at T₉ (NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 100%). The maximum Pore space (%) and Water holding capacity (%) of soil was recorded at T₉ (NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 100%) and the minimum was found in T₁ (control). The slight decrease of pH and EC was observed on application of the treatments, the maximum pH of soil was recorded at T₉ (NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 100%) and the minimum in T₁ (Absolute control). The maximum organic carbon (%) and increase of N, P, K of soil was recorded at T₉ (NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 100%) and the minimum was found in T₁ (Absolute control).

Keywords: *Fermented organic manures, Mycorrhiza, Green gram, Soil health, Physio-chemical properties, etc.*

INTRODUCTION

Soil is generally defined as the fine earth that covers the land surface, formed through various weathering processes either in place or by the accumulation of materials transported by wind, water, or ice. A distinctive feature of soil is the addition of organic material to this weathered substance. This organic matter can be both living and dead. Therefore, soil can be described as a mixture of mineral material and organic matter, which imparts its unique characteristics (**Nortcilff *et al.*, 2006**). Agricultural soil is a type of soil that is used for productive activities of various crops, considered of great importance in terms of environmental decontamination since plants apart from absorbing nutrients such as CO₂, also absorb different polluting compounds from soil[32,33,34].

Mycorrhiza are among the most widespread forms of symbiosis on Earth, involving a mutualistic relationship between plants and fungi that enhances nutrient and water uptake by plants (Johnson *et al.*, 2017). Operating at the interface of biological, chemical, and physical processes that underpin soil fertility, mycorrhizal fungi thrive in heterogeneous environments with highly variable conditions in soil and plant roots. The persistent use of chemical fertilizers can render soil unfit for crop cultivation by diminishing microbial activity and reducing microbial populations. In contrast, organic fertilizers play a crucial role in revitalizing soil health. Among these, biofertilizers are significant for maintaining soil health, with Arbuscular Mycorrhizal (AM) fungi forming associations with about 80% of plant species. The symbiotic relationship between AM fungi and plants has been extensively studied by researchers (**Rajashekar and Nagarajan, 2011**).

AM fungi, also known as Vesicular Arbuscular Mycorrhizal (VAM) fungi, are particularly effective at enhancing the absorption of micronutrients like zinc (Zn) and copper (Cu) and increasing plant resilience to various stresses, including drought, salinity, heavy metals, and water scarcity (**Arumugam *et al.*, 2010**). This symbiotic relationship boosts plant development by supplying essential nutrients such as nitrogen (N), phosphorus (P), potassium (K), and zinc. Mycorrhizal associations also help reduce plant susceptibility to certain soil-borne diseases, and the application of AM fungi has been shown to enhance nitrogenase activity (**Kobae *et al.*, 2019**).

Fermented organic manure over the past century, the evolution of agriculture has significantly increased agricultural output, with chemical fertilization playing a pivotal role. However, the increased use of fertilizers and intensive land use have led to numerous environmental issues, such as nitrogen runoff contributing to watershed pollution. Fermented organic manure has garnered attention for its environmental benefits. This type of microbial organic fertilizer not only supplies crops with essential nutrients but also

enhances crop growth conditions and activates intrinsic soil nutrients through numerous enzymatic activities (**Wang *et al.*, 2018**).

The composition of N-Carb, a type of fermented organic manure, includes a fermented organic extract that provides abundant organic nitrogen, phosphorus, potassium, and micronutrients. This enriches soil organic matter by rapidly enhancing organic carbon in the soil structure. Consequently, soil is protected from degradation and erosion, and soil aeration is improved. This environment promotes the activity of beneficial carbon-fixing microbes, which enrich soil organic carbon and support adequate photosynthesis in plants. The organic extracts of hydrolyzed material in N-Carb enable crop plants to absorb essential nutrients through their roots, even under adverse soil conditions. It also optimizes soil moisture retention at the root zone, helping plants withstand various biotic and abiotic stresses, including salinity, drought, and nutrient deficiency, making it suitable for both organic and conventional farming (**Dutta and Tambat, 2022**).

MATERIALS AND METHODS

The field trial was conducted during the Zaid season of 2023 at the Central Research Farm of the Department of Soil Science and Agricultural Chemistry at Sam Higginbottom University of Agriculture, Technology and Sciences (U.P.), located at 25°24'30'' N latitude, 81°51'10'' E longitude, and 98m above mean sea level. This location is part of the Agro-Ecological Sub Region [North Alluvium Plain Zone (0-1% slope)] and the Agro-Climatic Zone (Upper Gangetic Plain Region). Agro-climatically, Prayagraj District falls within the subtropical belt of Southeast Uttar Pradesh, experiencing extremely hot summers and cold winters. The maximum temperature ranges up to 46°C, rarely dropping below 4- 5°C, with relative humidity between 20-94%. The area receives an average annual rainfall of about 1100mm, primarily from July to the end of September, though occasional winter precipitation is not uncommon. During the crop season, temperatures ranged from a minimum of 21.38°C to a maximum of 37.82°C, while humidity varied from 46.42% to 96.85%

Experimental design

The experimental design was a 3x3 Randomized Block Design with three levels of fermented organic manure (0, 50, 100%), mycorrhiza (0, 50, 100%), and NPK at recommended doses (RDF). Each plot measured 2m x 2m, and the green gram variety used was PDM-139, aimed at evaluating soil properties and health.

Methodology

The soil in the experimental area was primarily sandy loam, classified as Inceptisol and part of the alluvium soil near the Ganga-Yamuna Basin. Before tillage operations, soil samples were collected randomly from five distinct locations within the experimental plot at a depth of 0-15cm. These samples were prepared for physical and chemical analysis by air-drying, coning, and quartering, sieving through a 2mm sieve, and storing in polythene bags. Post-harvest, soil samples were again collected according to different treatment combinations and analyzed for various parameters listed in table 1.

Table 1. Soil Parameter Analysis

I. Physical Properties

Sl. no.	Parameter	Method	Method
1.	Bulk Density	Graduated measuring cylinder	Muthuval <i>et al.</i> (1992)
2.	Particle density	Graduated measuring cylinder	Muthuval <i>et al.</i> (1992)
3.	Pore space	Graduated measuring cylinder	Muthuval <i>et al.</i> (1965)
4.	Water retention capacity	Graduated measuring cylinder	Muthuval <i>et al.</i> (1992)

II. Chemical properties

Sl. no.	Parameter	Unit	Method
1.	Soil pH	Digital pH meter	Jackson,1958
2.	Soil EC	Digital EC meter	Wilcox, 1950
3.	Organic Carbon	Wet Oxidation Method	Walkley and Black, 1947
4.	Available Nitrogen	Kjeldahl method	Subbiah and Asija, 1956
5.	Available Phosphorus	Colorimetric method	Olsen <i>et al.</i> , 1954

6.	Available Potassium	Flame Photometer method	Toth and Prince, 1949
----	---------------------	-------------------------------	-----------------------

The detailed about treatment combination of the conducted research wise given in Table 2.

Table 2. Treatment combination

Treatments	Treatment Combination	Symbol
T₁	Absolute control	L₀M₀F₀
T₂	NPK@RDF + M₀ + F₁ @ 50%	LM₀F₁
T₃	NPK@RDF + M₀ + F₂ @ 100%	LM₀F₂
T₄	NPK@RDF + M₁@ 50% + F₀	LM₁F₀
T₅	NPK@RDF + M₁@ 50% + F₁@ 50%	LM₁F₁
T₆	NPK@RDF + M₁@ 50% + F₂@ 100%	LM₁F₂
T₇	NPK@RDF + M₂@ 100% + F₀	LM₂F₀
T₈	NPK@RDF + M₂@ 100% + F₁@ 50%	LM₂F₁
T₉	NPK@RDF + M₂@ 100% + F₂@ 100%	LM₂F₂

Note: RDF = Recommended dose of fertilizer, M= Mycorrhiza and F= Fermented organic manure.

RESULT AND DISCUSSIONS

The study found that using fermented organic manure and mycorrhiza significantly enhances various soil physical parameters, such as particle density, water holding capacity, and porosity, thereby maintaining good soil health. Additionally, soil chemical properties improved with increased nutrient levels, including soil electrical conductivity (EC), organic carbon, available nitrogen, phosphorus, and potassium.

The field experiment revealed that soil bulk density and pH decreased with higher levels of fermented organic manure and mycorrhiza. The maximum bulk density (1.341 Mg m⁻³) and pH (7.06) were observed in the control treatment T₁[Absolute control], while the minimum pH (6.95) and bulk density (1.224 Mg m⁻³) were found in the T₉. [NPK@RDF + M₂@ 10Kg ha⁻¹ 100% + F₂@ 50 l ha⁻¹ 100%].

The T₉. [NPK@RDF + M₂@ 10Kg ha⁻¹ 100% + F₂@ 50 l ha⁻¹ 100%] treatment also resulted in the highest values for electrical conductivity (0.42 dS m⁻¹), particle density (2.41), pore space (48.72 %), water holding capacity (48.53 %), organic carbon (0.65 %), available nitrogen (297.67 kg ha⁻¹), available phosphorus (26.03 kg ha⁻¹), available potassium (211.07 kg ha⁻¹). In contrast, the control treatment T₁[Absolute control] had the lowest values for these parameters: minimum electrical conductivity (0.30 dS m⁻¹), pore space (45.43 %), water holding capacity (43.42 %), organic carbon (0.45 %), available nitrogen (274.67 kg ha⁻¹), available phosphorus (18.83 kg ha⁻¹), available potassium (178.30 kg ha⁻¹). This indicates that the T₉ treatment has the greatest potential to improve soil health and productivity.

Table 2. Effect of different levels of NPK fermented organic manure and mycorrhizal inoculation on bulk density (Mg m⁻³), particle density (Mg m⁻³) water holding capacity (%) and pore space (%).

Treatments	Bulk Density	Particle Density	Water Holding	Pore Space
	(Mg m ⁻³) 0-15 cm	(Mg m ⁻³) 0-15 cm	Capacity (%) 0-15 cm	(%) 0-15 cm
T ₁	1.341	2.424	43.52	44.68
T ₂	1.334	2.434	44.79	45.20
T ₃	1.325	2.446	45.20	45.83
T ₄	1.294	2.449	45.47	47.17
T ₅	1.281	2.490	46.12	48.56
T ₆	1.271	2.513	46.56	49.43
T ₇	1.261	2.514	47.52	49.85
T ₈	1.244	2.564	47.95	51.49
T ₉	1.224	2.597	48.53	52.87
F-Test	NS	NS	S	S
S.Em. (±)	0.02	0.02	0.21	0.14
C.D.@5%	0.05	0.05	0.45	0.29

Table 3. Effect of different levels of NPK fermented organic manure and mycorrhizal inoculation on pH, EC, organic carbon, Av. Nitrogen, Av. Phosphorus and Av. Potassium (kg ha⁻¹)

Treatments	pH	EC (dSm ⁻¹)	OC (%)	Av. Nitrogen (kg ha ⁻¹)	Av. P ₂ O ₅ (kg ha ⁻¹)	Av. K ₂ O (kg ha ⁻¹)
	0-15 cm	0-15 cm	0-15 cm	0-15 cm	0-15 cm	0-15 cm
T ₁	7.06	0.30	0.45	274.67	18.83	178.30
T ₂	7.05	0.32	0.46	276.33	19.60	181.52
T ₃	7.04	0.34	0.51	279.33	20.88	187.12
T ₄	7.05	0.34	0.53	283.33	21.48	192.18
T ₅	7.03	0.36	0.58	286.00	22.60	191.87
T ₆	7.01	0.38	0.60	289.33	23.56	193.91
T ₇	7.02	0.40	0.61	292.67	24.60	202.80
T ₈	6.99	0.41	0.63	296.33	25.08	207.68
T ₉	6.95	0.42	0.65	297.67	26.03	211.65
F-Test	S	S	S	S	S	S
S.Em. (±)	0.08	0.02	0.03	0.54	0.16	0.85
C.D.@5%	0.21	0.03	0.05	1.15	0.35	1.80




Fig 1. Graphical Representation of Av. Nitrogen, Av. Phosphorus and Av. Potassium vs Treatment Combination




Fig 2. Graphical representation of treatment combination vs EC, pH and OC (%)

Conclusion

It is revealed from trial that the various level of NPK fermented organic manure and mycorrhizal inoculation used from in the experiment, the treatment combination T₉ – [NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 100%] for green gram it could be recommended for profitable production of green gram (*Vigna radiata* L.) var. PDM-139 and treatment T₉ is best in terms of soil physical and chemical properties. Further it increased soil fertility and nutrient uptake by the plant. The major challenges faced during the research trial was infestation of white flies and yellow mosaic virus attack due to dry condition around the field. Overall, it proved to be beneficial both respect to soil health properties by application of fermented organic manures and inoculation of mycorrhiza in cultivation and growth of green gram.

ACKNOWLEDGEMENT

It gives me immense pleasure, a sense of thanks and gratitude to be the student of my respected advisor **Dr. Arun Alfred David**, Assistant Professor, Department of Soil Science and Agricultural Chemistry. I consider myself lucky to work under such a great person whose moral support and valuable guidance despite his hectic schedule have kept in this stead. I can't sufficiently thank for his thoughts provoking suggestions, for teaching me very important principles and values of life and for the constant encouragement he bestowed on me as no less than a parent during course of two years and especially in thesis work.

REFERENCE

1. **Arumugam, T., Rajsekharan, S. and Nagarajan, S. M. (2011)** Response of arbuscular mycorrhizal fungi and Rhizobium inoculation on growth chlorophyll content of *Vigna unguiculata* L. walp var Pusa 151. *Journal of Applied Sciences and Environmental management* 14.
2. **Athar, Mohammed (2015)** Nodulation of Native legumes in Pakistani Rangelands. *Agriculture Conspectus Scientificus (ACS)* 70.
3. **Anonymous (2009)** Annual report for 2008. IRRI, Los Banos, Philippines.
4. **Black, C.A. (ed) (1965)** Methods of Soil Analysis Vol. 2 American Society, Agro. Madison, Wisconsin, U.S.A.
5. **Bouyoucos, G.J. (1927)** The Hydrometer and new method for mechanical analysis of Soils. *Soil Science*, 23:393- 395.
6. **Brar, T.S., Thalkar, M.G., Singh, K., and Singh, R. (2019)** Effect of different phosphorus level and arbuscular mycorrhizal fungi on yield of green gram (*Vigna radiata* L.) crop. *Plant archives*, vol 19(2); 3165-3168.
7. **Chalk, P., Souza, R., Alves, B. & Boddery, R. (2006)** The role of arbuscular mycorrhiza in Legumes symbiotic performance, *Soil Biology and Biochemistry*, 38. 2944-2951.
8. **Datta, S.K. and Tambat, H. (2022)** The role of N-Carb, a fermented organic manure in reclamation of soil having low organic carbon, *International Journal of Scientific Research Publication*, Volume 12,3.
9. **Imran, R.A., Shahib, Y.J. and Khyuon, A. (2020)** use of three types of fermented and non-fermented organic fertilizers in the growth and production of lettuce (*Lactuca sativa* L.). *International Journal of Statistical Sciences.*, 16(1); 1681-1688
10. **Jackson, M.L., (1958)** Soil Chemical Analysis Prentice Hall of India, Private Ltd. New Delhi (1973)
11. **Jaiswal, P.C. (2006)** Soil Plant & Water analysis, Chemical analysis. pp. 74-82.
12. **Jat, P. C., Rathore, S. S. and Sharma, R. K. (2014)** Effect of integrated nitrogen management and intercropping systems on yield attributes and yield of maize. *Indian Journal of Hill Farming*, **27** (1): 91-99.
13. **Johnson, N.C., and Jansa, J. (2017)** Mycorrhizas: At the interface of biological, Soil and Earth Sciences. Mycorrhizal mediation in soil, fertility structure and carbon storage. *Phytoplasma disease in Asian countries*, 2: 1-6.

14. **Kobae, Y. (2019)** Dynamic phosphate uptake in Arbuscular Mycorrhizal Fungi roots under field condition.
15. **Lia, A., Raden, B., Elly, R.R., Raden, W.W. and Upit, K. (2020)** Fermented compos and N-fertilizer for enhancing the growth and productivity of purple egg- plant on vertisols. *Open Agriculture*,5; 898-904.
16. **Moebius-Clune, B.N., Moebius-Clune, D.J., and Abawi, G.S., (2016)** Comprehensive assessment soil
a. `health: The Cornell framework manual. (third ed.), Cornell University, Geneva.
17. **Muhr, G.R., N.P. Datta, H. Sankarasubramoney, V.K. and Donahue, R.L., (1965)** Soil testing in India:
a. U.S. agency for Indian, Development mission to India. New Delhi.
18. **Munsell, A.H., ed. 12. (1971)** *Color Notation* Baltimore, MD Munsell Color. Company. PP-65
19. **Muthuvel P., Udaysoorian, C., Natesan, R., and Ramaswami, P.P. (1992)** Introduction to Soil analysis, Tamil Nadu Agricultural, Coimbatore- 641002
20. **Natasha L. Hoover, Ji Yeow Law, Leigh Ann M. Long, Ramesh S. Kanwar and Michelle L. Soupir, (2019)** Long-term impact of poultry manure on crop yield, soil and water quality, and crop revenue. *Journal of Environmental Management*, 252: 109582
21. **Nortcliff, S., Hulpke, H., Bannick, C.G., and Terytze, K. (2006)** Soil, definition, Function and utilisation of soil. Ullman's Encyclopedia of Industrial Chemistry. Land utilisation and soil destruction
22. **Olsen, S.R., Watanale, F. S., (1954)** A method to determine a phosphorus adsorption measured by colorimetric method. *Soil Science Society of America Journal*. 21,144-149
23. **Rajasekaran, S., and Nagarajan, S.M. (2005)** Effect of dual inoculation (AM fungi and Rhizobium) on chlorophyll content on *Vigna unguiculata* L. var PUSA-151. *Mycorrhiza News* 17: 10-11.
24. **Rathore, G., Kaushal, R., Sharma, V., Sharma, G., Chaudhary, S. and Dhaliwal, S.S. (2023)** Evaluation of the Usefulness of Fermented Liquid Organic Formulations and Manures for Improving the Soil Fertility and Productivity *Agriculture* 2023, 13, 417.
25. **Ray, J.G., and Valsalakumar, N. (2010).** Arbuscular Mycorrhizal fungi and Piriformospora indica individually and in combination with rhizobium on green gram. *Journal of Plant Nutrition*, 33(2): 285-298.
26. **Sharma, K. L., Srinivas, K., Mandal, U. K., Vittal, K. P. R., Grace, K. J. and Maruthi, S.G.R. (2014)** Integrated Nutrient Management HT Strategies for Sorghum and Green gram in Semi-Arid Tropical Alfisol. *Indian Journal of Dryland Agricultural Research and Development*. 2017 19(1), 13-23.

27. **Shraddha, A., Shukla, Y.R., Thakur, K. and Kumari, J. (2023)** Impact of Fermented Organic formulations combined with inorganic fertilizers on broccoli (*Brassica oleraceae* var. *italica*) cv. *Palam Samridhhi*. *Science Direct Heliyon*,9(2); 2031.
28. **Singh, R.P., Bisen. J.S., Yadav, P.K., Singh, S.N., Singh, R. K. and Singh, J. (2008)** Integrated use of sulphur and molybdenum on growth, yield and quality of black gram. *Legume Research* 31:214-217
29. **Subbiah. B.V. and Asija (1956)** A rapid procedure for estimation of available nitrogen in soil. *a. Current Science*. 25:259.
30. **Toth, S. J. and Prince, A. L. (1949)** Estimation of Cation exchange capacity and exchangeable Ca, K & Na content of soil by Flame Photometer technique. *Soil science journal*, 67, 439-445.
31. **Vikram, A. and Hamzehzarghani, H. (2008)** Effect of Phosphate Solubilizing Bacteria on Nodulation and Growth Parameters of Green gram (*Vigna radiata* L. Wilczek), *Research Journal of Microbiology*, 3: 62-72.
32. Aechra S, Yadav BL, Doodhwal K, Bhinda R, Jat L. Yield and Total Nutrient Uptake Influenced by Soil Salinity, Phosphorus Sources and Biofertilizers in Cowpea (*Vigna unguiculata* L.). *J. Exp. Agric. Int.* [Internet]. 2021 Jun. 14 [cited 2024 Jun. 2];43(4):56-63. Available from: <https://journaljeai.com/index.php/JEAI/article/view/1801>
33. Wang-Bara B, Kaouvon P, Housseini JD, Alioum SP, Danra Djackba D. Effects of Fertilization Based on Chicken Manures and Mycorrhiza on Vegetative Parameters and Phenological Stages of Sorghum bicolor in Yagoua, Far-North Cameroon. *Int. J. Plant Soil Sci.* [Internet]. 2021 Dec. 21 [cited 2024 Jun. 2];33(24):375-83. Available from: <https://journalijpss.com/index.php/IJPSS/article/view/1593>
34. Aguilar R, Carreón-Abud Y, López-Carmona D, Larsen J. Organic fertilizers alter the composition of pathogens and arbuscular mycorrhizal fungi in maize roots. *Journal of Phytopathology*. 2017 Aug;165(7-8):448-54.