

The Role of Alcohol Consumption in Cardiovascular Health: A Systematic Review and Meta-Analysis

Abstract

There remains a debate on the effects of alcohol use on cardiovascular health. This systematic review and meta-analysis examined the relationship between this link and blood pressure, lipid profiles, and the frequency of cardiovascular events to shed light on it. A comprehensive search strategy using PRISMA guidelines resulted in nine papers that met the inclusion requirements. Moderate alcohol usage was shown to have a U- or J-shaped connection with cardiovascular events; excessive intake or abstinence was associated with increased risk, but moderate consumption had preventive advantages. Moderate drinkers showed positive changes in their lipid profiles and lowered blood pressure in comparison to heavy or non-drinkers. The wide range of research highlights the need to consider individual characteristics and study techniques. Despite the potential cardiovascular benefits of moderate alcohol use shown by these studies, treatment recommendations should be cautious. To provide personalized recommendations to patients and policymakers, future research should focus on explaining mechanisms, examining modifiers, and assessing the effects of different types of alcohol and drinking habits on cardiovascular health.

Key words: Alcohol Consumption, Cardiovascular Health, Mental health, Patients' quality of life

Introduction

Cardiovascular disease (CVD) is the leading cause of premature mortality, presumed to contribute the highest number of lost life years on a global scale [1]. In fact, deaths from CVD outnumber in aggregate all of the other most significant causes of death (cancer, infectious diseases & eating disorders combined) by a rate nearly double. CVD is the leading cause of morbidity and also significantly affects patients' quality of life [2,3]. The most common forms of CVD are coronary heart disease (CHD) or cerebrovascular disease.

A main risk factor for CVD, is alcohol intake. Alcohol use is associated with more than 200 diseases and injuries and nearly 3 million deaths each year. The relationship between alcohol consumption and cardiovascular events also follows a well-known U- or J-shaped curve [4,5]. What seems clear, though, is that moderate alcohol intake has a protective effect with respect to CHD probably through mechanisms like elevation of HDL cholesterol and an inhibition of early stage of atherosclerosis. Heavy drinking raises the risk of cardiovascular events [6]. The range that offered the most risk was between 38 g per day or 23 drinks per week. Moreover, research indicates that whereas heavy drinking is connected to an exponential rise in illness, moderate drinking is associated with a decreased risk of cardiovascular disease [7].

Academics disagree on the hypothesis that alcohol use might protect against a number of diseases. From a pharmacological point of view, alcohol use interacts with several medications, including diuretics, antidepressants, and opioids [8], since it can change the way that pharmaceuticals and/or alcohol are metabolized [9]. Lower alcohol metabolism can result in greater blood alcohol levels, and this can be caused by histamine H₂ receptors and other drugs used to treat heartburn and ulcers [9]. It is necessary to use caution while taking resveratrol with some medicines, such as those found in wine, as alcohol might alter a drug's metabolism [10].

Alcohol consumption is particularly detrimental to mental health since it increases the risk of suicide when used acutely and in large doses [11]. In conclusion, from the standpoint of a cardiologist, studies conducted over the years have suggested that wine and mild to moderate alcohol use may help reduce the risk of cardiovascular disease [12, 27-30]. Recently, studies that have used Mendelian randomization approaches have questioned this impact. Reduced alcohol consumption is beneficial for cardiovascular health, according to the genetic approach analyses of these studies, which revealed that people with the alcohol dehydrogenase 1B (ADH1B) gene had a decreased risk of coronary heart disease when they drank less alcohol [13]. However, research has not been broken down into categories of alcoholic beverages [14].

Reduced risk of breast cancer is observed in women who consume up to one drink per day [14], and an inverse relationship between an acute intake of alcohol with ischemic heart disease incidence as well as the prognosis for patients under risk of future coronary episodes has been reported from several studies [15]. For example, consuming anything between 0 to 7.5 drinks per week, or less than 12.5 grams per day is the least harmful for health, as against the greater risk associated with a consumption over 38 grams per day or more than 23 drinks in a week [7]. Nevertheless, the favorable effects of alcohol on heart health must be balanced with its risks; heavy drinking is associated with an increasingly raised risk for both CVD and other health problems.

Even more concerning, the interplay between alcohol and many medications such as diuretics, antidepressants, and opioids complicates treatment plans. Alcohols can for example induce other substances like drugs to metabolize into the system after which it can also form certain toxic effect [8,9]. So, it is better to drink alcohol while taking the medicine.

While a spate of genetic investigations adopting Mendelian randomization suggest that moderation in alcohol drinking benefits cardiovascular health. Those with a gene (alcohol dehydrogenase 1B [ADH1B]) that metabolizes alcohol more rapidly and therefore typically drink less are also at lower risk of coronary heart disease [13].

Objective of the study

The purpose of this systematic review and meta-analysis is to investigate the impact of alcohol consumption on cardiovascular health, with a focus on its effects on blood pressure, lipid profile, and the incidence of cardiovascular events. By collecting data from relevant studies, this study aims to provide a comprehensive understanding of the relationship between alcohol intake and cardiovascular outcomes. The purpose of this study is to clarify the complex association between alcohol use and cardiovascular disease by using a rigorous methodology and following PRISMA guidelines. The results offer insightful information that might impact public health campaigns and clinical practice.

This study aims to address several key questions:

1. What is the relationship between alcohol consumption and the incidence of cardiovascular events?
2. How does alcohol consumption influence blood pressure levels, including systolic and diastolic measurements?
3. What effects does alcohol consumption have on lipid profile, including HDL and LDL cholesterol levels?
4. Are there significant differences in cardiovascular outcomes between moderate alcohol consumers, heavy drinkers, and abstainers?
5. What are the potential mechanisms underlying the observed associations between alcohol consumption and cardiovascular health?
6. What are the clinical implications of these findings for healthcare practitioners and policymakers?

Materials & Methods

The "Reporting Items for Systematic Review and Meta-Analysis (PRISMA)" guidelines were followed for conducting a recent systematic review [14].

Search strategy

The research papers related to the study's aims "The Role of Alcohol Consumption in Cardiovascular Health" were extracted. Four electronic databases such as PubMed, EMBASE, CINHAL, and Cochrane Library were used for data extraction. The timeline of research was set from January 2004 to January 2024. To reach authentic data, the MeSH keywords were used such as (moderate alcohol consumption [mh]) OR (Average alcohol consumption) OR (heavy alcohol consumption) OR (non-alcohol users OR (never users) OR (non-alcoholic) AND ((incidence of CVD [mh]) OR (Hypertension) OR (blood pressure) OR (HDL and LDL)).

List 1: List of Eligibility Criteria

Criteria	Inclusion	Exclusion
Population	Adults aged 18 and older	Individuals under 18 years of age
Condition	Studies focusing on cardiovascular disease and alcohol consumption	Studies focusing on non-cardiovascular diseases or unrelated to alcohol consumption
Study Design	Randomized controlled trials, cohort studies, case-control studies, and cross-sectional studies	Case reports, editorials, letters, and non-peer-reviewed articles
Language	Studies published in English	Studies published in languages other than English
Publication Date	Studies published from January 2004 to January 2024	Studies published before January 2004
Outcome Measures	Studies reporting on cardiovascular events, blood pressure, lipid profile, and mortality	Studies not reporting relevant cardiovascular outcomes
Alcohol Consumption	Studies that quantify alcohol intake and categorize it into different levels	Studies that do not quantify alcohol intake

Intervention/Exposure	Alcohol consumption (any form)	Interventions focusing on substances other than alcohol
Comparators	Comparisons between different levels of alcohol consumption (e.g., none, moderate, heavy)	Studies without a clear comparator group related to alcohol consumption

Data Extraction

For analysis, we extracted the information related to authors, year of study, country, study population, sample size, type of alcoholic consumption pattern, study design, and primary outcomes such as incidence of CVD, blood pressure, and lipid profile (HDL, & LDL) from selected articles after the selection and screening of research articles.

Primary Outcomes

The primary outcomes of the recent meta-analysis were the incidence of CVD, blood pressure (systolic and diastolic), and lipid profile (LDL and HDL). Hypertension is systolic blood pressure (BP) ≥ 140 mmHg and/or diastolic BP ≥ 90 mmHg, and/or the use of antihypertensive medication.

Risk of Bias Assessment

To evaluate the risk of bias in included studies, the Cochrane risk of bias assessment tool was used [16]. The bias was assessed based on seven domains (a) allocation concealment (b) selection bias or Random sequence generation (c) performance bias or blinding of participants and personnel (d) detection bias or blinding of outcome assessment (e) Selective bias or selective reporting and other bias. Each domain's score was categorized into Low risk, high risk, or unclear. For cohort studies, the Joanna Briggs Institute (JBI) critical appraisal checklist used methodological quality assessment of included studies. The methodological quality of the included cross-

sectional studies and the strategies they employed to address and minimize bias were evaluated using the JBI critical assessment instrument [17].

Statistical analysis

Data from studies that were included in a recent meta-analysis and systematic review (16) were statistically analyzed using the RevMan 5.3 program. In statistical terms, a p-value of less than 0.05 was deemed significant, and findings were presented as odds ratios (ORs) with a 95% confidence interval (CI). Furthermore, the Q test and I² statistics were used to quantify the heterogeneity. If the heterogeneity test revealed no significant difference, two models—a fixed-effects model and a random-effects model—were used.

RESULTS

Included Studies

The selection and screening of research articles related to the study aim "The Role of Alcohol Consumption in Cardiovascular Health" was performed by following the PRISMA guidelines in the recent meta-analysis and systematic review. About 430 research articles were extracted from three electronic databases after applying the above-mentioned search strategy. Only 209 papers were screened, and 221 articles were excluded before screening. Among those, only 189 articles were assessed for eligibility criteria, and the final number of research articles after applying exclusion criteria was 9 for the recent systematic review and meta-analysis as mentioned below in Figure 1.




Figure 1: Prisma Flowchart

Quality Assessment of included studies

The Cochrane Library tool assessed the risk of 2, including randomized studies from recent meta-analysis and systematic review as mentioned below in Figures 2 and 3.




Figure 2: Risk of Bias Graph of Included Studies




Figure 3: Graph of Risk Bias Summary of Included Studies

As mentioned above, JBI was applied for our 7 included studies due to the involvement of the cohort or cross-sectional studies about the use of moderate alcohol consumption and associated risk factors of cardiovascular disease. The quality assessment by the JBI checklist is given below in Table 1.

Questions	Zhang et al., 2004 [19]	Bell et al., 2017 [20]	Zatońska et al., 2021 [21]	Onat et al., 2009 [22]	Blomster et al., 2014 [24]	Brügger-Andersen et al., 2009 [25]	Levantesi et al., 2013 [26]
Were the two groups similar and recruited from the same population?	Y	N	Y	Y	N/A	Y	Y
Were the exposures measured similarly to assign people to both exposed and unexposed groups?	Y	Y	N	Y	Y	N	Y
Was the exposure measured in a valid and reliable way?							

	Y	N	Y	N	Y	N	Y
Were confounding factors identified?	N	Y	N/A	Y	N/A	Y	Y
Were strategies to deal with confounding factors stated?	Y	N	Y	N/A	N	UN	Y
Was the follow up time reported and sufficient to be long enough for outcomes to occur?	N	Y	N/A	N	N/A	Y	Y
Was appropriate statistical analysis used?	N	Y	N/A	UN	N/A	Y	Y
Were strategies to address incomplete follow up utilized?	Y	N	Y	N/A	N	UN	Y
Was follow up complete, and if not, were the reasons to loss to follow up described and explored?	Y	UN	N	Y	N/A	Y	Y
Were the outcomes measured in a valid and reliable way?	Y	N/A	Y	N	Y	N	Y
Were the groups/participants free of the outcome at the start of the study?	N	Y	N/A	Y	N/A	Y	Y
Were strategies to address incomplete follow up utilized?	N	Y	N/A	Y	N/A	Y	Y

Table 1: *Quality Assessment of Included Studies By JBI*

N: No, yes: Y, N/A; Not applicable, Un: unclear

Characteristics of included studies

The included articles for recent meta-analysis and systematic review were published between 2004 and 2024. All included studies discussed the use of moderate alcohol consumption in comparison to heavy alcohol consumption or abstaining from alcohol consumption. To produce heterogeneity of results, the included studies belong to 8 different countries: 1 in Israel [4], 2 in China [19, 23], 1 in the United Kingdom [20], 1 in Poland [21], 1 in Turkey [22] 1 in Australia [24], 1 in Norway [25], and 1 in Italy [26]. Table 2.

Author, year	Country	Study population	Sample size	Type of design	Alcohol intake	Incidence of cardiovascular events	blood pressure	lipid profile
Gepner et al., 2015 [4]	Israel	224 CVD patients with DB type 2	73 red wine group 151 controls	Randomized controlled trial	Moderate versus abstaining	Nil	Red wine=(S) -4.30 (-9.00 to 0.27) (D) -3.00 (-5.80 to -0.21) Control = (S) -4.80 (-9.70 to 0.14) (D) -0.9 (-3.8 to 2.1)	HDL: Red wine= 4.0 (1.2) to 3.6 (1.9 - 5.3) Control= 4.3 (1.4) to -0.08 (-0.44 to 0.27) LDL: Red wine= 94.7 (31.2) to 0.18 (-7.20 to 7.50) Control= 93.9 (30.5) to 2.1 (-5.1 to 9.4)
Zhang et al., 2004 [5]	China	12,352 CVD patients	Wine group	Cohort study	Moderate versus abstaining	Wine: HR 1.96 (1.30-2.93) Control: 0.86 (0.57-1.27)		
Bell et al., 2017 [6]	United Kingdom	1 937 360 adults with no CVD	1 356 152 moderate drinking 581 208 in control group (heavy drinking)	Cohort study	Moderate drinking vs Heavy drinking	Wine: 1.00 (0.55-1.19) Control=1.33 (1.09-1.63) Wine: 3368 out of 6053 Control: 507 out of 6053	Moderate: (S) 133.5 (17.1) Control: 129.3 (19.0)	
Zatońska et al., 2021 [7]	Poland	2021 participants with no CVD	1360 drinker's vs 661 abstainers	Cohort study	Moderate drinking vs abstaining	Wine: OR 1.66, CI 1.27-2.18 Control: OR 1.76, CI 1.22-2.53		
Onat et al., 2009 [8]	Turkey	3,443 participants with no CVD	577 Moderate and 93 heavy drinkers	Prospective cohort study	Moderate vs Heavy	Moderate: 19% (109) Heavy: 30% (29)	Systolic Moderate: 131.7±1.5 Heavy: 137.1±2.3 Diastolic Moderate: 82.9±0.9 Heavy: 85.8±1.4	HDL Moderate: 44.7±0.9* Heavy: 46±1.3 LDL Moderate: 123±2.5 Heavy: 126±3.9

							Hypertension Moderate: 84; %37.5 Heavy: 49; %54.4	
Au Yeung et al., 2013 [9]	China	4,867 participants with no CVD	2384 moderate drinker's vs 2482 abstaining	Randomized controlled trial	Moderate vs abstaining	CVD: 661 out of 2384 1098 out 2482	Systolic Moderate: 131.1 (19.3) Abstainers: 132.7 (21.1) Diastolic: Moderate: 75.3 (10.1) Abstainers: 75.8 (13.9)	HDL Moderate: 0.02 Abstainers: 0.08 LDL Moderate: -0.03 Abstainers: 0.08
Blomster et al., 2014 [10]	Australia	11140 participants	3389 moderate alcohol user vs 7751 non-alcohol users	Cohort study	Moderate vs abstaining	292 among moderate 817 nonalcoholic		
Brügger-Andersen et al., 2009 [11]	Norway	5477 patients	2753 moderate alcohol 545 heavy users	Cohort study	Moderate vs heavy	403 CVD events among moderate 141 among heavy users		
Levantesi et al., 2013 [12]	Italy	11, 248 participants	37021 moderate alcohol users 74227 nonalcoholic	Cohort study	Moderate vs abstaining	1168 CVE 3098 CVE		

Table 2: Characteristics of Included Studies [4-12]

Primary Outcomes

Incidence of Cardiovascular Events

Among the 9 included studies, about 8 studies discussed the incidence of cardiovascular events among moderate alcohol users and heavy or non-alcohol users [5-12]. There were significantly lower rates of CVD events among moderate alcoholic consumption in comparison to placebo (non-user or heavy users) which was evaluated through an odd ratio (Odds Ratio= 0.49; CI: 0.29 to 0.84; $p > 0.00001$), and heterogeneity was found ($df = 5$; $I^2 = 99\%$) as shown in Figure 4 and 5. The mean difference was evaluated in the given hazard risk ratio for three studies included [5, 6], as shown in Figure 6.




Figure 4: Forest Plot of CVD Incidence Among Moderate Alcohol Consumption and Heavy or Non-Alcohol Users [5,12]




Figure 5: Funnel Plot of CVD Incidence Among Moderate Alcohol Consumption and Heavy or Non-Alcohol Users [5, 12]




Figure 6: Mean Difference of CVD Hazard Risks Among Moderate Alcohol Consumption and Heavy or Non-Alcoholic Users [5, 6]

Blood Pressure

a) Systolic blood pressure

Among the 9 included studies, about 4 studies discussed the systolic and diastolic blood pressure among moderate alcohol users and heavy or non-alcohol users [4, 6, 8, 9]. There were significantly lower rates of systolic blood pressure (Mean difference = -4.10; CI: -4.12 to -4.04; p > 0.00001,) and diastolic blood pressure (Mean difference = 0.49; CI: 0.29 to 0.84; p > 0.00001,) among moderate alcoholic consumption in comparison to placebo (non-user or heavy users) and heterogeneity was found (df = 5; I² = 99%) as shown below in Figure 7 and 8.




Figure 7: Mean Difference of Systolic Blood Pressure Among Experimental And Control Groups [4, 6, 8, 9]

b) Diastolic blood pressure




Figure 8: Mean Difference of Systolic Blood Pressure Among Experimental And Control Groups [4, 6, 8, 9].

Lipid Profile

Among the 9 included studies, about 3 studies discussed the systolic and diastolic blood pressure among moderate alcohol users and heavy or non-alcohol users [4, 6, 8, 9]. There were significantly lower rates of LDL (Mean difference -4.39; CI: -4.50 to -4.28; p>0.00001,) and HDL (Mean difference = -1.49; CI: 1.14 to 0.93; p>0.00001,) among moderate alcoholic consumption in comparison to placebo (non-user or heavy users) and heterogeneity was found (df = 2; I² = 89%) as shown below in Figure 9 and 10.

a) LDL




Figure 9: Alcoholic Consumption in Comparison to Placebo (Non-User Or Heavy Users)
[4,6,8,9]

b) HDL




Figure 10: Alcoholic Consumption in Comparison to Placebo (Non-User Or Heavy Users)
[4,6,8,9]

Discussion

The study's main goal was to determine the association between patterns of alcohol intake and cardiovascular health outcomes, such as changes in lipid profiles, blood pressure measurements, and the frequency of cardiovascular events. The study found a strong correlation between moderate alcohol consumption and a lower risk of cardiovascular events using a thorough meta-analysis.

The statistical method of odds ratio analysis, which determines the probability of an occurrence, showed a strong protective effect of moderate alcohol use. In particular, moderate drinkers had a significantly reduced risk of cardiovascular events than heavy or non-drinkers. A computed Odds Ratio of 0.49 (CI: 0.29 to 0.84; $p > 0.00001$) demonstrated that this link was strong enough since it significantly shows the benefit of lower chances in cardiovascular events for moderate alcohol use.

An examination of the blood pressure assessments provided even more information to show the advantages of taking an average amount of alcohol in terms of cardiovascular health. The diastolic as well as the systolic values of the blood pressure had similar trends which favored moderate drinkers. For people with moderate drinking patterns, there was a decrease in both if it's a plus. This significant decrease in systolic and diastolic blood pressure was seen among those who were drinking moderately compared to those who never drank alcohol or did so heavily. The systolic and diastolic blood pressure were -4.10 (-4.12 to -4.04; $p < 0.00001$) and 0.49 (0.29 to 0.84; $p < 0.00001$) from each other which demonstrated the favorable effects of moderate alcohol intake on cardiovascular health as well as blood pressure control.

Secondly, the analysis of the lipid profile gave more understanding of the physiological influence that alcohol has on cholesterol levels. The moderate drinkers had lower levels of low-density lipoprotein (bad cholesterol) and high-density lipoprotein (good cholesterol) than heavy drinkers or non-drinkers. Our analysis revealed significant and favorable alterations in lipid profile associated with modest alcohol intake; mean differences were observed: -1.49 mmol/L (CI: -1.57 to -1.41, $p > 0.00001$) for high-density lipoprotein cholesterol (HDL-C) and -4.39 mmol/L (CI: -4.40 to -4.38, $p > 0.00001$) for low-density lipoprotein cholesterol (LDL-C).

Although there are statistically significant observations on alterations in lipid profiles, the variations identified in the studies ($df = 2$; $I^2 = 89\%$) highlight how complicated the relationship between alcohol consumption and lipid metabolism is. In this regard, more studies must be undertaken to comprehend this complexity well enough since the lipid profile changes can be influenced by other factors but not necessarily consumption of alcohol.

Implications and Future Research Directions

The findings inform medical advice and public health policy on clinical health promotion methods such as whether or not drinking moderately could be beneficial to our hearts. Doctors should be talking to their clients about moderate drinking habits because it could help control high blood pressure in ways other than taking medication. Nevertheless, one should be careful because the

study was diverse or had many dissimilar types of effects which means that we could still have some unknown or unfixed problems from a different source.

Limitations

Despite the benefits of research, it's important to understand its limitations. Results may not be as broadly applicable as they could otherwise be because of the possible bias introduced by self-reported alcohol intake data, and disparities in study design or methodology, among other things. Given that research findings are so varied, it underscores just how complex the association is between alcohol consumption and cardiovascular health; hence there is a need for more studies.

Conclusion

To sum up, our meta-analysis and systematic review help explain the intricate relationship between alcohol consumption and cardiovascular health, so that a sensible approach is necessary notwithstanding that mild to moderate alcohol has been linked with some advantages over either heavy or no drinking at all, including low risk of cardiovascular events, normal blood pressure, and favorable modifications in lipid profiles. People are different and learn differently, hence the need for person-centered alcohol consumption advice with the different risk profiles that exist among different patients in mind. Comparing and contrasting likely confounders will go a long way in substantiating this. To comprehend this work, future research is needed to test the interaction effect between gender and personality traits. Future research can fill up these gaps and offer more complex insights that will direct evidence-based cardiovascular health strategies.

COMPETING INTERESTS

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

1. Mendis S, Davis S, Norrving B. Organizational Update: The World Health Organization Global Status Report on Noncommunicable Diseases 2014; One More Landmark Step in the Combat Against Stroke and Vascular Disease. *Stroke*. 2015, 46:e121–2. <https://doi.org/10.1161/strokeaha.115.008097>
2. Puddu PE, Menotti A. Coronary heart disease differences in Eastern versus Western Europe: A demanding situation. *International Journal of Cardiology*. 2016, 217:S60–3. <https://doi.org/10.1016/j.ijcard.2016.06.225>
3. Parks D, Booyse F. Moderate Wine and Alcohol Consumption: Beneficial Effects on Cardiovascular Disease. *Thrombosis and Haemostasis*. 2001, 86:517–28. <https://doi.org/10.1055/s-0037-1616080>
4. Parks D, Booyse F. Moderate Wine and Alcohol Consumption: Beneficial Effects on Cardiovascular Disease. *Thrombosis and Haemostasis*. 2001, 86:517–28. <https://doi.org/10.1055/s-0037-1616080>
5. Wood AM, Kaptoge S, Butterworth AS, et al. Risk thresholds for alcohol consumption: combined analysis of individual-participant data for 599 912 current drinkers in 83 prospective studies. *The Lancet*. 2018, 391:1513–23. [https://doi.org/10.1016/s0140-6736\(18\)30134-x](https://doi.org/10.1016/s0140-6736(18)30134-x)

6. Xu W, Wang H, Wan Y, et al. Alcohol consumption and dementia risk: a dose–response meta-analysis of prospective studies. *European Journal of Epidemiology*. 2017, 32:31–42. <https://doi.org/10.1007/s10654-017-0225-3>
7. Biddinger KJ, Emdin CA, Haas ME, et al. Association of Habitual Alcohol Intake With Risk of Cardiovascular Disease. *JAMA Network Open*. 2022, 5:e223849–9. <https://doi.org/10.1001/jamanetworkopen.2022.3849>
8. Harmful Interactions | National Institute on Alcohol Abuse and Alcoholism (NIAAA). www.niaaa.nih.gov. Accessed: April 11, 2024. <http://pubs.niaaa.nih.gov/publications/Medicine/medicine.htm>.
9. Breslow RA, Dong C, White A. Prevalence of Alcohol-Interactive Prescription Medication Use Among Current Drinkers: United States, 1999 to 2010. *Alcoholism: Clinical and Experimental Research*. 2015, 39:371–9. <https://doi.org/10.1111/acer.12633>
10. Agrawal P, Halaweish F, Dwivedi C. Antioxidant Effects and Drug Interactions of Resveratrol Present in Wine. *Journal of Wine Research*. 2007, 18:59–71. <https://doi.org/10.1080/09571260701660839>
11. Borges G, Bagge CL, Cherpitel CJ, et al A meta-analysis of acute use of alcohol and the risk of suicide attempt. *Psychological Medicine*. 2016, 47:949–57. <https://doi.org/10.1017/s0033291716002841>
12. Haseeb S, Alexander B, Baranchuk A. Response by Haseeb et al to Letter Regarding Article, ‘Wine and Cardiovascular Health: A Comprehensive Review’. *Circulation*. 2018, 137:1880–1. <https://doi.org/10.1161/circulationaha.117.033086>
13. Holmes MV, Dale CE, Zuccolo L, et al.: Association between alcohol and cardiovascular disease: Mendelian randomisation analysis based on individual participant data. *BMJ*. 2014, 349:g4164–4. <https://doi.org/10.1136/bmj.g4164>
14. Chiva-Blanch G, Arranz S, Lamuela-Raventos RM, et al. Effects of Wine, Alcohol and Polyphenols on Cardiovascular Disease Risk Factors: Evidences from Human Studies. *Alcohol and Alcoholism*. 2013, 48:270–7. <https://doi.org/10.1093/alcalc/agt007>
15. Page MJ, McKenzie JE, Bossuyt PM, et al.: The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *British Medical Journal*. 2021, 372: <https://doi.org/10.1136/bmj.n71>

16. Sarkis-Onofre R, Catalá-López F, Aromataris E, et al. How to Properly Use the PRISMA Statement. *Systematic Reviews*. 2021, 10: <https://doi.org/10.1186/s13643-021-01671-z>
17. Santos WM dos, Secoli SR, Püschel VA de A. The Joanna Briggs Institute approach for systematic reviews. *Revista Latino-Americana de Enfermagem*. 2018, 26: <https://doi.org/10.1590/1518-8345.2885.3074>
18. Gepner Y, Golan R, Harman-Boehm I, et al.: Effects of Initiating Moderate Alcohol Intake on Cardiometabolic Risk in Adults With Type 2 Diabetes: A 2-Year Randomized, Controlled Trial. *Annals of internal medicine*. 2015, 163:569–79. <https://doi.org/10.7326/M14-1650>
19. Zhang L, Zhao L, Zhou B, et al. [Alcohol consumption and incidence of ischemic stroke in male Chinese]. *PubMed*. 2004, 25:954–7.
20. Bell S, Daskalopoulou M, Rapsomaniki E, et al.: Association between clinically recorded alcohol consumption and initial presentation of 12 cardiovascular diseases: population based cohort study using linked health records. *BMJ*. 2017, 356:j909. <https://doi.org/10.1136/bmj.j909>
21. Zatońska K, Psikus P, Basiak-Rasała A, et al.: Patterns of Alcohol Consumption in the PURE Poland Cohort Study and Their Relationship with Health Problems. *International Journal of Environmental Research and Public Health*. 2021, 18:4185. <https://doi.org/10.3390/ijerph18084185>
22. Altan Onat, Gülay Hergenç, Zekeriya Küçükdurmaz, et al. Moderate and heavy alcohol consumption among Turks: long-term impact on mortality and cardiometabolic risk. *PubMed*. 2009, 37:83–90.
23. Au Yeung SL, Jiang C, Cheng KK, et al.: Moderate Alcohol Use and Cardiovascular Disease from Mendelian Randomization. *PLoS ONE*. 2013, 8:e68054. <https://doi.org/10.1371/journal.pone.0068054>
24. Blomster JI, Zoungas S, Chalmers J, et al.: The Relationship Between Alcohol Consumption and Vascular Complications and Mortality in Individuals With Type 2 Diabetes. *Diabetes Care*. 2014, 37:1353–9. <https://doi.org/10.2337/dc13-2727>
25. Brügger-Andersen T, Pönitz V, Snapinn S, et al. Moderate alcohol consumption is associated with reduced long-term cardiovascular risk in patients following a complicated

- acute myocardial infarction. *International Journal of Cardiology*. 2009, 133:229–32. <https://doi.org/10.1016/j.ijcard.2007.12.046>
26. Levantesi G, Marfisi R, Mozaffarian D, et al. Wine consumption and risk of cardiovascular events after myocardial infarction: Results from the GISSI-Prevenzione trial. *International Journal of Cardiology*. 2013, 163:282–7. <https://doi.org/10.1016/j.ijcard.2011.06.053>
27. Male, C. K. V. L. S. N. A., Swathi, V., Kumar, D. S., Karishma, . and Yunus, M. S. (2021) “Hyperlipidemia Condition and Novel-Drug Therapies: A Overall Study”, *Journal of Pharmaceutical Research International*, 33(40B), pp. 171–186. doi: 10.9734/jpri/2021/v33i40B32277.
28. Mashahit, Mohamed, Eman Ezzat, and Eman Hanafy. 2017. “Prevalence of Non-Communicable Diseases in Patients With Type 2 Diabetes Mellitus in Egypt; A Retrospective Survey”. *Asian Journal of Medicine and Health* 7 (1):1-9. <https://doi.org/10.9734/AJMAH/2017/36089>.
29. Mukamal KJ, Rimm EB. Alcohol consumption: risks and benefits. *Current atherosclerosis reports*. 2008 Dec;10(6):536-43.
30. Peele S, Brodsky A. Exploring psychological benefits associated with moderate alcohol use: a necessary corrective to assessments of drinking outcomes?. *Drug and alcohol dependence*. 2000 Nov 10;60(3):221-47.