

EMERGING ~~PATHOGENS-DISEASES~~ AND THEIR IMPACT ON GLOBAL CROP PRODUCTION

ABSTRACT

Emerging ~~pathogens-diseases in agriculture~~ are not a new concern ~~in agriculture~~ but have ~~recently~~ gained ~~heightened-more~~ attention due to their increasing frequency and severity. A well-known example is the potato late blight, which was brought on by the oomycete *Phytophthora infestans* and is infamous for causing the Irish Potato Famine in the 1840s. ~~A major factor is that~~ ~~crops are~~ have become more vulnerable to diseases due to ~~changes-the fluctuations~~ in the growing environments ~~brought about~~ caused by climate change. ~~Changes-Variations~~ in precipitation patterns and ~~rising-increasing~~ temperatures can foster an environment ~~that is~~ conducive to the growth and ~~dissemination-spread~~ of diseases. Furthermore, ~~Agricultural-agricultural~~ intensification, ~~characterized by~~ with its emphasis on monocropping and high-density planting, ~~provides-creates~~ an ideal environment for the rapid spread of pathogens ~~to spread rapidly~~. ~~The-Due to the~~ lack of genetic diversity in monocultures, ~~means that~~ once a pathogen infects a crop, it can quickly decimate entire fields. ~~Mechanisms of disease spread and innovative approaches of management of emerging diseases are discussed~~. By prioritizing research, implementing sustainable practices, and fostering global cooperation, it is possible to mitigate the impact of these pathogens on global crop production and ensure food security for future generations.

Formatted: Justified

Formatted: Font: Italic

Keywords: pathogens, global crop production, *Fusarium oxysporum*, wheat blast

1. INTRODUCTION

Background

In the contemporary era of agriculture, the rise of emerging ~~pathogens-disease~~ presents a significant threat to global crop production. These ~~pathogens-diseases caused by, encompassing~~ -fungi, bacteria, viruses, and nematodes, can lead to devastating crop losses, affecting both the yield and quality of agricultural produce. Historically, the agricultural sector has ~~continuously~~ ~~ually~~ battled ~~with~~ various plant diseases; On the other hand, the ~~advent occurrence~~ and dissemination of new infections have been expedited by trade ~~globalisation~~ ~~globalization~~ and climate change. The aforementioned phenomena present a significant obstacle to both food security and economic stability, especially in areas that strongly depend on agriculture.

Emerging pathogens in agriculture are not a new concern but have gained ~~heightened-more~~ attention due to their increasing frequency and severity. A well-known example is the potato late blight, which was brought on by the oomycete *Phytophthora infestans* and is infamous for causing the Irish Potato Famine in the 1840s. Similar concerns also exist today due to the emergence of new, ~~and more~~ ~~deadly~~ ~~virulent~~ strains of existing infections or their ~~comeback-appearance~~ in ~~hitherto-unexplored~~ ~~new~~ ~~areas~~ ~~locations~~. The reappearance of Panama disease in bananas due to *Fusarium oxysporum* f. sp. *cubense* TR4 and the wheat blast produced by the fungus *Magnaporthe oryzae*, ~~for example, highlight~~ ~~how~~ illustrate persistent and dynamic ~~nature of~~ these threats ~~are~~ (Savary et al., 2019).

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Importance of Crop Production in Global Food Security

~~The-Crop production is the~~ foundation of global food security and the source of ~~nutrition for~~ the world's ~~population's~~ ~~nourishment is~~ crop production. Since plant-based meals provide more than 80% of human calories, ~~the Food and Agriculture Organisation (FAO) states that~~ it is imperative ~~that to~~

~~consistently produce~~ crops ~~including such as~~ wheat, rice, maize, and soybeans ~~are produced consistently~~ (FAO, 2020). These staple crops are not only vital for direct human consumption but also serve as essential feed for livestock, linking crop production directly to meat and dairy production.

Strong crop production systems are crucial because of the growing world population, which is expected to reach 9.7 billion by 2050 and increase food consumption (United Nations, 2019). However, emerging ~~pathogens diseases~~ threaten to destabilize these systems, leading to potential food shortages and increased food prices. The impact is ~~disproportionately severe~~ significantly greater in developing ~~countries nations~~ where agricultural practices may be less advanced and economic resilience ~~lower weaker~~. The 2008 global food crisis, exacerbated by crop failures in major producing regions, illustrated the fragile nature of food security and the far-reaching consequences of disruptions in crop production.

In addition to food security, crop production is integral to economic stability and development. Agriculture employs a significant portion of the global workforce, particularly in developing nations where it can account for up to 60% of employment (World Bank, 2021). Crop failures, therefore, have a cascading effect, leading to unemployment, reduced incomes, and increased poverty levels. Thus, ~~safeguarding protecting~~ crop production from emerging ~~pathogens diseases~~ is crucial for ~~not only a matter of~~ food security but also economic and social stability.

Definition and Scope of Emerging ~~Pathogens Diseases~~

Emerging ~~pathogens diseases~~ are broadly defined as infectious agents that have recently increased in incidence or geographic range, or are newly recognized in a specific crop host (Anderson et al., 2004). These ~~pathogens diseases~~ can be fungi, bacteria, viruses, nematodes, or oomycetes ~~that cause diseases~~ ~~may be previously unknown~~ new or re-emerging diseases that were once under control. The emergence of these ~~pathogens diseases~~ can be attributed to ~~several factors, including~~ climate change, international trade, agricultural intensification, and changes in land use.

Commented [GN1]: Mentioned in the introduction. Repetition. May be removed.

~~Crops have become more vulnerable to diseases due to the fluctuations in the growing environments caused by climate change. Variations in precipitation patterns and increasing temperatures can foster an environment conducive to the growth and spread of diseases. A major factor is that crops are more vulnerable to diseases due to changes in the growing environments brought about by climate change. Changes in precipitation patterns and rising temperatures can foster an environment that is conducive to the growth and dissemination of diseases.~~ For instance, the geographic range of rust diseases in wheat has expanded due to changing climatic conditions, posing new challenges for disease management (Chakraborty Newton, 2011).

~~International trade facilitates the movement of pathogens across borders, often inadvertently. The globalization of agricultural trade means that crops and their associated pathogens can move quickly from one region to another. The introduction of *Xylella fastidiosa* to Europe, which devastated olive groves in Italy, is a stark reminder of how international trade can facilitate the spread of harmful pathogens (EFSA, 2019).~~

Formatted: Font: Italic

Agricultural intensification, ~~with its emphasis on monocropping and high-density planting, creates characterized by monocropping and high density planting, provides an ideal environment for the rapid spread of pathogens to spread rapidly. Due to the lack of genetic diversity in monocultures, once a pathogen infects a crop, it can quickly decimate entire fields. The lack of genetic diversity in monocultures means that once a pathogen infects a crop, it can quickly decimate entire fields.~~ This was observed in the outbreak of cassava mosaic disease in Africa, which spread rapidly through fields of genetically uniform cassava plants (Legg et al., 2014).

Table 1: Key Emerging Pathogens Affecting Global Crop Production

Crop	Pathogen	Disease	Impact
Wheat	<i>F. oxysporum</i> f. sp. <i>cubense</i>	Fusarium wilt	Threat to modern banana varieties, causing devastation.

Formatted: Font: Italic

Formatted: Justified

Formatted: Font: Italic

Wheat	<i>Pyricularia oryzae</i> pathotype Triticum	Wheat blast	High yield losses, potential spread to new regions.
Banana	<i>Xanthomonas vasicola</i> pv. <i>musacearum</i>	BXW (Banana Xanthomonas Wilt)	Major threat to banana production worldwide.
Potato	<i>Phytophthora</i> <i>infestans</i>	Late blight	Devastating potato disease, responsible for historical famines.
Various	<i>Phakopsora</i> <i>pachyrhizi</i> & <i>Zymoseptoria tritici</i>	Soybean rust & Wheat blotch	Fungal diseases causing significant yield losses in major crops.

- Formatted: Font: Italic
- Formatted: Justified
- Formatted: Font: Italic
- Formatted: Justified
- Formatted: Font: Italic
- Formatted: Font: Italic
- Formatted: Justified
- Formatted: Font: Italic
- Formatted: Justified
- Formatted: Font: Italic

Changes in land use, such as deforestation and urbanization, also contribute to the emergence of new ~~pathogens~~diseases. These activities disrupt natural ecosystems and can lead to the spillover of pathogens from wild plants to crops. The emergence of *Pseudomonas syringae* pv. *actinidiae* in kiwifruit orchards in New Zealand is one such example, where changes in land use and global trade played a role in the pathogen's spread (Vanneste et al., 2011).

- Formatted: Font: Italic
- Formatted: Font: Italic

Understanding the scope of emerging ~~pathogens~~diseases requires a multidisciplinary approach, encompassing plant pathology, climate science, trade policies, and agricultural practices. Addressing the challenges posed by these pathogens involves integrated disease management strategies, including the development of resistant crop varieties, the implementation of strict biosecurity measures, and the adoption of sustainable farming practices.

2. OVERVIEW OF EMERGING ~~PATHOGENS~~DISEASES

Definition and Classification

~~Emerging pathogens are microorganisms, including bacteria, fungi, viruses, and nematodes, that have recently increased in incidence, geographical distribution, or host range, or have developed new strains with enhanced virulence. These pathogens can significantly impact crop production by reducing yields, affecting quality, and increasing the costs of management and control measures (Anderson et al., 2004). The emergence of these pathogens can be attributed to various factors, including environmental changes, agricultural practices, and global trade.~~

Commented [GN2]: Definition has been discussed abundantly in previous sub division. Redundant.

Emerging ~~pathogens~~diseases can be classified based on ~~their taxonomic~~the taxonomic groups of the ~~causative agents~~, such as bacterial, viral, fungal, or nematode pathogens. Another classification criterion involves their mode of transmission and spread. For example, some pathogens are soil-borne, while others are seed-borne or vector-borne, each having different implications for control strategies. Understanding the classification helps in devising appropriate management practices tailored to the specific pathogen type (Fisher et al., 2012).

Mechanisms of Emergence

The emergence of new pathogens or new strains of existing pathogens can be driven by multiple mechanisms. These mechanisms include:

Genetic Changes: Pathogens can evolve through mutation, recombination, or horizontal gene transfer, leading to new strains with increased virulence or resistance to existing control measures (Fisher et al., 2012). Genetic diversity in pathogens can result in the emergence of strains capable of overcoming plant resistance genes, ~~making~~rendering previously resistant crop varieties susceptible.

Environmental Changes: Climate change, including shifts in temperature and precipitation patterns, can create favorable conditions for pathogen survival, reproduction, and spread (Bebber et al., 2013). For instance, warmer temperatures can extend the growing season and geographic range of certain pathogens.

Agricultural Practices: Changes in agricultural practices, such as the intensification of farming, monoculture planting, and the overuse of chemical inputs, can lead to the emergence of pathogens. Monocultures, in particular, can provide a large, uniform host population that facilitates pathogen spread (Anderson et al., 2004).

Commented [GN3]: Repeated.

Global Trade and Movement: The globalization of trade and the movement of plant materials across borders can introduce pathogens to new regions where they previously did not exist. This can result in outbreaks in areas where crops lack resistance to these newly introduced pathogens (Bebber et al., 2013).

Changes in Host Susceptibility: The introduction of new crop varieties, which may be more susceptible to certain pathogens, can also play a role in the emergence of these diseases. Additionally, the breakdown of resistance genes in crops due to genetic changes in pathogens can make previously resistant crops vulnerable (Fisher et al., 2012).

Commented [GN4]: These have been adequately mentioned in previous paragraphs. So, they can be presented concisely

Examples of Recent Emerging Pathogens/diseases Affecting Crops

Commented [GN5]: Please refer to the comment on 7. CASE STUDIES OF EMERGING PATHOGENS. Please combine both to avoid repetition.

Wheat Blast (*Magnaporthe M. oryzae* Triticum): Initially reported in Brazil in the mid-1980s, wheat blast is a fungal disease caused by a particular strain that has subsequently moved to Bangladesh and Zambia. Because it can quickly and severely reduce output under the right circumstances, this pathogen is a serious danger to the world's wheat supply (Islam et al., 2016).

Banana Fusarium Wilt (*Fusarium F. oxysporum* f. sp. cubense TR4): The deadly Panama disease in bananas is caused by the Tropical Race 4 (TR4) strain of *Fusarium oxysporum* f. sp. cubense. TR4 has spread from Southeast Asia to other banana-growing regions, threatening global banana production due to its persistence in the soil and the lack of effective control measures (Ploetz, 2015).

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Cassava Brown Streak Disease (CBSD): Two viruses—the Ugandan cassava brown streak virus (UCBSV) and the cassava brown streak virus (CBSV)—cause CBSD. Due to the substantial tuber loss and impact on the region's food security, this disease has become a severe danger to East Africa's cassava output (Nichols, 1950; Alicai et al., 2007).

Pierce's sickness (*Xylella X. fastidiosa*): Recently, we have seen the emergence of several diseases caused by this bacterial pathogen, which is impacting many different crops, such as citrus, grapes, and olives is observed. In grapes, *Xylella X. fastidiosa* is the source of illnesses including Pierce's sickness and Olive Quick Decline Syndrome. Concerns have been expressed over its possible effects on agricultural output and trade as a result of its spread throughout Europe, especially in Italy and Spain (European Food Safety Authority, 2015).

Formatted: Font: Italic

Formatted: Font: Italic

Tomato Brown Rugose Fruit Virus (ToBRFV): Since its discovery in Israel in 2014, ToBRFV has quickly spread to a number of nations, including the US, Mexico, and several European states. This virus affects tomatoes and peppers, causing severe fruit symptoms and reducing marketable yields. Its rapid spread and the lack of resistant varieties pose significant challenges to growers (Salem et al., 2016).

In conclusion, emerging pathogens present a significant threat to global crop production, necessitating continuous monitoring, research, and the development of integrated management strategies to mitigate their impact. Understanding the mechanisms driving their emergence and implementing effective control measures are crucial for safeguarding food security in the face of amidst these evolving challenges.

Emerging pathogens/diseases present a significant threat to global crop production, impacting agricultural systems worldwide. These pathogens include bacteria, viruses, fungi, and nematodes that cause diseases leading to severe economic losses, reduced yield and quality, and increased costs for management and control. This paper explores the economic consequences, effects on yield and quality, and provides case studies of major crop pathogens/diseases, categorized into bacterial, viral, fungal, and nematode pathogens based on the pathogens.

Economic Consequences

~~Emerging crop pathogens have profound economic impacts on global agriculture.~~ The costs associated with the ~~se~~ ~~pathogens~~ ~~emerging~~ ~~diseases~~ include direct losses from reduced yields and quality, as well as indirect costs related to management and control measures.

Direct Economic Losses: ~~Crop losses due to emerging pathogens can be devastating. For example,~~ ~~€~~The global annual loss due to plant diseases is estimated to be around 10-16% of the total crop production, translating to billions of dollars annually (Savary et al., 2019). The impact is particularly severe in developing countries where agricultural practices may be less advanced, and resources for disease management are limited.

Indirect Costs: Indirect costs include expenses for disease management, such as the development and application of pesticides, investment in breeding for disease-resistant crop varieties, and the implementation of quarantine measures. The cost of developing resistant varieties can be substantial, and these varieties may not always be fully effective due to the rapid evolution of pathogens (Anderson et al., 2004).

Market and Trade Disruptions: Pathogen outbreaks can lead to trade restrictions and reduced market access for affected countries. For instance, the outbreak of Citrus Greening Disease (Huanglongbing) in Florida caused significant economic losses, not only due to decreased production but also because of export restrictions imposed by trading partners (Bové, 2006).

Table 2: Economic Impact of Emerging ~~diseases~~ ~~Pathogens~~ on Crop Production

Causative Pathogen/agent	Economic Losses (in \$ millions)	Affected Regions	Year
<i>Phytophthora P. infestans</i>	1,000	North America, Europe	2019
<i>Fusarium F. oxysporum</i>	500	Asia, Africa	2020
<i>Xylella X. fastidiosa</i>	1,200	Europe, South America	2021
<i>Puccinia P. graminis</i>	800	Worldwide	2022
<i>Magnaporthe M. oryzae</i>	950	Southeast Asia, Africa	2023
<i>Ralstonia R. solanacearum</i>	750	South America, Asia	2023
Spodoptera frugiperda	1,100	Africa, Americas	2023

Effects on Yield and Quality

Emerging pathogens affect crop yield and quality in multiple ways, leading to significant challenges for global food security.

Reduced Yield: ~~Pathogens-Diseases~~ can cause a substantial reduction in crop yield by damaging plant tissues, disrupting physiological processes, and reducing photosynthetic efficiency. For example, the wheat rust fungus (*Puccinia spp.*) can lead to yield losses of up to 70% in susceptible wheat varieties (Singh et al., 2016).

Quality Deterioration: The quality of crops can be severely compromised by ~~pathogens~~ ~~diseases~~. This includes changes in physical appearance, nutritional content, and storability. For instance, potato late blight ~~caused by Phytophthora infestans not only~~ reduces yield ~~but also affects~~ and tuber quality, leading to significant post-harvest losses (Fry, 2008).

Toxin Production: ~~Some~~ ~~During some infections, the~~ pathogens produce toxins that contaminate food crops, posing health risks to humans and animals. A notable example is the aflatoxin produced by *Aspergillus flavus* in crops like maize and peanuts, which can lead to serious health issues and economic losses due to reduced marketability (Wu, 2006).

Case Studies of Major Crop Pathogens

To illustrate the impact of emerging pathogens on global crop production, we examine specific examples of bacterial, viral, fungal pathogens, and nematodes.

Bacterial Pathogens

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Highlight

Commented [GN6]: This is an insect pest and thus, is beyond the scope of discussion here.

Formatted: Font: Italic

Formatted: Font: Italic

Commented [GN7]: Many portions of this section are repetitive of the discussion under "Examples of Emerging diseases Affecting Crops". Please combine them.

Please italicize the names of organisms.

Formatted: Highlight

Xylella fastidiosa: This bacterium affects a wide range of plants, including olive trees, grapevines, and citrus. The disease caused by *X. fastidiosa* has led to significant losses in southern Europe, particularly in Italy where it has devastated olive groves (Saponari et al., 2019). Management strategies include removing infected plants and controlling insect vectors, but these measures are costly and not always effective.

Ralstonia solanacearum: This soil-borne pathogen causes bacterial wilt in many crops, including potatoes, tomatoes, and bananas. It is particularly problematic in tropical and subtropical regions. The disease can result in total crop loss, and the bacterium's ability to persist in the soil makes it challenging to manage (Álvarez et al., 2010).

Viral Pathogens

Tomato Brown Rugose Fruit Virus (ToBRFV): This newly emerged virus has caused severe damage to tomato crops in various countries. It leads to mottling and brown spots on fruits, significantly reducing their market value. The virus spreads rapidly through mechanical transmission and contaminated seeds, making it difficult to control (Panno et al., 2019).

Banana Bunchy Top Virus (BBTV): BBTV is a major threat to banana production worldwide. It causes stunted growth and "bunchy" appearance of leaves, leading to significant yield losses. The virus is transmitted by aphids, and control measures include using virus-free planting material and managing vector populations (Blomme et al., 2017).

Fungal Pathogens

Fusarium oxysporum: This fungus causes wilt illnesses in a variety of crops, such as tomatoes and bananas (also known as Panama disease or Fusarium wilt). There have been catastrophic losses in banana farms in Asia, Africa, and Latin America due to the *Fusarium* tropical race 4 (TR4) strain. Because the fungus lingers in the soil, it is quite challenging to remove (Ploetz, 2015).

Magnaporthe oryzae: This fungus is one of the most harmful diseases affecting rice production worldwide and is the cause of rice blast disease. It results in severe yield losses by causing lesions on panicles, stems, and leaves. The disease can reduce rice production by up to 30%, with severe outbreaks causing even higher losses (Dean et al., 2012).

Nematodes

Meloidogyne spp. (Root-Knot Nematodes): These nematodes infect the roots of many crops, causing galls that interfere with water and nutrient uptake. Crops like tomatoes, potatoes, and carrots are particularly vulnerable. Root-knot nematodes can cause yield losses of up to 50% in heavily infested fields (Jones et al., 2013).

Heterodera spp. (Cyst Nematodes): These nematodes form cysts on the roots of host plants, disrupting their growth and development. Soybeans and sugar beets are major crops affected by cyst nematodes. Management strategies include crop rotation and resistant varieties, but these measures are not always effective (Nicol et al., 2011).

4. MECHANISMS OF PATHOGEN SPREAD

Global Trade and Movement of Plant Material

The globalization of trade has significantly increased the movement of plant materials across borders, facilitating the spread of plant pathogens on an unprecedented scale. The international trade of seeds, plants, and agricultural products can introduce pathogens to new regions where they previously did not exist, leading to severe outbreaks that can devastate local agriculture.

Introduction of Pathogens Through Trade

The trade of plants and plant products is a primary vector for the spread of pathogens. For instance, the accidental introduction of *Phytophthora infestans*, the causative agent of potato late blight, to Europe in the 19th century led to the Irish Potato Famine (Ristaino et al., 2001). Similarly, the global trade of

Commented [GN8]: Please see the above comment. Either remove the previous section on "Mechanism....." or combine both.

Commented [GN9]: Sub sections may be numbered.

Formatted: Highlight

Commented [GN10]: Repetitive

ornamental plants has been linked to the spread of the bacterium *Xylella fastidiosa*, which causes diseases in multiple crops, including olives and citrus (EFSA Panel on Plant Health, 2015).

Commented [GN11]: Repetitive

Contaminated Plant Material

Contaminated seeds and nursery stocks are common pathways for pathogen dissemination. Seed-borne pathogens, such as fungi, bacteria, and viruses, can remain dormant within seeds and become active when conditions are favorable for growth (Franceschini et al., 2019). The movement of nursery stocks, especially those that are asymptomatic, presents a significant risk, as pathogens can remain undetected until after the plants have been distributed and planted.

Regulatory Challenges

Regulating the international trade of plant materials is challenging due to the vast volume of trade and the complexity of monitoring and inspecting shipments. Although phytosanitary measures are in place, the effectiveness of these regulations varies by country, and compliance can be inconsistent (Mumford et al., 2016). The implementation of stringent biosecurity protocols and the development of rapid diagnostic tools are crucial to mitigate the risk of pathogen spread through global trade.

Climate Change and Its Role in Pathogen Emergence

Climate change is altering the dynamics of plant-pathogen interactions, leading to the emergence and re-emergence of plant diseases. Changes in temperature, precipitation patterns, and atmospheric CO₂ levels can influence pathogen survival, reproduction, and dispersal, thereby impacting crop production globally.

Commented [GN12]: Repetitive

Altered Pathogen Life Cycles

Rising temperatures and changing precipitation patterns can extend the growing seasons of crops, which may provide more opportunities for pathogens to complete their life cycles and increase their populations. For instance, the increased incidence of rust diseases in cereals has been linked to warmer temperatures and altered rainfall patterns (Chakraborty Newton, 2011). Additionally, some pathogens may expand their geographical range to higher latitudes and altitudes as previously inhospitable climates become suitable for their growth (Bebber et al., 2013).

Enhanced Virulence and Resistance Breakdown

Climate change can also affect the virulence of pathogens and the resistance of host plants. Elevated CO₂ levels can enhance the virulence of certain pathogens, such as fungal pathogens, by increasing their growth rates and sporulation (Chakraborty et al., 2000). Moreover, changes in environmental conditions can stress plants, making them more susceptible to infections and reducing the effectiveness of resistance genes (Burdon Zhan, 2020).

Interaction with Other Stress Factors

Climate change often interacts with other environmental stressors, such as drought and nutrient deficiencies, to exacerbate plant disease outbreaks. For example, drought-stressed plants are more vulnerable to pathogen attacks, as their defense mechanisms are compromised (Ramegowda Senthil-Kumar, 2015). Understanding the complex interactions between climate change, plant physiology, and pathogen biology is essential for developing effective disease management strategies.

Agricultural Practices and Monocultures

Modern agricultural practices, particularly the widespread use of monocultures, have created environments conducive to the spread of pathogens. The simplification of agroecosystems, combined with intensive farming practices, has led to increased susceptibility of crops to diseases.

Commented [GN13]: Repetitive

Monocultures and Genetic Uniformity

In monocultures, a single crop species is grown throughout huge lands, resulting in genetic homogeneity among farms. Due to a uniform vulnerability to pathogens brought on by this lack of genetic variation, diseases can spread quickly and spark serious epidemics. For instance, the extensive planting of genetically uniform hybrid maize, which was extremely sensitive to the pathogen *Bipolaris maydis*,

Formatted: Font: Italic

contributed to the Southern Corn Leaf Blight outbreak in the United States throughout the 1970s (Ullstrup, 1972).

Commented [GN14]: Discussed already. Please combine the examples.

Intensification and Crop Rotation

Intensive farming methods can degrade soil health and increase pathogen pressure, such as continuous cropping and reduced crop rotation. When the same species is continuously farmed, the soil is deprived of the chance to break pathogen life cycles, which might result in the accumulation of soilborne infections. For example, the take-all disease caused by *Gaeumannomyces graminis* var. *tritici* may become more common if wheat is continuously cultivated (McDonald Stukenbrock, 2016).

Formatted: Font: Italic

Formatted: Font: Italic

Use of Chemical Inputs

The dynamics of infections can also be impacted by an over-reliance on chemical inputs like pesticides and fertilisers. While the use of fungicides might promote the growth of resistant pathogen strains, excessive fertiliser use can cause nutritional imbalances that impair plant defences (Lucas et al., 2015). Sustainable treatment of plant diseases requires the application of integrated pest management (IPM) techniques, which integrate cultural practices, biological control, and the sparing use of pesticides.

Commented [GN15]: Integrated Disease Management (IDM)???

~~The spread of emerging pathogens and their impact on global crop production are influenced by a multitude of factors, including global trade, climate change, and agricultural practices. Understanding these mechanisms is crucial for developing effective strategies to mitigate the risks and ensure global food security.~~ Collaborative efforts among researchers, policymakers, and farmers are essential to address the challenges posed by emerging plant pathogens diseases in a rapidly changing world.

5. DETECTION AND DIAGNOSIS

Commented [GN16]: Sub sections may be numbered.

Table 3: Pathogen Detection Methods

Method	Pathogen Detected	Accuracy (%)	Time Required	Cost	Description
PCR (Polymerase Chain Reaction)	Various viruses, fungi	95	3-4 hours	High	A molecular technique used to amplify and detect DNA sequences.
ELISA (Enzyme-Linked Immunosorbent Assay)	Bacteria, viruses	90	2-3 hours	Medium	A plate-based assay technique for detecting and quantifying substances.
LAMP (Loop-mediated Isothermal Amplification)	Bacteria, viruses	92	1-2 hours	Medium	A single-tube method that amplifies DNA while maintaining a steady temperature.
qPCR (Quantitative PCR)	Various pathogens	98	2-3 hours	High	A PCR-based method to quantify DNA or RNA in a sample.
Immunofluorescence	Viruses, bacteria	85	1-2 hours	High	Uses antibodies labeled with a fluorescent

					dye to detect antigens.
Microarray Analysis	Multiple pathogens simultaneously	93	4-6 hours	Very High	A high-throughput method to study many genes at once for variations and expression levels.
Serological Tests	Bacteria, viruses	88	1-2 hours	Low	Detects antibodies or antigens in blood, often used for viral infections.
Next-Generation Sequencing (NGS)	All types of pathogens	99	24-48 hours	Very High	A high-throughput method to determine the sequence of nucleotides in DNA.
Biosensors	Specific bacteria, viruses	90	30 minutes - 1 hour	Medium	Uses a biological component to detect the presence of various pathogens.
Lateral Flow Assays	Viruses, bacteria	80	10-30 minutes	Low	Simple device intended to detect the presence of a target substance in a liquid sample.

Traditional Methods

Traditional methods for detecting and diagnosing plant pathogens have been foundational in plant pathology for decades. These methods primarily include visual ~~inspections~~inspections (Schumann D'Arcy, 2006), cultural ~~techniques~~techniques (Agrios, 2005), and biochemical ~~tests~~tests (Clark Adams, 1977).

Visual Inspection

~~Visual inspection is the oldest and most straightforward method for detecting plant pathogens. Agronomists and farmers rely on the presence of symptoms such as wilting, chlorosis, necrosis, and abnormal growth patterns to identify diseases. Nevertheless, visual inspection is frequently arbitrary and susceptible to environmental influences, rendering it unreliable for accurate pathogen identification and early detection (Schumann D'Arcy, 2006).~~

Cultural Techniques

Cultural techniques involve isolating pathogens on nutrient media under laboratory conditions. These techniques allow for the identification of fungi, bacteria, and viruses based on their growth patterns, morphology, and physiological characteristics. For instance, the use of selective media can help in isolating specific pathogens like *Fusarium* spp. or *Pseudomonas* spp. To effectively interpret the results, this procedure necessitates skilled staff and is time-consuming (Agrios, 2005).

Formatted: Font: Italic

Formatted: Font: Italic

Biochemical Tests

Biochemical tests, such as enzyme-linked immunosorbent assay (ELISA) and other serological assays, are used to detect specific proteins or antigens associated with plant pathogens. ELISA has been widely used due to its sensitivity and ability to process multiple samples simultaneously. Despite its effectiveness, ELISA requires the availability of specific antibodies and can sometimes produce false positives due to cross-reactivity (Clark Adams, 1977).

Advanced Molecular Techniques

Advanced molecular techniques such as Polymerase Chain Reaction (Schena et al., 2013), Next-Generation Sequencing (Massart et al., 2014), Loop-Mediated Isothermal Amplification (Notomi et al., 2000)

have revolutionized the detection and diagnosis of plant pathogens, offering high sensitivity, specificity, and rapid results.

Polymerase Chain Reaction (PCR)

PCR-based methods, including conventional PCR, real-time PCR (qPCR), and reverse transcription PCR (RT-PCR), are among the most widely used molecular techniques. These methods amplify specific DNA or RNA sequences of pathogens, enabling their detection even at low concentrations. Real-time PCR, in particular, allows for quantitative analysis and has been used to detect and quantify various pathogens such as *Phytophthora* spp. and *Xanthomonas* spp. (Schena et al., 2013). PCR techniques are highly reliable but require specialized equipment and expertise.

Formatted: Font: Not Bold

Next-Generation Sequencing (NGS)

NGS technologies have further enhanced pathogen detection by enabling comprehensive analysis of plant microbiomes. Metagenomics, a subfield of NGS, allows for the identification of both known and unknown pathogens by sequencing all genetic material in a sample. This approach has been pivotal in identifying emerging pathogens that were previously undetectable by traditional methods (Massart et al., 2014). NGS is powerful but expensive and generates large volumes of data that require complex bioinformatic analysis.

Formatted: Font: Not Bold

Loop-Mediated Isothermal Amplification (LAMP)

LAMP is a rapid, cost-effective molecular technique that amplifies DNA at a constant temperature, eliminating the need for thermal cycling equipment. It is highly specific and sensitive, making it suitable for on-site testing and resource-limited settings. LAMP has been successfully applied in detecting pathogens like *Ralstonia solanacearum* and *Fusarium oxysporum* (Notomi et al., 2000). However, like other molecular techniques, LAMP requires initial sample preparation and validation.

Formatted: Font: Not Bold

Remote Sensing and Digital Tools

Large-scale early diagnosis and monitoring of plant diseases is now possible because to the integration of Remote sensing (Rumpf et al., 2010), Drones, or unmanned aerial vehicles, or UAVs (Zhang et al., 2020), Digital Disease Diagnostic Tools including Mobile applications and digital platforms (Barbedo, 2019) in agriculture. digital tools and remote sensing in agriculture.

Remote Sensing

Remote sensing involves the use of satellite or aerial imagery to monitor crop health and detect disease outbreaks. Techniques such as multispectral and hyperspectral imaging capture data across various wavelengths, revealing changes in plant physiology that may indicate disease presence. For example, diseased plants often show distinct spectral signatures due to stress-induced changes in pigment composition (Rumpf et al., 2010). Remote sensing provides a non-invasive, large-scale monitoring solution but requires sophisticated data processing and ground truth validation.

Unmanned Aerial Vehicles (UAVs)

Drones, or unmanned aerial vehicles, or UAVs, are becoming indispensable in precision farming. UAVs with sophisticated sensors can fly over fields to take detailed pictures and gather information about crop conditions. This technology allows for timely identification of disease hotspots, enabling targeted interventions. UAV-based remote sensing has been effectively used to detect early signs of diseases such as late blight in potatoes and stripe rust in wheat (Zhang et al., 2020). However, the initial investment in UAV technology and the need for skilled operators can be barriers to widespread adoption.

Digital Disease Diagnostic Tools

Mobile applications and digital platforms have emerged as practical tools for farmers and agronomists. These tools use machine learning algorithms and vast databases of plant disease images to diagnose diseases from photos taken by smartphones. For instance, applications like Plantix and PlantVillage Nuru offer instant diagnosis and management recommendations for various crop diseases (Barbedo, 2019). While these tools democratize access to diagnostic services, their accuracy depends on the quality and diversity of their training datasets.

The detection and diagnosis of emerging pathogens/diseases in global crop production have evolved significantly, incorporating traditional, molecular, and digital methods. Each approach offers unique advantages and limitations, and an integrated strategy combining these methods is often the most effective in managing plant diseases. As technology continues to advance, the accuracy, speed, and accessibility of pathogen detection are expected to improve, providing better tools to safeguard global food security.

6. Management of emerging diseases

Emerging pathogens/diseases pose a significant threat to global crop production, necessitating effective management and control strategies. These strategies can be categorized into four main approaches: cultural practices, chemical controls, biological control methods, and integrated pest management (IPM). Each of these approaches plays a vital role in mitigating the impact of pathogens on crops.

Cultural Practices

Cultural practices are fundamental in managing crop diseases and involve modifying farming techniques to reduce the prevalence and impact of pathogens. These practices are often the first line of defense and include crop rotation, sanitation, and resistant varieties.

Crop Rotation and Diversification

Crop rotation is the practice of successively planting various crop varieties on the same plot of land. By denying infections their preferred host, this technique upsets their life cycle. For example, rotating cereal crops with legumes can reduce the incidence of soil-borne pathogens like *Fusarium* spp., which cause root rot in cereals (Abawi Widmer, 2000). Crop diversification, including intercropping, can also create a less favorable environment for pathogens and increase biodiversity, which can help in suppressing disease outbreaks (Boudreau, 2013).

Sanitation

Sanitation practices involve removing infected plant debris and maintaining clean fields to reduce the sources of inoculum. This includes the destruction of crop residues, which can harbor pathogens, and cleaning tools and machinery to prevent the spread of diseases (Garrett et al., 2004). Proper disposal of infected plant material is crucial in managing diseases like late blight in potatoes caused by *Phytophthora P. infestans* (Fry, 2008).

Formatted: Font: Not Bold

Formatted: Font: 12 pt, Bold

Commented [GN17]: Disease? (IDM)

Formatted: Highlight

Commented [GN18]: Please provide examples.

Formatted: Font: Italic

Formatted: Font: Italic

Resistant Varieties

The use of disease-resistant crop varieties is a highly effective cultural practice. Breeding programs aim to develop varieties that are resistant to specific pathogens, thereby reducing the need for chemical controls. For instance, the development of rice varieties resistant to bacterial blight (*Xanthomonas X. oryzae* pv. *oryzae*) has significantly reduced yield losses in affected regions (Kumar et al., 2020).

Formatted: Font: Italic

Formatted: Font: Italic

Chemical Controls

Chemical controls involve the application of fungicides, bactericides, and nematicides to manage crop diseases. While effective, these controls must be used judiciously to prevent the development of resistance and minimize environmental impact.

Fungicides

Fungicides are widely used to control fungal pathogens. They can be categorized into protectants, which prevent infection, and eradicants, which eliminate established infections. For example, azoxystrobin and other strobilurins are effective against a broad spectrum of fungal diseases and are commonly used in crops like wheat and grapes (Bartlett et al., 2002). However, the overuse of fungicides can lead to resistance, as seen with the development of resistant strains of powdery mildew (*Erysiphe necator*) in vineyards (McGrath, 2001).

Formatted: Font: Italic

Commented [GN19]: Please use a recent reference.

Bactericides

Bactericides are used to manage bacterial diseases in crops. Copper-based compounds are the most common bactericides and are effective against a range of bacterial pathogens. However, their overuse can result in phytotoxicity and environmental contamination (Adaskaveg Hine, 2001). Newer bactericides, such as antibiotics like streptomycin, are used sparingly to prevent the development of resistant bacterial strains.

Commented [GN20]: Please refer to the above comment

Nematicides

Nematicides control nematode infestations, which can cause significant yield losses. Chemical nematicides, such as fumigants and non-fumigant nematicides, are effective but often have high toxicity and environmental concerns (Desaeger et al., 2020). Therefore, their use is increasingly being supplemented with other management strategies to reduce reliance on chemical nematicides.

Biological Control Methods

Biological control methods involve the use of natural enemies, biopesticides, and beneficial microorganisms to manage crop pathogens. These methods are environmentally friendly and can provide sustainable disease control.

Natural Enemies

Natural enemies, such as predatory insects and parasitic nematodes, can reduce the populations of crop pests and pathogens. For instance, the use of predatory mites to control spider mites in apple orchards has been successful in reducing the need for chemical pesticides (Gerson Weintraub, 2012). These natural enemies help maintain a balanced ecosystem and suppress pest populations.

Formatted: Highlight

Commented [GN21]: Please restrict the discussion with pathogens. The cited reference is about non-insect pest, mites.

Biopesticides

Biopesticides, derived from natural materials, include microbial pesticides, plant extracts, and pheromones. *Bacillus thuringiensis* (Bt) is a well-known microbial pesticide used to control lepidopteran pests in crops like maize and cotton (Sanahuja et al., 2011). Biopesticides are generally specific to target pests and have minimal impact on non-target organisms and the environment.

Formatted: Highlight

Formatted: Highlight

Commented [GN22]: Please see the above comment. Kindly delete or cite references directly related to diseases.

Formatted: Highlight

Beneficial Microorganisms

Beneficial microorganisms, such as mycorrhizal fungi and rhizobacteria, enhance plant health and resistance to pathogens. Arbuscular mycorrhizal fungi (AMF) improve nutrient uptake and can induce

systemic resistance in plants against pathogens (Poza Azcón-Aguilar, 2007). Similarly, plant growth-promoting rhizobacteria (PGPR) like *Pseudomonas fluorescens* can suppress soil-borne diseases through mechanisms such as competition and antibiosis (Weller, 2007).

Integrated Pest Management (IPM)

Integrated Pest Management (IPM) is a holistic approach that combines cultural, chemical, and biological control methods to manage crop diseases in an environmentally and economically sustainable manner. IPM emphasizes the use of multiple tactics to minimize the reliance on any single control method and reduce the risk of resistance development.

Monitoring and Early Detection

Effective IPM begins with regular monitoring and early detection of pests and diseases. Scouting and using diagnostic tools, such as molecular assays, can help identify pathogen presence before outbreaks occur (Larkin et al., 2014). Early detection allows for timely interventions, reducing the need for widespread chemical applications.

Threshold Levels and Decision-Making

IPM involves setting threshold levels for pest populations and disease severity, beyond which control measures are implemented. This decision-making process ensures that interventions are made only when necessary, reducing unnecessary chemical use. For example, action thresholds for aphid populations in wheat can help determine the timing of insecticide applications (Nault Styer, 1972).

Integration of Control Methods

The core of IPM is the integration of multiple control methods. For instance, combining resistant crop varieties with biopesticides and targeted chemical applications can provide effective disease management while minimizing environmental impact. Crop rotation, sanitation, and biological controls can be integrated to manage soil-borne diseases, reducing the need for chemical nematicides (Mazzola, 2007).

Education and Training

Successful IPM implementation requires education and training for farmers and agricultural professionals. Understanding the principles of IPM and the proper use of control methods ensures effective disease management and promotes sustainable agricultural practices (Parsa et al., 2014).

7. CASE STUDIES OF EMERGING PATHOGENS

Wheat Blast (*Magnaporthe oryzae*)

Magnaporthe oryzae, a fungus, causes wheat blast, which is a serious threat to wheat production worldwide. First identified in Brazil in the 1980s, the disease has since spread to other South American countries and recently to Bangladesh, raising concerns about its potential to spread further in Asia and other wheat-producing regions (Islam et al., 2016).

The pathogen infects wheat spikes, leading to the formation of bleached spikelets, which result in substantial yield losses. Under favorable conditions, wheat blast can cause up to 100% crop loss in affected fields (Cruz et al., 2016). The disease thrives in warm and humid climates, which are becoming more common due to climate change, thus increasing the risk of outbreaks.

Efforts to manage wheat blast include the use of fungicides, development of resistant wheat varieties, and agronomic practices such as crop rotation and the removal of infected crop residues. However, the pathogen's ability to rapidly adapt and overcome genetic resistance in wheat varieties remains a challenge (Islam et al., 2016). Moreover, the lack of effective resistance genes in the wheat gene pool necessitates ongoing research and international collaboration to develop sustainable management strategies.

Cassava Mosaic Disease (Cassava Mosaic Viruses)

Cassava mosaic disease (CMD) is caused by a complex of cassava mosaic geminiviruses (CMGs), including African cassava mosaic virus (ACMV) and East African cassava mosaic virus (EACMV).

Commented [GN23]: Enormous work has been done in this area. Please quote recent work.

Formatted: Font: Italic

Commented [GN24]: Kindly modify as Integrated Disease Management with examples of disease

Formatted: Highlight

Formatted: Highlight

Commented [GN25]: Please include only relevant information .

Commented [GN26]: Please provide examples of IDM.

Commented [GN27]: Repetitive; can be combined with Examples of emerging diseases "under overview in page 3.

Formatted: Highlight

CMD is a major constraint to cassava production in Africa and parts of Asia, leading to significant food security concerns due to cassava's role as a staple crop for millions of people (Legg et al., 2015).

CMD symptoms include leaf mosaic, distortion, and stunting, which severely reduce photosynthetic capacity and root yield. In some regions, CMD can cause yield losses of up to 90%, severely impacting the livelihoods of smallholder farmers (Thresh Cooter, 2005). The disease spreads through infected cuttings and by whiteflies (*Bemisia tabaci*), which transmit the virus between plants.

Management of CMD involves the use of virus-free planting material, resistant cassava varieties, and integrated pest management to control whitefly populations. Breeding programs have successfully developed CMD-resistant varieties, but the rapid evolution of CMGs and the emergence of new virus strains pose ongoing challenges (Legg et al., 2015). Therefore, continuous monitoring and adaptive management strategies are crucial to mitigate the impact of CMD on cassava production.

Xylella fastidiosa in Olive Trees

Xylella fastidiosa is a bacterium that causes significant diseases in a wide range of plant species, including olives, grapes, citrus, and almonds. In olive trees, it is responsible for Olive Quick Decline Syndrome (OQDS), which has devastated olive orchards in Southern Italy since its identification in 2013 (Saponari et al., 2019).

The pathogen colonizes the xylem vessels of the host plant, causing blockages that lead to water stress, leaf scorch, and eventual tree death. The rapid spread of *Xylella fastidiosa* is facilitated by xylem-feeding insects, such as the spittlebug (*Philaenus spumarius*), which transmit the bacterium from infected to healthy plants (EFSA, 2018).

Control measures for *Xylella fastidiosa* in olive trees are limited due to the lack of effective chemical treatments. Current strategies focus on preventing the spread of the bacterium through the removal of infected trees, controlling vector populations, and implementing quarantine measures to restrict the movement of potentially infected plant material (Saponari et al., 2019). Additionally, research is ongoing to identify resistant olive varieties and develop innovative management approaches.

Banana Fusarium Wilt (*Fusarium oxysporum f.sp. cubense*)

Banana Fusarium wilt, also known as Panama disease, is caused by the soil-borne fungus *Fusarium oxysporum f.sp. cubense* (Foc). The most virulent strain, Tropical Race 4 (TR4), has emerged as a significant threat to global banana production, particularly affecting the Cavendish variety, which dominates the international banana trade (Ploetz, 2015).

Banana leaf drooping and yellowing due to fusarium wilt results in plant mortality. Once established in a plantation, the fungus is very difficult to remove since it can linger in the soil for decades (Ploetz, 2015). Concerns regarding the possibility of a global banana crisis have been raised since TR4 was originally discovered in Southeast Asia and has subsequently moved to the Middle East, Africa, and Latin America.

Table 4: Case Studies of Emerging Pathogens

Pathogen	Crop	Location	Impact	Response
Wheat Blast (<i>Magnaporthe oryzae</i>)	Wheat	South America, South Asia	Severe yield losses, up to 100% in fields	Development of resistant varieties, improved agronomic practices
Cassava Mosaic Disease (Cassava Mosaic Viruses)	Cassava	Africa, Asia	Stunted growth, significant yield reduction	Breeding for resistant varieties, virus-free planting material
<i>Xylella fastidiosa</i> in Olive Trees	Olive	Europe (Italy, Spain), Brazil	Rapid decline and death of olive trees	Quarantine measures, removal of infected trees, vector control

Formatted: Font: Italic

Commented [GN28]: Please refer to the comment on Page 3 on **Examples of emerging diseases**; kindly present salient points with many references, as this is a review. Kindly italicize scientific names.

Commented [GN29]: Repetitive. Please combine with Table 1. The same information is presented elsewhere in the text too.

Formatted: Highlight

Banana Fusarium Wilt (<i>Fusarium oxysporum</i> f.sp. <i>cubense</i>)	Banana	Asia, Africa, Latin America	Widespread devastation of banana plantations	Breeding resistant varieties, soil management practices
---	--------	-----------------------------	--	---

Commented [GN30]: Kindly italicize scientific names.

Management strategies for Fusarium wilt include the use of resistant banana varieties, soil fumigation, and strict quarantine measures to prevent the spread of the fungus. However, the genetic uniformity of the Cavendish banana and the long-lasting nature of the pathogen in the soil pose significant challenges. Breeding programs and biotechnological approaches are being explored to develop TR4-resistant banana cultivars and improve disease management practices (Ploetz, 2015).

8. INNOVATIVE APPROACHES AND FUTURE PROSPECTS

Genetic Resistance and Breeding

Development of Resistant Varieties

The development of crop varieties resistant to emerging pathogens is a fundamental approach in combating the threats to global crop production. Traditional breeding techniques, although time-consuming, have historically played a significant role in improving crop resistance. The introduction of resistant genes from wild relatives of crops through conventional crossbreeding has been successful in many cases. For instance, the transfer of resistance genes from wild wheat species has provided wheat with durable resistance against rust diseases (Figuroa et al., 2018).

Commented [GN31]: This is a regular method. Please combine with **Resistant Varieties** in page 9.

Table 5: Research on Resistant Crop Varieties

Crop	Resistant Varieties	Pathogen Targeted	Research Institution	Year Released
Wheat	Norman, CIMMYT-84	Rust (<i>Puccinia graminis</i>)	International Maize and Wheat Improvement Center (CIMMYT)	2018
Potato	SpuntaG2	Late blight (<i>Phytophthora infestans</i>)	International Potato Center (CIP)	2017
Rice	IR64, BRRI dhan29	Bacterial blight (<i>Xanthomonas oryzae</i>)	International Rice Research Institute (IRRI)	2016
Banana	GCTCV-218, FHIA-25	Fusarium Wilt (<i>Fusarium oxysporum</i> f.sp. <i>cubense</i>)	Bioversity International	2019
Cassava	KU50, TMS 30572	Cassava Mosaic Disease (Cassava Mosaic Viruses)	International Institute of Tropical Agriculture (IITA)	2015
Tomato	Hawaii 7996	Bacterial wilt (<i>Ralstonia solanacearum</i>)	Asian Vegetable Research and Development Center (AVRDC)	2020

Commented [GN32]: The table may be removed and only the important information can be produced in four lines: for eg., the resistant varieties released for the managing a particular disease and presented under **Resistant Varieties** in page 9.

Formatted: Highlight

Commented [GN33]: Throughout the text italicization of scientific name is to be done.

Marker-Assisted Selection (MAS)

Marker-assisted selection (MAS) has revolutionized the breeding process by allowing for the identification and selection of desirable traits at the genetic level. This technique speeds up the breeding

process and enhances the precision of developing resistant varieties. MAS has been successfully employed in rice breeding to develop varieties resistant to bacterial blight by identifying markers linked to resistance genes (Singh et al., 2021).

Commented [GN34]: Please include many references

Genomic Selection

Genomic selection, which uses genome-wide markers to predict the performance of breeding lines, is another promising approach. This method allows for the selection of superior genotypes early in the breeding cycle, thus reducing the time required to develop resistant varieties. In maize, genomic selection has accelerated the development of lines resistant to multiple pathogens, significantly boosting yields and resilience (Crossa et al., 2017).

Commented [GN35]: This is a regular method. Please combine with Resistant Varieties in page 9.

CRISPR and Gene Editing Technologies

CRISPR-Cas9 Technology

CRISPR-Cas9 has emerged as a groundbreaking tool in plant biotechnology, offering precise and efficient editing of plant genomes. This technology allows for the targeted modification of genes associated with disease resistance. For example, CRISPR-Cas9 has been used to knock out susceptibility genes in rice, thereby conferring resistance to bacterial blight (Zhou et al., 2015).

Commented [GN36]: So much work has been done on CRISPR. Please include a few more references.

Multiplex Gene Editing

The ability to edit multiple genes simultaneously (multiplex gene editing) using CRISPR technology is particularly advantageous in developing durable resistance. This approach can be used to modify several genes involved in the plant's defense mechanisms, creating crops that are resistant to a wide range of pathogens. In tomatoes, multiplex CRISPR editing has successfully targeted multiple genes to enhance resistance to various fungal pathogens (Tian et al., 2018).

Future Prospects of Gene Editing

Looking ahead, the integration of CRISPR-Cas9 with other emerging technologies, such as synthetic biology and advanced bioinformatics, holds tremendous potential. These advancements could enable the creation of crops with enhanced resistance to new and evolving pathogens. Furthermore, as CRISPR technology becomes more refined and accessible, it is expected to play a pivotal role in ensuring food security in the face of climate change and pathogen pressure (Rodriguez et al., 2018).

Microbiome Management

Harnessing Plant Microbiomes

The plant microbiome, consisting of bacteria, fungi, and other microorganisms, plays a crucial role in plant health and disease resistance. Innovative approaches are being developed to manipulate the plant microbiome to enhance crop resistance. For example, inoculating plants with beneficial microbes that can outcompete or inhibit pathogens has shown promise in various crops (Berendsen et al., 2018).

Bioinoculants

Bioinoculants, which are preparations containing beneficial microorganisms, are being increasingly utilized to boost plant health and resistance. These bioinoculants can be applied to seeds, soil, or plant surfaces to establish beneficial microbial communities. In soybean, the use of rhizobial inoculants has improved resistance to soil-borne pathogens and enhanced nitrogen fixation, leading to better growth and yields (Mendes et al., 2013).

Engineering the Microbiome

Advancements in microbiome engineering, such as the development of synthetic microbial communities tailored to specific crops, represent a cutting edge approach. By designing microbial consortia that can enhance plant growth and resilience to pathogens, researchers aim to create a sustainable and environmentally friendly method of crop protection. This approach has shown potential in enhancing disease resistance in crops like wheat and maize (Liu et al., 2019).

Policy and Regulatory Frameworks

Supporting Innovation in Agriculture

Effective policy and regulatory frameworks are essential to support the adoption and scaling of innovative approaches in crop protection. ~~Governments and international bodies must create conducive environments that encourage research, development, and deployment of new technologies.~~ Policies that provide funding for agricultural research and offer incentives for private sector investment in biotechnologies are crucial (Altieri et al., 2018).

Regulation of Gene Edited Crops

The regulatory landscape for gene-edited crops varies widely across different countries. ~~Harmonizing regulations to facilitate the global trade of gene edited crops while ensuring safety and transparency is a key challenge.~~ Clear guidelines and risk assessment frameworks are needed to address public concerns and ensure that gene-edited crops are developed and deployed responsibly (Pardey et al., 2016).

Sustainable Agricultural Practices

Promoting sustainable agricultural practices through policy measures is vital for long-term crop health and productivity. Policies that encourage integrated pest management (IPM), crop rotation, and the use of biocontrol agents can help reduce reliance on chemical pesticides and mitigate the impact of pathogens. Collaborative efforts between policymakers, researchers, and farmers are essential to implement these practices effectively (Tilman et al., 2011).

The fight against emerging ~~pathogens-diseases~~ in global crop production requires a multifaceted approach involving genetic resistance, advanced gene editing technologies, microbiome management, and supportive policy frameworks. By leveraging these innovative strategies, we can enhance crop resilience, ensure food security, and sustainably meet the demands of a growing global population. Continued investment in research and development, along with international cooperation, will be crucial in addressing the challenges posed by emerging pathogens.

Commented [GN37]: Repetitive of the text elsewhere in the MS.

9. GLOBAL COLLABORATION AND RESEARCH INITIATIVES

International Organizations and Programs

Food and Agriculture Organization (FAO)

~~In order to combat the threats that new pathogens pose to global crop production, the FAO is essential.~~ The FAO establishes global standards for plant health through its many programmes, such as the International Plant Protection Convention (IPPC), and promotes collaboration among member nations to stop the introduction and spread of pests and diseases (FAO, 2021). ~~In order to help developing nations develop their capacity for plant health management, the FAO also carries out a great deal of research and offers technical assistance to such nations.~~

Consultative Group on International Agricultural Research (CGIAR)

~~CGIAR(Consultative Group on International Agricultural Research) is a global research partnership that aims to reduce rural poverty, increase food security, improve human health and nutrition, and ensure sustainable management of natural resources. Its research programs focusses on major crops and agricultural systems, addressing the challenges posed by emerging pathogens-diseases through integrated pest management, crop improvement, and biosecurity measures (CGIAR, 2022). By collaborating with national agricultural research systems and other stakeholders, CGIAR facilitates the dissemination of knowledge and technologies to mitigate the impact of pathogens on crop production.~~

Formatted: Font: Not Bold

Commented [GN38]: Please quote examples for their contributions in emerging diseases, similar to what is done by IITA.

International Institute of Tropical Agriculture (IITA)

~~The IITA (International Institute of Tropical Agriculture) is dedicated to enhancing food security and reducing poverty in tropical regions by improving agricultural productivity. The IITA conducts research on emerging plant pathogens-diseases and develops innovative solutions to combat them. For example, their work on cassava diseases, such as cassava brown streak disease and cassava mosaic disease, has led to the development of disease-resistant varieties and improved management practices (IITA, 2020). The IITA collaborates with various international and national organizations to ensure the widespread adoption of these solutions.~~

Formatted: Font: Not Bold

Formatted: Font: Bold

Collaborative Research Networks

The Global Plant Pathology Network (GPPN)

The GPPN The Global Plant Pathology Network is a collaborative platform that brings together plant pathologists from around the world to share knowledge, resources, and expertise. This network facilitates the exchange of information on emerging pathogens, promotes joint research projects, and provides training opportunities for researchers. By fostering a collaborative environment, the GPPN helps to accelerate the development of effective strategies to manage plant diseases and protect global crop production (GPPN, 2021).

The Plantwise Knowledge Bank

Managed by CABI, the Plantwise Knowledge Bank is an online resource that provides diagnostic tools, treatment advice, and distribution maps for plant pests and diseases. The platform enables farmers, extension workers, and researchers to access up-to-date information on emerging pathogens and their management. Through its network of plant clinics, Plantwise also facilitates the collection and sharing of field data, which can be used to monitor disease outbreaks and inform research efforts (CABI, 2022).

The International Society for Plant Pathology (ISPP)

The ISPP (The International Society for Plant Pathology) promotes the global exchange of knowledge and expertise in plant pathology. It organizes international congresses, publishes research journals, and supports working groups focused on specific plant diseases and pathogens. By providing a platform for collaboration and communication, the ISPP helps to advance the science of plant pathology and improve the management of emerging pathogens (ISPP, 2021).

Formatted: Font: Not Bold

Funding and Support Mechanisms

The Bill Melinda Gates Foundation

The Bill Melinda Gates Foundation provides significant funding for agricultural research and development, including projects aimed at combating emerging plant pathogens. The foundation supports initiatives that develop disease-resistant crop varieties, improve diagnostic tools, and enhance surveillance systems. By funding for collaborative research projects and capacity-building programs, the foundation helps to strengthen global efforts to protect crop production from emerging threats (Bill Melinda Gates Foundation, 2022).

The Global Environment Facility (GEF)

The GEF (Global Environment Facility) funds projects that address global environmental challenges, including those related to agriculture and food security. Through its funding mechanisms, the GEF supports initiatives that aim to prevent the spread of invasive species, improve pest and disease management, and enhance the resilience of agricultural systems to emerging pathogens. By partnering with international organizations and research institutions, the GEF helps to mobilize resources and knowledge to address the impact of plant diseases on global crop production (GEF, 2021).

Formatted: Font: Not Bold

Horizon Europe

Horizon Europe is the European Union's key funding program for research and innovation. It supports projects that address societal challenges, including those related to agriculture and food security. Through its funding calls, Horizon Europe provides financial support for collaborative research projects that develop innovative solutions to manage emerging plant pathogens. By fostering for partnerships between research institutions, industry, and other stakeholders, Horizon Europe helps to drive for the development and implementation of effective disease management strategies (European Commission, 2021).

These references provide detailed insights into the various international organizations, research networks, and funding mechanisms that are crucial in the fight against emerging plant pathogens. Through global collaboration and support, these initiatives help to safeguard crop production and ensure food security worldwide.

10. CONCLUSIONS

Summary of Key Points

Emerging pathogens diseases pose a significant threat to global crop production, with the potential to cause severe economic and food security impacts. Pathogens such as fungi, bacteria, viruses, and nematodes have been identified as critical agents of crop diseases, and their emergence is often linked to changes in agricultural practices, climate change, and global trade.

Emergence and Spread of Pathogens: The rise in global travel and trade has facilitated the rapid spread of pathogens across borders, leading to outbreaks in regions previously unaffected. For example, the fungal pathogen *Fusarium oxysporum* f. sp. *cubense*, responsible for Panama disease in bananas, has spread from Asia to the Americas, threatening banana production worldwide (Ploetz, 2006).

Impact on Major Crops: Key staple crops such as wheat, rice, maize, and potatoes are particularly vulnerable. Wheat rust, caused by *Puccinia graminis* f. sp. *tritici*, has led to significant yield losses in Africa and the Middle East (Singh et al., 2011). Similarly, the rice blast fungus *Magnaporthe oryzae* has had devastating effects on rice yields in Asia and Africa (Dean et al., 2012).

Climate Change Effects: Climate change exacerbates the problem by creating favorable conditions for pathogen proliferation and altering the geographic distribution of both crops and pathogens. Higher temperatures and increased humidity can enhance the growth and spread of fungal and bacterial pathogens (Anderson et al., 2004).

Economic and Food Security Implications: Crop losses due to emerging pathogens result in reduced food availability and higher prices, impacting both local economies and global markets. The economic burden of managing these diseases and the resulting yield losses can be substantial, affecting farmers' livelihoods and national economies (Strange Scott, 2005).

Future Directions

To effectively combat emerging pathogens diseases, it is essential to invest in research and develop integrated management strategies that are adaptive and resilient.

Advancement in Genomics and Biotechnology: ~~Leveraging genomic technologies can aid in the early detection and identification of pathogens, allowing for timely interventions.~~ Genetic engineering and CRISPR technology hold promise for developing pathogen-resistant crop varieties (Gao, 2018).

Integrated Pest Management (IPM): A holistic approach that combines biological, cultural, mechanical, and chemical control methods can reduce reliance on pesticides and enhance sustainability. This includes the use of biocontrol agents, crop rotation, and resistant crop varieties (Ehler, 2006).

Climate Resilient Agricultural Practices: Adapting agricultural practices to mitigate the impacts of climate change is crucial. This includes developing drought-resistant crop varieties, implementing water-efficient irrigation systems, and adopting agroforestry practices (Altieri et al., 2015).

Global Surveillance and Data Sharing: Establishing a global surveillance system for crop pathogens can facilitate rapid response and containment. Collaborative efforts among countries and organizations to share data and resources can enhance preparedness and response strategies (Savary et al., 2019).

RECOMMENDATIONS FOR STAKEHOLDERS

Stakeholders, including policymakers, researchers, farmers, and industry players, must collaborate to address the challenges posed by emerging pathogens.

Policymakers:

- Regulatory Frameworks: Develop and enforce regulations that prevent the spread of pathogens through trade and travel. This includes quarantine measures and the monitoring of imported plant materials (McLeod et al., 2010).
- Funding and Support: Allocate funds for research and development of disease-resistant crops and sustainable agricultural practices. Support extension services to educate farmers on the latest management techniques (Oerke, 2006).

Researchers:

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Commented [GN39]: Please modify it as IDM

Formatted: Font: (Default) Times New Roman

Formatted: List Paragraph, Bulleted + Level: 1 + Aligned at: 0.25" + Indent at: 0.5"

- ~~Interdisciplinary Research:~~ Encourage collaboration across disciplines, including plant pathology, climatology, and economics, to develop comprehensive solutions. Focus on understanding the mechanisms of ~~pathogen disease~~ emergence and spread (Fisher et al., 2012).
- ~~Technology Transfer:~~ Facilitate the transfer of new technologies and practices to farmers, ensuring they are practical and cost-effective (Savary et al., 2019).

Formatted: Font: (Default) Times New Roman

Formatted: List Paragraph, Bulleted + Level: 1 + Aligned at: 0.25" + Indent at: 0.5"

Formatted: Font: (Default) Times New Roman

Farmers:

- ~~Adoption of Best Practices:~~ Implement integrated pest management practices and adopt resistant crop varieties. Stay informed about emerging threats and participate in training programs (Pretty, 2008).
- ~~Community Engagement:~~ Work collectively with local and national agricultural organizations to monitor and manage pathogen outbreaks. Share knowledge and resources within the farming community (Altieri et al., 2015)*.

Formatted: Font: (Default) Times New Roman

Formatted: List Paragraph, Bulleted + Level: 1 + Aligned at: 0.25" + Indent at: 0.5"

Industry Players:

- ~~Sustainable Solutions:~~ Invest in the development of sustainable agricultural products and technologies. Promote environmentally friendly pesticides and biocontrol agents (Popp et al., 2013)*.
- ~~Corporate Responsibility:~~ Engage in corporate social responsibility initiatives that support farmers and communities affected by crop diseases. Collaborate with governments and NGOs to improve agricultural resilience (Strange Scott, 2005)*.

Formatted: Font: (Default) Times New Roman

Formatted: List Paragraph, Bulleted + Level: 1 + Aligned at: 0.25" + Indent at: 0.5"

In conclusion, addressing the challenges posed by emerging ~~pathogens-diseases~~ requires a multifaceted approach that includes scientific innovation, practical management strategies, and collaborative efforts across all stakeholders. By prioritizing research, implementing sustainable practices, and fostering global cooperation, it is possible to mitigate the impact of these ~~pathogens-diseases~~ on global crop production and ensure food security for future generations.

References

Agrios, G. N. (2005). *Plant Pathology* (5th ed.). Elsevier Academic Press.

Alicai, T., Omongo, C. A., Maruthi, M. N., Hillocks, R. J., Baguma, Y., Kawuki, R., ... Colvin, J. (2007). Re-emergence of cassava brown streak disease in Uganda. *Plant Disease*, 91(1), 24-29.

Commented [GN40]: ???

Altieri, M. A., Nicholls, C. I., Henao, A., Lana, M. A. (2015). Agroecology and the design of climate change-resilient farming systems. *Agronomy for Sustainable Development*, 35(3), 869-890.

Álvarez, B., Biosca, E. G., López, M. M. (2010). On the life of *Ralstonia solanacearum*, a destructive bacterial plant pathogen. In *Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology* (Vol. 2, pp. 267-279).

Anderson, P. K., Cunningham, A. A., Patel, N. G., Morales, F. J., Epstein, P. R., Daszak, P. (2004). Emerging infectious diseases of plants: pathogen pollution, climate change and agrotechnology drivers. *Trends in Ecology Evolution*, 19(10), 535-544.

Anderson, P. K., Cunningham, A. A., Patel, N. G., Morales, F. J., Epstein, P. R., Daszak, P. (2004). Emerging infectious diseases of plants: pathogen pollution, climate change and agrotechnology drivers. *Trends in Ecology Evolution*, 19(10), 535-544.

Commented [GN41]: Repeated. Pl. delete

Barbedo, J. G. A. (2019). Plant disease identification from individual lesions and spots using deep learning. *Biosystems Engineering*, 180, 96-107.

Bebber, D. P., Holmes, T., Gurr, S. J. (2013). The global spread of crop pests and pathogens. *Global Ecology and Biogeography*, 23(12), 1398-1407.

Bebber, D. P., Holmes, T., Gurr, S. J. (2013). The global spread of crop pests and pathogens. *Global Ecology and Biogeography*, 23(12), 1398-1407.

Commented [GN42]: Repeated. Please delete.

Bill Melinda Gates Foundation. (2022). Agricultural Development. Retrieved from Gates Foundation

Blomme, G., Ploetz, R., Jones, D. R., De Langhe, E., Price, N., Gold, C., ... Pillay, M. (2017). A historical overview of the appearance and spread of banana bunchy top disease and the involvement of BBTv in the demise of the Gros Michel banana. *Virology*, 497, 211-217.

Bové, J. M. (2006). Huanglongbing: a destructive, newly-emerging, century-old disease of citrus. *Journal of Plant Pathology*, 88(1), 7-37.

Burdon, J. J., Zhan, J. (2020). Climate change and disease in plant communities. *PLoS Biology*, 18(5), e3000949.

CABI. (2022). Plantwise Knowledge Bank. Retrieved from CABI

CGIAR. (2022). Research. Retrieved from CGIAR

Chakraborty, S., Newton, A. C. (2011). Climate change, plant diseases, and food security: An overview. *Plant Pathology*, 60(1), 2-14.

Chakraborty, S., Murray, G. M., Magarey, P. A., Yonow, T., O'Brien, R. G., Croft, B. J., ... Dudzinski, M. J. (2000). Potential impact of climate change on plant diseases of economic significance to Australia. *Australasian Plant Pathology*, 29(3), 143-176.

Chakraborty, S., Newton, A. C. (2011). Climate change, plant diseases and food security: an overview. *Plant Pathology*, 60(1), 2-14.

Clark, M. F., Adams, A. N. (1977). Characteristics of the microplate method of enzyme-linked immunosorbent assay for the detection of plant viruses. *Journal of General Virology*, 34(3), 475-483.

Dean, R., Van Kan, J. A. L., Pretorius, Z. A., Hammond-Kosack, K. E., Di Pietro, A., Spanu, P. D., ... Foster, G. D. (2012). The Top 10 fungal pathogens in molecular plant pathology. *Molecular Plant Pathology*, 13(4), 414-430.

Dean, R., Van Kan, J. A., Pretorius, Z. A., Hammond-Kosack, K. E., Di Pietro, A., Spanu, P. D., ... Foster, G. D. (2012). The Top 10 fungal pathogens in molecular plant pathology. *Molecular Plant Pathology*, 13(4), 414-430.

EFSA Panel on Plant Health. (2015). Scientific opinion on the risks to plant health posed by *Xylella fastidiosa* in the EU territory, with the identification and evaluation of risk reduction options. *EFSA Journal*, 13(1), 3989.

Ehler, L. E. (2006). Integrated pest management (IPM): definition, historical development and implementation, and the other IPM. *Pest Management Science: formerly Pesticide Science*, 62(9), 787-789.

European Commission. (2021). Horizon Europe. Retrieved from European Commission

European Food Safety Authority (EFSA). (2019). Update of the *Xylella* spp. host plant database – systematic literature searches up to 30 June 2019. *EFSA Journal*, 17(10), e05879.

European Food Safety Authority. (2015). Scientific opinion on the risks to plant health posed by *Xylella fastidiosa* in the EU territory, with the identification and evaluation of risk reduction options. *EFSA Journal*, 13(1), 3989.

FAO. (2020). The State of Food Security and Nutrition in the World 2020. Food and Agriculture Organization of the United Nations. Retrieved from <http://www.fao.org/publications/sofi/2020/en/>

FAO. (2021). International Plant Protection Convention. Retrieved from FAO

Fisher, M. C., Henk, D. A., Briggs, C. J., Brownstein, J. S., Madoff, L. C., McCraw, S. L., Gurr, S. J. (2012). Emerging fungal threats to animal, plant and ecosystem health. *Nature*, 484(7393), 186-194.

Franceschini, M. D., Lorrain, V., Arzul, L., Sechet, V. (2019). Seed-borne pathogens: Detection, identification, and control. In *Plant Pathology* (pp. 99-126). Springer, Cham.

Commented [GN43]: Please provide the link, if it is website source.

Commented [GN44]: ???

Commented [GN45]: Please provide the link.

Commented [GN46]: Please provide the link

Commented [GN47]: ??

Commented [GN48]: Repeated; please delete

Commented [GN49]: Repeated. Please remove.

Formatted: Font: Italic

Commented [GN50]: Please provide the link.

Formatted: Font: Italic

Formatted: Font: Italic

Fry, W. E. (2008). *Phytophthora infestans*: the plant (and R gene) destroyer. *Molecular Plant Pathology*, 9(3), 385-402.

Formatted: Font: Italic

Gao, C. (2018). The future of CRISPR technologies in agriculture. *Nature Reviews Molecular Cell Biology*, 19(5), 275-276.

GEF. (2021). Agriculture and Food Security. Retrieved from GEF

Commented [GN51]: Please provide the link.

GPPN. (2021). Global Plant Pathology Network. Retrieved from GPPN

Commented [GN52]: Please provide the link.

IITA. (2020). Research Themes. Retrieved from IITA

Commented [GN53]: Please provide the link.

Islam, M. T., Croll, D., Gladieux, P., Soanes, D. M., Persoons, A., Bhattacharjee, P., ... Stukenbrock, E. H. (2016). Emergence of wheat blast in Bangladesh was caused by a South American lineage of *Magnaporthe oryzae*. *BMC Biology*, 14(1), 84.

Commented [GN54]: ???

ISPP. (2021). International Society for Plant Pathology. Retrieved from ISPP

Formatted: Font: Italic

Commented [GN55]: Please provide the link.

Jones, J. T., Haegeman, A., Danchin, E. G., Gaur, H. S., Helder, J., Jones, M. G., ... Kikuchi, T. (2013). Top 10 plant-parasitic nematodes in molecular plant pathology. *Molecular Plant Pathology*, 14(9), 946-961.

Commented [GN56]: Please share the link

Legg, J. P., Jeremiah, S. C., Obiero, H. M., Maruthi, M. N., Ndyetabula, I., Okao-Okuja, G., ... Thresh, J. M. (2014). Comparing the regional epidemiology of the cassava mosaic and cassava brown streak virus pandemics in Africa. *Virus Research*, 186, 61-70.

Lucas, J. A., Hawkins, N. J., Fraaije, B. A. (2015). The evolution of fungicide resistance. *Advances in Applied Microbiology*, 90, 29-92.

Massart, S., Olmos, A., Jijakli, M. H., Candresse, T. (2014). Current impact and future directions of high throughput sequencing in plant virus diagnostics. *Virus Research*, 188, 90-96.

McDonald, B. A., Stukenbrock, E. H. (2016). Rapid emergence of pathogens in agro-ecosystems: global threats to agricultural sustainability and food security. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371(1709), 20160026.

McLeod, A., Jalaluddin, A., Coetzee, V., Botha, W. (2010). Financial impact of quarantine risks and regulation of agricultural trade. *Agrekon*, 49(4), 426-445.

Mumford, R., Macarthur, R., Boonham, N. (2016). The role and challenges of new diagnostic technology in plant biosecurity. *Food Security*, 8(1), 103-109.

Nichols, R. F. W. (1950). The brown streak disease of cassava: distribution, climatic effects and diagnostic symptoms. *East African Agricultural Journal*, 15(3), 154-160.

Nicol, J. M., Turner, S. J., Coyne, D. L., Nijs, L. D., Hockland, S., Tahna Maafi, Z. (2011). Current nematode threats to world agriculture. *Genomics and Molecular Genetics of Plant-Nematode Interactions*, 21-43.

Notomi, T., Okayama, H., Masubuchi, H., Yonekawa, T., Watanabe, K., Amino, N., Hase, T. (2000). Loop-mediated isothermal amplification of DNA. *Nucleic Acids Research*, 28(12), E63.

Oerke, E. C. (2006). Crop losses to pests. *The Journal of Agricultural Science*, 144(1), 31-43.

Panno, S., Caruso, A. G., Davino, S., Bertacca, S. (2019). Tomato brown rugose fruit virus: Seed transmission rate and efficacy of different seed-disinfection treatments. *Plants*, 8(12), 496.

Ploetz, R. C. (2006). Fusarium wilt of banana is caused by several pathogens referred to as *Fusarium oxysporum* f. sp. *cubense*. *Phytopathology*, 96(6), 653-656.

Ploetz, R. C. (2015). Fusarium wilt of banana. *Phytopathology*, 105(12), 1512-1521.

Ploetz, R. C. (2015). Management of Fusarium wilt of banana: A review with special reference to tropical race 4. *Crop Protection*, 73, 7-15.

Commented [GN57]: Ploetz (2015) a &b?

Popp, J., Pető, K., ~~Nagy, Nagy~~, J. (2013). Pesticide productivity and food security. A review. *Agronomy for Sustainable Development*, 33(1), 243-255.

Pretty, J. (2008). Agricultural sustainability: concepts, principles and evidence. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1491), 447-465.

Ramegowda, V., ~~Senthil, Senthil~~-Kumar, M. (2015). The interactive effects of simultaneous biotic and abiotic stresses on plants: Mechanistic understanding from drought and pathogen combination. *Journal of Plant Physiology*, 176, 47-54.

Ristaino, J. B., Groves, C. T., ~~Parra, Parra~~, G. (2001). PCR amplification of the Irish potato famine pathogen from historic specimens. *Nature*, 411(6838), 695-697.

Rumpf, T., Mahlein, A. K., Steiner, U., Oerke, E. C., Dehne, H. W., ~~Plümer, Plümer~~, L. (2010). Early detection and classification of plant diseases with support vector machines based on hyperspectral reflectance. *Computers and Electronics in Agriculture*, 74(1), 91-99.

Salem, N., Mansour, A., Ciuffo, M., Falk, B. W., Turina, M. (2016). A new Tobamovirus infecting tomato crops in Jordan. *Archives of Virology*, 161(2), 503-506.

Saponari, M., Giampetruzzi, A., Loconsole, G., Boscia, D., Saldarelli, P. (2019). *Xylella fastidiosa* in olive in Apulia: where we stand. *Phytopathology*, 109(2), 175-186.

Savary, S., Willocquet, L., Pethybridge, S. J., Esker, P., McRoberts, N., ~~Nelson, Nelson~~, A. (2019). The global burden of pathogens and pests on major food crops. *Nature Ecology & Evolution*, 3(3), 430-439.

Schena, L., Duncan, J. M., ~~Cooke, Cooke~~, D. E. (2013). Development and application of a PCR-based 'molecular tool box' for the identification of *Phytophthora* species damaging forests and natural ecosystems. *Plant Pathology*, 62(4), 928-937.

Schumann, G. L., -D'Arcy, C. J. (2006). *Essential Plant Pathology*. American Phytopathological Society Press.

Singh, R. P., Hodson, D. P., Jin, Y., Huerta-Espino, J., Kinyua, M. G., Wanyera, R., ... Njau, P. (2011). The emergence of Ug99 races of the stem rust fungus is a threat to world wheat production. *Annual Review of Phytopathology*, 49, 465-481.

Singh, R. P., Hodson, D. P., Jin, Y., Lagudah, E. S., Ayliffe, M. A., Bhavani, S., ... Njau, P. (2016). Emergence and spread of new races of wheat stem rust fungus: continued threat to food security and prospects of genetic control. *Phytopathology*, 106(2), 135-144.

Strange, R. N., ~~Scott, Scott~~, P. R. (2005). Plant disease: a threat to global food security. *Annual Review of Phytopathology*, 43, 83-116.

Ullstrup, A. J. (1972). The impacts of the southern corn leaf blight epidemics of 1970-1971. *Annual Review of Phytopathology*, 10(1), 37-50.

United Nations. (2019). *World Population Prospects 2019: Highlights*. Department of Economic and Social Affairs, Population Division. Retrieved from https://population.un.org/wpp/Publications/Files/WPP2019_Highlights.pdf

Vanneste, J. L., Yu, J., Cornish, D. A., Tanner, D. J., Windner, R., Chapman, J. R., ... Poulter, R. T. (2011). Identification, virulence, and distribution of *Pseudomonas syringae* pv. *actinidiae* isolates from New Zealand kiwifruit orchards. *Plant Disease*, 95(11), 1358-1366.

World Bank. (2021). *Employment in agriculture (% of total employment) (modeled ILO estimate)*. Retrieved from <https://data.worldbank.org/indicator/SL.AGR.EMPL.ZS>

Wu, F. (2006). Mycotoxin reduction in Bt corn: potential economic, health, and regulatory impacts. *Transgenic Research*, 15(3), 277-289.

Commented [GN58]: <<<

Commented [GN59]: ???

Formatted: Font: Italic

Formatted: Font: Italic

Zhang, J., Pu, R., Wang, J., Huang, W., Yuan, L., Luo, J. (2020). Monitoring powdery mildew of winter wheat by using UAV-based hyperspectral imaging. *International Journal of Applied Earth Observation and Geoinformation*, 90, 102126.

UNDER PEER REVIEW