

Does upper Arm length, Arm Span, and foot length serve as good predictors for stature? A cross-sectional study among Northern Nigerians.

Abstract

Background: Stature estimation is an important identification process necessary when dissembled body parts are found. The study is important in the field of physical and forensic anthropology and for industrial design. The study aims to determine if upper arm length, foot length, and arm span could be a good predictor for stature among northern Nigerians.

Method: The study adopted a descriptive cross-sectional design using a multi-stage random sampling technique. 400 northern Nigerians (200 males and 200 females) were involved in the research and data were collected via direct anthropometric method using a stadiometer, and meter rule. The data obtained were analyzed using the IBM SPSS and probability was stated ($p < 0.05$).

Results: The findings of the study show that males have a mean value for the standing height, upper arm length, foot length, and arm span as 171.97 ± 6.68 cm, 32.62 ± 2.55 cm, 25.76 ± 1.17 cm, 181.72 ± 7.83 cm respectively, and females were 163.33 ± 5.96 cm, 35.23 ± 2.19 cm, 23.04 ± 1.51 cm, and 168.67 ± 6.68 cm for standing height, upper arm length, foot length, and arm span which has shown sexual differences and age-related differences ($p < 0.05$). The upper arm was found a weak predictor for stature ($R = 0.045$, $SEE = 7.66$), foot length and arm span were good predictors of stature ($R = 0.73$, $SEE = 5.24$ and $R = 0.83$, $SEE = 4.25$) univariately. Multivariate, the upper arm length, foot length, and arm span show a better predictor for stature ($R = 0.85$, $SEE = 4.07$).

Conclusion: The foot length and arm span are good predictors of stature and the findings and will be useful to forensic anthropologists

Keywords: forensic anthropology, upper arm length, stature estimation, Northern Nigerians, foot length and arm span

Introduction

Anthropometric measures have advanced in building biological profiles such as stature, sex, and ancestry for identification and classification purposes [1]. Building a biological profile has been validated in various places on different races due to well-defined measurements of the population that have been established. However, there is a concern that some ethnic, races, and countries have a dearth in the literature relating to the metric dimensions of the body and this has further resulted a numerous challenges where implantation becomes rarely impossible because no established dimensions. Environmental, genetic, cultural, diet, and hormonal factors could influence an anthropometric dimension of a specific population rendering it inaccurate to use obtainable data in another population[2-4]. This study aims to generate anthropometric dimensions and evaluate how well are they in the prediction of stature among the northern Nigerians.

The upper arm length, foot length, and arm span are parameters of interest in this study, the upper arm length is a vertical measurement that runs from the lateral tip acromion down to the distal part of the arm (lateral and medial epicondyle) and has been used for some medical application such as nutritional assessment, body proportions, prosthetic fittings, muscle wasting and so on. Several authors have conducted anthropometric studies and have reported inconsistent values across different populations [5-10].

The foot length has been explored widely for various reasons; mostly for shoe design and forensic approaches. It also has some medical applications in pediatric growth assessment and prosthetic designs. Among different populations, the findings are inconsistent [11-14]. The arm span is not excluded, findings also have variation in different populations and time [15-17].

However, the variation observed further suggests that even in a particular population, diet, and lifestyle could also be attributed to the variations observed. Though, considering the numerous works that abound in different populations and the dearth of literature on northern Nigeria there is a need for continuous re-evaluation of the anthropometric dimension of a population to remain current and relevant. This has motivated the interest of this study to generate anthropometric dimensions and predict stature using the parameter for forensic and industrial applications.

Materials and Methods

Study Design

This study employed a cross-sectional descriptive approach to collect data on standing height, upper arm length, foot length, arm span, and age in northern Nigeria. The study included four hundred Nigerian northerners now residing in Kano State, Nigeria, including two hundred males and two hundred females. The study's participants range in age from 17 to 40 years. The study lasted six months (January 2024-June 2024), with the first three months spent collecting data and the second three months managing and designing the study's article. Kano City in Kano State, Nigeria was used as a sample frame to draw subjects from different northern states of Nigeria.

Sampling size and techniques

Four hundred Nigerian northerners were recruited via multi-stage random sampling techniques to ensure all subjects had an equal chance of being selected and the minimum sampling size was determined using the Taro Yamane formula for quantitative research.

Selection Criteria

In the course of the study, the researcher only recruited subjects whose grandparents and parents were of Nigeria's northern origin, between the age interval of 17 years to 40 years, and had no morphological defect that could alter the stature or the areas of measurements. Lastly, only subjects who consented to participate by filling out the consent form were allowed. The study further excluded subjects whose origin is from northern Nigeria and had some morphological defect that could alter stature and the measured part of interest. The subjects who failed to consent were also excluded from the study.

Method of Data collection

Data were collected via a semi-structured descriptive questionnaire following a personal interview and direct measurements of the body dimensions. The questionnaire was designed in two sections; A containing the bio data of the subjects, and Section B, the under-listed parameters of interest and the scoring system. The personal interview was adopted to ensure the subjects met the inclusion criteria and also to validate that the information given was correct. Using a ZT-160 stadiometer the height of the subjects was measured, and a non-stretchable and a mega-size caliper were used to measure the upper arm length and foot length. A meter rule was also used to measure the arm span. All measurement follows the international standard of anthropometric measurements.

Anthropological landmarks

- **Standing height:** it is a vertical measurement of the human body that runs from the vertex of the head to the sole when the subject is standing upright in an anatomical position.
- **Upper arm length:** this measurement runs from the lateral tip acromion down to the distal part of the arm (lateral and medial epicondyle)
- **Foot length.** This is the measurement of the foot from the big toe to the distal talus of the leg
- **Arm span;** this is a horizontal measurement of the human body from the tip of the third finger/digit to the opposite tip of the third finger/digit

Method of Data Analysis

Data from the study were analyzed using IBM SPSS version 25 and presented as mean±standard deviation. The T-test and ANOVA were employed as inferential statistics to assess sexual and age-related differences, and univariate and multivariate analyses were performed to estimate stature using individual and combination factors. Statistical significance was defined as $p < 0.05$, with 95% confidence range.



Figure 1. Measurement of the foot length (*source: field works*)



Figure 2. Measurement of standing height (*source: fieldwork*)

Results

The study comprised four hundred northern subjects between the ages 17- 40 years and Table 1, shows the descriptive statistics of the parameters for Northern Nigeria. The results present that

the minimum and maximum standing heights were 148 cm and 191.2cm respectively with a mean value of 167.65cm, the upper arm length has a mean value of 33.93cm whereas the minimum and maximum values are 23cm and 45.3cm. The foot length was measured and our result presented that the minimum, maximum, and mean values were 18.6 cm, 34.1cm, and 24.40cm respectively. The arm span had a mean value of 175.19 cm with minimum and maximum values of 150cm and 201 cm respectively.

Table 2. explores the sexual differences observed in the studied parameters among Northern Nigeria and the results present that male has a mean value for standing height, upper arm length, foot length, and arm span as 171.97 ± 6.68 cm, 32.62 ± 2.55 cm, 25.76 ± 1.17 cm, 181.72 ± 7.83 cm respectively. While the females show 163.33 ± 5.96 cm, 35.23 ± 2.19 cm, 23.04 ± 1.51 cm, and 168.67 ± 6.68 cm for standing height, upper arm length, foot length, and arm span. The comparison between the males and the females further shows that there are statistical differences observed in standing height, upper arm length, foot length, and arm span ($p < 0.05$).

Table 3, explores the age-related differences in standing height, upper arm length, foot length, and arm span and our findings show that the mean value increases with the increase in age. However, the findings have displayed statistical age-related differences in Standing height, upper arm length, foot length, and arm span ($p < 0.05$).

Table 4. Present the univariate regression for stature estimation (S) among northern Nigeria and our findings explore that upper arm length is a weak predictor for stature ($R=0.045$, $SEE=7.66$), the foot length showed a strong predictor for stature where the R was observed as 0.73 and SEE was 5.24. and the arm span also shows a strong correlation and a good predictor for stature ($R=0.83$, $SEE=4.25$). The collinearity has shown that upper arm length, foot length, and arm span are not closely related to stature and could be used as a predictive value ($VIF < 2.5$). From

the univariate regression, our study generated models for stature estimation using the upper arm length, stature could be predicted as $S=(163.34) +Al(0.13)$, using foot length, $S=(96.49)+FL(2.92)$ and using Arm span= $(53.40)+AS(0.65)$.

Table 5 showed the multivariate linear regression for stature estimation (S) among northern Nigeria across the sexes and the finding from our study shows that upper arm length, foot length, and arm span are good predictors of stature ($R=0.85$, $VIF<2.5$ and $SEE<1$). For the female population, the result showed that upper arm length, foot length, and arm span were good predictors for stature ($R=0.77$) with SEE of the studied parameters observed ($SEE<1$) which further indicates the accuracy of the prediction and the VIF of the upper arm length and foot length was $VIF<2$ indicating that the parameter is not closely related to the stature and could serve as a good predictors for stature. Among the male population, the upper arm length, foot length, and arm span display a good predictor for stature estimation ($R=0.78$) with each parameter having SEE less than 1. The collinearity has also shown that upper arm length, foot length, and arm span are accurate and better predictors for stature ($VIF<2$).

Table 1. Descriptive Statistics of the Parameters for Northern Nigeria

	min	max	Mean	SEM	SD
Age(year)	17	40	22.458	0.2002	4.0042
S (cm)	148	191.2	167.657	0.3832	7.664
AL (cm)	23	45.3	33.928	0.1358	2.7169
Fl (cm)	18.6	34.1	24.402	0.0959	1.9175
AS (cm)	150	201	175.198	0.4888	9.776

Min= minimum, max=maximum, S=standing height, AL=upper Arm Length, FL=foot length, AS=Arm span, SEM=Standard Error Mean, SD=standard deviation.

Table 2. Sexual differences in the parameters among Northern Nigeria

SEX	Male	Female	T score	p-value	Inference
S	171.97±6.68	163.33±5.96	13.65	0.0001	Sig
AL	32.62±2.55	35.23±2.19	-11.01	0.001	Sig
FL	25.76±1.17	23.04±1.51	20.19	0.001	Sig
AS	181.72±7.83	168.67±6.68	17.92	0.0001	Sig

S=standing height, AL=upper Arm Length, FL=foot length, AS=Arm span, Sig= significant ($p < 0.05$)

Table 3. Age-related differences in the parameters among Northern Nigeria

	<= 17.0	18.0 - 24.7	24.8 - 32.3	32.4+	F	p-value	Inference
S	160.75±5.30	166.95±7.69	170.06±7.06	170.96±6.98	4.88	0.002	Sig
AL	33.00±1.41	34.26±2.73	32.80±2.53	32.92±1.31	7.04	0.0001	Sig
FL	22.80±1.69	24.15±1.92	25.20±1.63	25.89±1.67	9.90	0.0001	Sig
AS	166.70±6.08	173.93±9.47	179.55±9.62	180.33±9.22	9.07	0.001	Sig

S=standing height, AL=upper Arm Length, FL=foot length, AS=Arm span, Sig= significant ($p < 0.05$)

Table 4 univariate linear regression for Stature estimation (S) among Northern Nigeria

parameters	B	R	R ²	SEE	Sig. Change	F	Collinearity Tolerance	VIF
(constant)	163.34							

AL	0.13	0.045	0.002	7.6658	0.368	0.92	1.08
(constant)	96.49						
FL	2.92	0.73	0.53	5.24	0.0001	0.34	2.50
(constant)	53.40						
AS	0.65	0.83	0.69	4.25	0.0001	0.41	2.46

S=standing height, AL=upper Arm Length, FL=foot length, AS=Arm span

Table 5 Multivariate linear regression for stature estimation (S) among Northern Nigeria

Model	Parameters	B	SEE	R	R ²	Sig	Tolerance	VIF
All	(Constant)	49.595	4.402			.000		
sexes	AL	.169	.078	0.85	0.72	.032	.922	1.085
	FL	1.048	.169			.000	.398	2.514
	AS	.495	.033			.000	.407	2.460
Female	(Constant)	53.651	6.972			.000		
	AL	.318	.173	0.77	0.59	.068	.518	1.929
	FL	1.249	.213			.000	.724	1.381
	AS	.413	.062			.000	.430	2.325
Males	(Constant)	43.721	7.771			.000		
	AL	.012	.133	0.78	0.60	.928	.792	1.263
	FL	.945	.314			.003	.678	1.475
	AS	.570	.051			.000	.581	1.720

a. Dependent Variable: STANDING HEIGHT

S=standing height, AL=upper Arm Length, FL=foot length, AS=Arm span

Discussion

The present study took account of the sex-related differences in the study parameters of the Northern Nigeria currently residing in Kano city in Kano State, Nigeria. Our findings present

that male has a mean value for standing height, upper arm length, foot length, and arm span of 171.97 ± 6.68 cm, 32.62 ± 2.55 cm, 25.76 ± 1.17 cm, and 181.72 ± 7.83 cm respectively. While the females show 163.33 ± 5.96 cm, 35.23 ± 2.19 cm, 23.04 ± 1.51 cm, and 168.67 ± 6.68 cm for standing height, upper arm length, foot length, and arm span. The implication of the finding shows that males have higher mean values in standing height, upper arm length, foot length, and arm span compared to females. The comparison between the males and the females further shows that there are statistical differences observed in standing height, upper arm length, foot length, and arm span ($p < 0.05$).

The reason for the variation observed could be attributed to genetic or hormonal factors. However, males and females have distinct makeup that guides the growth patterns and could lead to noticeable physical variations. Hormones on the other hand influence growth in males during puberty by the surge increase of testosterone that influence the bone and muscle growth leading to variation in physical dimensions among sexes [4, 5]. From an evolutionary standpoint, these morphological distinctions have developed to support diverse functions and sexual dimorphism, or the intrinsic disparities in size and appearance between men and females, has developed under natural and sexual selection forces.[5].The findings of this study agree with the findings from Asiwe et al., [5] that upper arm length is significantly different in sexes among the Mgbidi population of Imo State, Nigeria. Among northern India, Singh et al., [18] also reported that stature was statistically significant with sex, in Iran and Pakistan, Gh et al., [19] findings were consistent with our study that upper limb extremity measurements were significant with sex. Krauss et al., [20] on the anthropometry of the foot reported that sexual differences exist in foot measurements, and among the Ikwerre and Okirrika ethnic groups of Rivers State, Asiwe et al., [3] have reported that standing height has a significant difference between males and females.

Table 3, explores the age-related differences in standing height, upper arm length, foot length, and arm span. Our findings display statistical age-related differences in Standing height, upper arm length, foot length, and arm span ($p < 0.05$). However, Between the ages of 17 and 40, changes in standing height, upper arm length, foot length, and arm span are usually minimal compared to the more dramatic growth seen in childhood and adolescence. By 17 most bones have ossified, our bodies can alter in subtle ways throughout our lives. Posture, spinal compression, and muscular mass are all potential contributing factors. Poor posture, muscular imbalances, and the effects of gravity can all contribute to small height loss over time, especially as we approach our late 30s and early 40s. Our muscle mass normally peaks in our late teens and early twenties, but it might progressively drop if we don't exercise regularly. This drop can have an impact on dimensions including upper arm size, foot length, and arm span. The findings of the study agree with Li et al., [21], Tomassoni et al., [22], and Echeita et al., [23] that there are age-related differences observed in the foot. Rezende et al., [24], Perissinotto et al., [25] and López-Ortega and Arroyo, [26] findings were consistent with our study.

The studied parameter has shown a high level of sexual and age-related difference providing a better understanding of the parameters. However, the study explores to evaluate if upper arm length, foot length, and arm span could serve as a good predictor for stature. Table 4, shows the univariate regression of upper arm length, foot length, and arm span and the findings present that upper arm length is a weak predictor for stature ($R=0.045$, $SEE=7.66$), the foot length was a strong predictor for stature where the R was observed as 0.73 and SEE was 5.24. and the arm span also shows a strong correlation and a good predictor for stature ($R=0.83$, $SEE=4.25$). The collinearity has shown that upper arm length, foot length, and arm span are not closely related to stature and could be used as a predictive value ($VIF < 2.5$). Our findings are consistent with

Gardasevic et al., [27], Arifi et al., [28], Supare et al., [29], Gupta et al., [30], Dorjee and Sen, [31], and Esomonu et al., [32] that arm span is indeed a good predictor for stature. Kanchan et al., [33], Sanli et al., [13], Ozden et al., [34], and Sen and Ghosh, [35] further reported in their study on the estimation of stature using foot length that foot length is a predictor for stature estimation. However, the findings from our study disagree with Asiwe et al., [5] whose study reported upper arm length was a good predictor for stature among the Mgbidi population of Imo State, Nigeria. The findings could be contradicting but according to Asiwe et al., [5], they emphasize that muscle mass and body composition could affect and influence the estimation of stature.

Table 5, evaluates the multivariate regression for stature estimation, and the findings present that good predictors of stature include upper arm length, foot length, and arm span ($R=0.85$, $VIF<2.5$, and $SEE<1$) when combined. Upper arm length, foot length, and arm span were good predictors of stature ($R=0.77$) for the female population. The SEE of the studied parameters was observed ($SEE<1$), indicating prediction accuracy. The VIF of the upper arm length and foot length was $VIF<2$, suggesting that the parameters are not closely related to stature and could be useful predictors of stature. With each measure having a SEE smaller than 1, the male population's upper arm length, foot length, and arm span all show promise as predictors for stature estimate ($R=0.78$). The findings further suggest that for both males and females, the upper arm length, foot length, and arm span are good predictors for stature and this agrees with Asiwe et al., [5], Gardasevic et al., [27], Arifi et al., [28], Supare et al., [29], Gupta et al., [30], Dorjee and Sen, [31], Esomonu et al., [32], Kanchan et al., [33], Sanli et al., [13], Ozden et al., [34], and Sen and Ghosh, [35] among the different population.

Conclusion

The study has shown sexual and age-related differences in standing height, upper arm length, foot length, and arm span among the northern Nigerians residing in Kano. The univariate regression revealed that foot length and arm span are better predictors of stature. The multivariate regression revealed that both for males and females, the upper arm length, foot length, and arm span could serve as better predictors for stature estimation.

Consent

A written consent was distributed to all the subjects explaining the nature of the research and only those who consented were allowed to participate in the study. The consents were retrieved and preserved by the authors.

Ethical consideration

The study was approved by the research and ethics committee of the University of Port Harcourt, Port Harcourt Nigeria.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of manuscripts.

References

1. Anjani RC, Artaria MD, Singuwan P, Arunorat J, Mahakkanukrauh P. Biological Identification of Skulls in Indonesian and Thai Populations: Ancestry Estimation, Sex

Determination, Stature Estimation, and Age Estimation. *International Journal of Morphology*. 2024 Jan 1;42(1).

2. Di Fusco SA, Mocini E, Gulizia MM, Gabrielli D, Grimaldi M, Oliva F, Colivicchi F. ANMCO (Italian Association of Hospital Cardiologists) scientific statement: obesity in adults—an approach for cardiologists. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*. 2024 Jan 2;29(1):1.
3. Asiwe N, Irozulike FC, Wariboko LI, Adheke OM. Health Assessment of the Ikwerres and Okrikas Ethnic Group of Rivers State, Nigeria: Using Body Mass Index and Waist-to-Hip Ratio. *Journal of Complementary and Alternative Medical Research*. 2023 Jul 31;23(2):27-35.
4. Fewehinmi HB, Okoh PD, Oghenemavwe LE, Amadi MA, Ebieto CE, Bobbo KA, Asiwe N. Facial Analysis of the Igbo Ethnic Group of Nigeria for the Evaluation of Sexual Dimorphism. *Asian Journal of Medicine and Health*. 2023 Dec 20;21(12):67-75.
5. Asiwe N, Adheke OM, Ezeah I, Okon M, Filima PL, Buseni OV. Discriminant and Multiple Linear Regression Analysis for Sex and Stature Estimation Using Upper Arm and Forearm-Hand Length: A Study among Mgbidi Population of Imo State Nigeria. *Asian Journal of Medical Principles and Clinical Practice*. 2024 Jun 3;7(1):295-305.
6. Navid S, Mokhtari T, Alizamir T, Arabkheradmand A, Hassanzadeh G. Determination of stature from upper arm length in medical students. *Anatomical Sciences Journal*. 2014 Aug 10;11(3):135-40.
7. Hennessey WJ, Falco FJ, Goldberg G, Braddom RL. GENDER AND ARM LENGTH: INFLUENCE ON NERVE CONDUCTION PARAMETERS IN THE UPPER LIMB. *American Journal of Physical Medicine & Rehabilitation*. 1993 Aug 1;72(4):235.

8. Belt-Niedbala BJ, Ekvall S, Cook CM, Oppenheimer S, Wessel J. Linear growth measurement: A comparison of single arm-lengths and arm-span. *Developmental Medicine & Child Neurology*. 1986 Jun;28(3):319-24.
9. Datta Banik S, Azcorra H, Valentín G, Bogin B, Dickinson F. Estimation of stature from upper arm length in children aged 4.0 to 6.92 y in Merida, Yucatan. *The Indian Journal of Pediatrics*. 2012 May;79:640-6.
10. Marshall WA, Ahmed L. Variation in upper arm length and forearm length in normal British girls: photogrammetric standards. *Annals of Human Biology*. 1976 Jan 1;3(1):61-70.
11. Agnihotri AK, Purwar B, Googoolye K, Agnihotri S, Jeebun N. Estimation of stature by foot length. *Journal of forensic and legal medicine*. 2007 Jul 1;14(5):279-83.
12. Masanovic B, Gardasevic J, Arifi F. Relationship between foot length measurements and body height. *Anthropologie (1962-)*. 2019 Jan 1;57(2):227-34.
13. Sanli SG, Kizilkanat ED, Boyan N, Ozsahin ET, Bozkir MG, Soames R, Erol H, Oguz O. Stature estimation based on hand length and foot length. *Clinical Anatomy: The Official Journal of the American Association of Clinical Anatomists and the British Association of Clinical Anatomists*. 2005 Nov;18(8):589-96.
14. Hill M, Naemi R, Branthwaite H, Chockalingam N. The relationship between arch height and foot length: Implications for size grading. *Applied ergonomics*. 2017 Mar 1;59:243-50.
15. Quanjer PH, Capderou A, Mazicioglu MM, Aggarwal AN, Banik SD, Popovic S, Tayie FA, Golshan M, Ip MS, Zelter M. All-age relationship between arm span and height in different ethnic groups. *European Respiratory Journal*. 2014 Oct 1;44(4):905-12.

16. Reeves SL, Varakamin C, Henry CJ. The relationship between arm-span measurement and height with special reference to gender and ethnicity. *European journal of clinical nutrition*. 1996 Jun 1;50(6):398-400.
17. Mohanty SP, Babu SS, Nair NS. The use of arm span as a predictor of height: A study of South Indian women. *Journal of Orthopaedic Surgery*. 2001 Jun;9(1):19-23.
18. Singh B, Kaur M, Kaur J, Singh M, Batra A. Estimation of stature from forearm length in north Indians—an anthropometric study. *International Journal of Basic and Applied Medical Sciences*. 2013;3(1):201-4.
19. Gh M, Shiasi M, Hassanzadeh G, Alaghebandha N, Dehbashipour A, Zeidi H, Barbarestani M. Anthropometric characteristics of upper limb in Iranian and Pakistani subjects. *Journal of Gorgan University of Medical Sciences*. 2014 Oct 10;16(3):80-5.
20. Krauss I, Grau S, Mauch M, Maiwald C, Horstmann T. Sex-related differences in foot shape. *Ergonomics*. 2008 Nov 1;51(11):1693-709.
21. Li PL, Yick KL, Zhang LY, Keung YC. Evaluation of Age-related Differences in Foot Anthropometry among Women. *AATCC Journal of Research*. 2023 Jan 26:24723444221147977.
22. Tomassoni D, Traini E, Amenta F. Gender and age-related differences in foot morphology. *Maturitas*. 2014 Dec 1;79(4):421-7.
23. Echeita JA, Hijmans JM, Smits S, Van der Woude LH, Postema K. Age-related differences in women's foot shape. *Maturitas*. 2016 Dec 1;94:64-9.
24. Rezende FA, Ribeiro AQ, Priore SE, Franceschini SD. Anthropometric differences related to genders and age in the elderly. *Nutrición Hospitalaria*. 2015;32(2):757-64.

25. Perissinotto E, Pisent C, Sergi G, Grigoletto F, Enzi G, ILSA Working Group. Anthropometric measurements in the elderly: age and gender differences. *British Journal of Nutrition*. 2002 Feb;87(2):177-86.
26. López-Ortega M, Arroyo P. Anthropometric characteristics and body composition in Mexican older adults: age and sex differences. *British Journal of Nutrition*. 2016 Feb;115(3):490-9.
27. Gardasevic J, Rasidagic F, Krivokapic D, Corluca M, Bjelica D. Stature and Its Estimation Utilizing Arm Span Measurements in Male Adolescents from Federation of Bosnia and Herzegovina Entity in Bosnia and Herzegovina. *Montenegrin Journal of Sports Science & Medicine*. 2017 Mar 1;6(1).
28. Arifi F, Sermahaj S, Gardasevic J, Alaj I, Metaj Z. Stature and its estimation utilizing arm span measurements of both gender adolescents from southern region in Kosovo. *Sport Mont*. 2018 Jun 1;16(2):51-4.
29. Supare MS, Bagul AS, Pandit SV, Jadhav JS. Estimation of stature from arm span in medical students of Maharashtra, India. *Annals of Medical and Health Sciences Research*. 2015 May 26;5(3):218-21.
30. Gupta C, Palimar V, Guru P. Estimation of stature from the arm span of an individual in South Indian population. *Annals of Bioanthropology*. 2016 Jan 1;4(1):17.
31. Dorjee B, Sen J. Estimation of stature from arm span, arm length and tibial length among Bengalee. *Human Bio Review*. 2016;5(2):128-45.
32. Esomonu UG, Ijomone OM, Mba C, Oranusi A. Estimation of stature using arm span length amongst Bekwara ethnic group of Cross River State, Nigeria. *Annals of Bioanthropology*. 2015 Jul 1;3(2):55.

33. Kanchan T, Menezes RG, Moudgil R, Kaur R, Kotian MS, Garg RK. Stature estimation from foot dimensions. *Forensic Science International*. 2008 Aug 6;179(2-3):241-e1.
34. Ozden H, Balci Y, Demirüstü C, Turgut A, Ertugrul M. Stature and sex estimate using foot and shoe dimensions. *Forensic Science International*. 2005 Jan 29;147(2-3):181-4.
35. Sen J, Ghosh S. Estimation of stature from foot length and foot breadth among the Rajbanshi: an indigenous population of North Bengal. *Forensic Science International*. 2008 Oct 25;181(1-3):55-e1.

UNDER PEER REVIEW