

Effect of different concentrations of IBA on rooting and growth factors in apical shoot cuttings of *Callistemon lanceolatus* under polytunnel condition

Commented [EA1]: Suggested topic "Impact of Various IBA concentrations on Rooting and Growth in Apical Shoot Cuttings of *Callistemon lanceolatus* (Sm.) Sweet in Poly tunnel Conditions"

Abstract. The study was undertaken to standardize the optimal concentration of rooting hormone suitable for propagation of *Callistemon lanceolatus* and it was conducted at the Vanavarayar Institute of Agriculture, Pollachi, Coimbatore, India from December 2022 to September 2023. Thirteen different treatments were tried viz., T₁- Apical shoot cuttings with IAB at 1000 ppm; T₂- Apical shoot cuttings with IAB at 2000 ppm; T₃- Apical shoot cuttings with IAB at 3000 ppm; T₄- Apical shoot cuttings with IAB at 4000 ppm; T₅- Apical shoot cuttings with IAB at 5000 ppm; T₆- Apical shoot cuttings with IAB at 6000 ppm; T₇- stem cuttings with IAB at 1000 ppm; T₈- stem cuttings with IAB at 2000 ppm; T₉- stem cuttings with IAB at 3000 ppm; T₁₀- stem cuttings with IAB at 4000 ppm; T₁₁- stem cuttings with IAB at 5000 ppm; T₁₂- stem cuttings with IAB at 6000 ppm; T₁₃- control (treated with distilled water). The studies were conducted in a completely randomized block design (CRD). The results of the study depicted that the maximum rooting percentage and root length were observed in T₆ (73.28±8.87%) and (12.54±0.00cm) due to the exogenous application of auxin.

Commented [EA2]: IBA or IAB?

Commented [EA3]: Rewrite the abstract to be summary of findings and not statement of results

Commented [EA4]: Include conclusive statements that depicts importance of findings

Keywords: *Callistemon lanceolatus*, rooting hormone, IBA, apical shoot cuttings, stem cuttings,

Introduction

Callistemon lanceolatus, which belongs to the Myrtaceae family, native to Australia, is widely distributed in subtropical and tropical regions (Sowndhararajan *et al.*, 2021). This tree, characterized by its striking bright red flowers resembling bottlebrushes, is commonly cultivated in gardens throughout India as an ornamental plant (Kumar *et al.*, 2020). Referred to as the "lemon bottlebrush" due to its distinctive cylindrical brush-like red flowers (Singh *et al.*, 2020), this plant is known to possess various biological activities in its aerial parts, including antimicrobial (Nazreen *et al.*, 2020), antioxidant, antidiabetic (Ahmad *et al.*, 2018; Kumar *et al.*, 2011), anti-inflammatory (Kumar *et al.*, 2011), and anti-proliferative (Park *et al.*, 2018) properties. Notably, essential oils extracted from the leaves of *C. lanceolatus* exhibit antimicrobial and anti-inflammatory properties (Shukla *et al.*, 2012; Sudhakar *et al.*, 2004).

The propagation of ornamental plants has traditionally relied on seed germination and vegetative propagation techniques (Hartman *et al.*, 2002; Shokri, 2012). However, these conventional methods come with constraints related to time, labour, and genetic fidelity (Singh *et al.* 2013). Seed germination can be unpredictable and time-consuming, resulting in a wide range of progeny diversity due to genetic recombination (Müller-Starck *et al.*, 2005; Acquah, 2015). To address these challenges and meet the increasing demand for genetically identical, high-quality plant material, clonal propagation techniques have gained prominence (Muchugi *et al.*, 2023). Clonal propagation enables the mass production of plants with identical genetic characteristics, ensuring uniformity in desirable attributes such as flower color, scent, and growth habits (Gradziel, 2012; Cardoso *et al.* 2023). Some species necessitate vegetative propagation for short-term regeneration (Leakey, 2004; Mei *et al.*, 2023). Over the past decade, vegetative propagation approaches have gained traction for the bulk propagation of enhanced genetic material, particularly for tree species with problematic seeds (Kantarli, 1993). Notably, some tropical trees have demonstrated the potential of

Commented [EA5]: used too often change using another synonym

vegetative propagation, which improves yield potential by reducing variation among the ramets (Parthiban *et al.*, 1999). This progress in vegetative propagation has evolved into clonal technology, emerging as an alternative to traditional seed-based propagation (Parthiban and Seenivasan, 2017). When it comes to vegetative multiplication through cuttings, the use of crown branches of advanced ontogenetic age has been observed to reduce the percentage of propagule roots (Ciriello and Mori, 2015). However, the utilization of spontaneous or induced basal sprouts has yielded excellent rooting indexes, resulting in enhanced efficiency in vegetative proliferation (Rickli *et al.*, 2015). Compared to seed propagation, clonal propagation is a simpler, faster, and more efficient way to rapidly expand millions of cuttings from elite germplasms (Gonin *et al.*, 2019; Solgi *et al.*, 2022). ("Exogenous hormones supplementation improve adventitious root formation in woody plants")

Callistemon lanceolatus trees pose a unique challenge as they require a substantial amount of time for seed release, ranging from one year to several years, with some even requiring fire to activate the capsules. Pre-treatment of the seeds is often necessary to disrupt this dormancy (Brophy *et al.*, 2013; Lamont *et al.*, 2020). Furthermore, these trees exhibit variable seed production in different regions, creating operational challenges in meeting annual planting material demands. The proposed clonal technology offers a solution to these challenges while ensuring uniformity in growth and production. It enables the large-scale multiplication of genetically identical ramets, promoting the propagation of elite genotypes and the bulk production of high-quality planting material for commercial ornamental trees. In this context, clonal technology is a game-changer in boosting the generation of improved genotypes in *Callistemon lanceolatus* trees, facilitating large-scale commercial production.

Vegetative propagation of the bottlebrush plant poses challenges due to the inherent difficulty in rooting stem cuttings, which necessitates specific hormonal and environmental interventions to induce root formation (Zarinbal *et al.*, 2005). Root induction in stem cuttings of challenging-to-root species has been addressed in numerous studies, employing methods such as auxin treatment, bottom heat, wounding and mist systems (Dawson and King, 1994; Hartman *et al.*, 2002; Dominik and Gregor, 2007; Shokri, 2012). Top of Form The imperative for mass propagation of *C. lanceolatus* through cloning has emerged as a significant priority in the realms of horticulture and conservation. Clonal propagation techniques, which involve replicating individuals with identical genetic characteristics as the parent tree, hold the promise of preserving and proliferating the remarkable attributes of this species. This approach ensures that the enchanting qualities, both in terms of appearance and fragrance, remain consistent across the newly propagated species.

This paper aims to explore the methods of clonal propagation for *C. lanceolatus*, examining the different concentrations of IBA on the growth. By understanding and implementing these techniques, horticulturists and enthusiasts can ensure the continued cultivation and conservation of this iconic and culturally significant tree, which not only adds beauty to our surroundings but also holds a special place in our traditions and heritage.

Materials and Methods

Study site: The study was conducted at the Vanavarayar Institute of Agriculture, Pollachi, Coimbatore, India from December 2022 to September 2023. The location experiences a warm, humid climate with an average annual rainfall of about 2100 to 2500 mm. The average temperature ranges from 23.8°C to 34.6°C with a relative humidity of 51 to 82%.

Experimental material: The study was undertaken to standardize the optimal size of cuttings suitable for propagation. The cuttings collected from identified tree, were grouped into apical shoot cuttings and stem cuttings without an apical shoot). These cuttings were collected from the tree by conventional cutting methods Schwambach *et al.* (2008); Wendling *et al.* (2010). The cuttings were harvested using sterile pruning scissors in the early morning. After harvesting, they were screened for desired length (8- 10cm) by using scale) and

Commented [EA6]: 2017? Check

Commented [EA7]: Plagiarized material rephrase

Commented [EA8]: Rephrase I do not think you mean that the tree itself is a problem

Commented [EA9]: Not listed in references

Commented [EA10]: Not Listed in references

Commented [EA11]: Not listed, include in reference list

Commented [EA12]: rephrase for clarity

Commented [EA13]: Were screened or cut to size?

diameter (2-4 mm) by using a calibrated vernier calliper following which they were placed in ice box to prevent damage during transportation to laboratory. The cuttings were treated with an aqueous solution of 0.1% Bavistin for 15 minutes and subsequently washed with distilled water (Sabari, 2017). Since the discovery of natural plant rooting hormones for propagation, it has been intuitive to apply these substances to the basal end of cuttings to initiate new root formation. Six concentrations of aqueous rooting solutions viz. 1000, 2000, 3000, 4000, 5000 and 6000 mg L⁻¹ IBA were prepared in 1N NaOH with distilled water. Untreated cutting was considered as control (cuttings were dipped in distilled water which served as a control). 15 cuttings were prepared following the above procedure in each treatment with 3 replications.

After auxin treatment, the cuttings were planted in the polybags containing coir pith as a rooting medium. One third of the basal cut portion was inserted in different rooting mediums. The medium has to be kept damp and the surface even. Holes should be punched into the medium to allow the insertion of the cuttings without damaging the cambium or removing the rooting hormone. After the cuttings were stuck, the rooting medium was pressed slightly around them. The cuttings were kept in polytunnel for maintaining the relative humidity in the range of 60 to 80% and the optimal temperature (25-30°C / 15-20°C day/night). The cuttings were regularly watered and drenched with 0.1% Bavistin at a 15-day interval to avoid desiccation damage and the attack of fungal pathogens. The rooting experiment was carried out for 60 days. Rooted cuttings were transferred to polythene bags (15 x 25 cm) containing soil + poultry manure (5:1) and kept in a shade house for 30 days.

Experimental Design and Data Collection

Thirteen different treatments were tried viz., T₁- Apical shoot cuttings with IAB at 1000 ppm; T₂- Apical shoot cuttings with IAB at 2000 ppm; T₃- Apical shoot cuttings with IAB at 3000 ppm; T₄- Apical shoot cuttings with IAB at 4000 ppm; T₅- Apical shoot cuttings with IAB at 5000 ppm; T₆- Apical shoot cuttings with IAB at 6000 ppm; T₇- stem cuttings with IAB at 1000 ppm; T₈- stem cuttings with IAB at 2000 ppm; T₉- stem cuttings with IAB at 3000 ppm; T₁₀- stem cuttings with IAB at 4000 ppm; T₁₁- stem cuttings with IAB at 5000 ppm; T₁₂- stem cuttings with IAB at 6000 ppm; T₁₃- control (treated with distilled water). The studies were conducted in a completely randomized block design (CRD) with three replications and 15 cuttings per treatment per replication.

The biometric parameters of epicormic shoots such as sprouting per cent, rooting per cent, root length, number of roots per cutting, shoot length (cm), number of leaves, leaf length (cm), leaf width (cm), survival per cent and vigour index were measured and recorded.

Results

The sprouting percentage can vary depending on the plant species, the quality of the plant parts, and the environmental conditions (such as temperature, humidity, and soil quality). It is an important metric in agriculture, horticulture, and plant propagation, as it indicates the success of sprouting process. The results of the study revealed that the maximum sprouting percent of *C. lanceolatus* was observed in T₆ (80.87±0.86%), T₁₂ (79.15±3.43%), T₅ (75.73±4.53%), T₁₁ (73.13±2.89%) whereas the minimum sprouting per cent was noticed in T₁₃ (32.73±1.80%) (Table 1). Shoot length refers to the measurement of the length of the above-ground part of a plant, specifically the stem or shoots. It is a common parameter used in plant biology and agriculture to assess the growth and development of plants. Measuring shoot length can provide valuable information about a plant's response to various environmental factors, such as light, water, and nutrients, and can be an important indicator of overall plant health and productivity. The study reported that the highest shoot length of *C. lanceolatus* was recorded in T₆ (12.72±0.73cm) followed by T₁₂ (11.68±0.05cm), T₅ (11.65±0.17cm) whereas the lowest was reported in T₁₃ as (8.23±0.36cm) (Table 2).

The number of leaves per cutting can vary widely depending on the plant species, the size of the cutting, and the specific growth conditions. Some cuttings may have only a few

Commented [EA14]: Not listed

Commented [EA15]: delete

Commented [EA16]: Change to words

Commented [EA17]: ? Nursery bags?

Commented [EA18]: State the different rooting media applied

Commented [EA19]: Has to be or was kept? Be specific about what was done

Commented [EA20]: Rephrase stating exactly what was done

Commented [EA21]: I think you mean Indole-3-Butyric acid (IBA)? Please correct to reflect this

Commented [EA22]: ?

Commented [EA23]: ?

Commented [EA24]: ?

Commented [EA25]: Discussion?

Commented [EA26]: Delete not necessary

Commented [EA27]: These statements are discussion and too elementary to be stated here

Commented [EA28]: Rephrase

Commented [EA29]: Confusing how many plant species was used for this study?

leaves, while others can have several leaves. It's essential to consider the type of plant and its specific requirements when propagating through cuttings, as the number of leaves can influence the success of propagation. The highest number of leaves per cutting was observed in T₆ as 9.45±0.26 followed by T₁₂ (8.56±0.03) whereas the lowest was reported in T₁₃ (2.35±0.06). The result of present investigation clearly exhibits that the leaf length of *C. Lanceolatus* varied significantly among the treatments, the longest leaf was observed in T₁₂ (5.87±0.32cm), T₁₁ (5.74±0.24cm), T₆ (5.67±0.27cm) while the smallest leaf was noticed in T₁₃ (1.11±0.02cm).

Rooting efficiency

Rooting percentage often referred to as "rooting success" or "rooting efficiency," is a crucial metric in horticulture and plant propagation. It measures the success rate of stem cuttings developing roots, a key step in clonal propagation. Rooting percentage is the proportion of stem cuttings that successfully form roots compared to the total number of cuttings used. The results of the study depicted that the maximum rooting percentage was observed in T₆ (73.28±8.87%) and T₁₂ (72.27±1.97%), followed by T₁₁ (57.39±0.03%), while the minimum percentage was noticed in T₁₃ (20.19±1.27%). The highest root length was reported in T₆ (12.54±0.00cm); T₁₂ (12.01±0.29cm) and the lowest was recorded in T₁₃ (1.76±0.02cm). The result of the study showed that the number of roots per cutting varied significantly between the treatments. Among the treatments the maximum number of roots was noticed in T₁₂ (6.45±0.03), T₆ (6.45±0.36) whereas the minimum was recorded in T₁₃ (1.02±0.01).with regard to survival per cent the highest was reported in T₆ (69.84±0.65%) followed by T₁₂ (63.54±0.48%) and the lowest was registered in T₁₃ (20.45±0.20%) (Fig 4&5).

Discussion

Plant propagation by cuttings has several advantages including maintenance of the specific characteristics of the plant from which the cutting was collected, giving true-to-type plant materials, being simple and easy to conduct, it does not need a large space and being a cheaper propagation method compared to layering, grafting or budding. The success of plant propagation by cuttings is shown by root and shoot formation and growth which is well documented to be affected by several factors, including genotypes, physiological and ontogenetic age of cuttings, endogenous hormone contents, type of wood, carbohydrate contents, preconditioning treatment of cuttings and external factors such as micro-environment of cuttings and the use of root promoting substances (Dirr & Heuser, 2006; Hartmann *et al.*, 2011). One of the most important and widely reported factors is the use of auxins such as IBA as root promoting substances (Abu-Zahra *et al.*, 2013; Rahdari *et al.*, 2014; Costa *et al.*, 2015). IBA at 6000ppm concentrations significantly stimulated adventitious root formation as shown by the marked increases in rooting percentage. It was also shown that regardless of the types of auxins applied, increasing concentrations from 1000 to 6000 ppm caused significant increases in the number of roots per cutting, all with a 73% rooting percentage (Fig 1&2). These findings agree with the previous statement that for more than 70 years, IBA has been effectively used as a root promoting substance for cuttings from various species commercially and has been the backbone of cutting propagation success (Dirr & Heuser, 2006).

Auxins are widely recognized for their significant role in promoting the development of adventitious roots in vegetative propagules. Both exogenously applied and endogenous auxins influence the rooting potential and the quantity of roots produced in cuttings (Wen *et al.*, 2016; Gonin *et al.*, 2019). Notably, the impact of externally administered growth hormones on the rooting process exhibits significant variation between the different concentrations of IBA. The sprouting percentage of stem cuttings refers to the proportion of stem cuttings from a plant species that successfully develop new sprouts or shoots, typically

Commented [EA30]: Delete or rephrase for clarity

Commented [EA31]: Change to "and"

Commented [EA32]: Statistical inference and should be properly stated to reflect that

Commented [EA33]: This explanations are out of place here, they are of benefit in discussion

Commented [EA34]: Rephrase for clarity

as a result of vegetative propagation. This percentage is a measure of the cloning or propagation success and can vary depending on factors such as the species, the specific method of propagation, and the conditions under which the cuttings are kept. Higher sprouting percentages indicate more successful propagation. It is an important metric in horticulture, agriculture, and forestry when reproducing plants asexually (Yusnita *et al.*, 2017; Husen & Pal, 2006).

Days taken to start sprouting and days taken to attain fifty percent sprouting of cuttings were significantly affected by different growth regulators but the earliest sprouting of cuttings was noted in IBA at 6000ppm whereas, latest sprouting of cuttings was noted in control. (“EFFECT OF GROWTH REGULATORS AND BIOFERTILIZERS ON SURVIVAL OF ...”) The percentage of success of cuttings was also found significantly higher at IBA 6000ppm, whereas the minimum percentages of success of cutting were recorded in control. This may be due to the increased level of Auxins, which resulted in earlier completion of physiological processes in rooting and sprouting of cuttings. (“Effect of growth regulators and biofertilizers on survival of ...”) A plethora of workers have also reported similar values in *Citrus limonburn* as 81.90% (Kumar *et al.*, 2015); *Calliandra haematocephala* as 53.33% (Kaur and Grewal, 2017). IBA at 6000ppm produced the maximum shoot length, but the minimum shoot length was observed in control. The current study was consistent with the findings of Sevik & Guney (2013) in *Melissa officinalis* as 6.71cm when treated with IBA at 1000ppm. The study is consistent with the findings of Yusnita *et al.* (2017) in *Syzygium malaccense* as 3.30cm treated with IBA at 4000ppm; Yeshiwas *et al.* (2015) in rose stem cuttings treated with IBA at 2500ppm (15.47cm). The total number of leaves per cutting was also found to be significantly higher in IBA at 6000ppm, while the lowest number of leaves per cutting was noted in control. Although, growth regulators significantly influenced the shoot length and maximum number of leaves per cutting it was observed that the maximum was at IBA6000ppm. This may be due to the vigorous root system which increased nutrients uptake under this treatment. (“EFFECT OF GROWTH REGULATORS AND BIOFERTILIZERS ON SURVIVAL OF ...”) It is in accordance with the findings of Yusnita *et al.* (2017) in *Syzygium malaccense* as and it was lower than the earlier reports of Yeshiwas *et al.* (2015) in rose stem cuttings as 58.50; Damar *et al.* (2014) in *Punica granatum* treated with IBA at 2000ppm (37.77). Poor performance of cuttings as regards the percentage of success and number of leaves per shoot and per cuttings were found to be under control (T₁₃). These results may be attributed to the fact that growth attributes in terms of root and shoot growth parameters are affected by the exogenous application of required growth regulators. This is depicted in lowest physiological activity for triggering root initiation and development and finally all other growth parameters of cuttings were seriously affected. (“EFFECT OF GROWTH REGULATORS AND BIOFERTILIZERS ON SURVIVAL OF ...”) This may be due to low activity of the growth substance and low physiological activity.

Rooting percentage is a critical indicator of the success of clonal propagation methods. It directly influences the number of new plants that can be produced from a mother plant, which can be crucial for agricultural, horticultural, and forestry practices (Sarkar *et al.*, 2023). As regards root characters like rooting per cent, root length, number of roots per cutting, the maximum value in IBA at 6000 ppm. Whereas minimum rooting per cent, root length, number of roots per cutting were observed in control (Fig. 3). The longest root was noted in IBA 6000ppm, while the shortest root was observed in control in this experiment. This may be due to the higher accumulation of photosynthates, metabolites and nutrients under this treatment. (“EFFECT OF GROWTH REGULATORS AND BIOFERTILIZERS ON SURVIVAL OF ...”) Joel *et al.* (2023) suggested that IBA at 400ppm gives maximum rooting success in *Chrysanthemum* (94.62%). Topacoglu *et al.* (2016); Sevik & Guney

Commented [EA35]: A bit of confusion here 3.30cm and 15.47cm are these your results or the authors cited, just stating that the results are consistent with the findings of others should

Commented [EA36]: It will help if where the results are situated is stated here, say “Table 1 or Figure 1”

Commented [EA37]: Please do rephrase for clarity and just state that the IBA 6000ppm treatment recorded optimal number of leaves

Commented [EA38]: Remove the ambiguity here by stating that the IBA treatment produced optimal results because of increased root production. And this is in agreement with the work of Yusnita *et al.*; Now the comparison been made here seems not correct if the plants are not of the same species and the concentration of IBA used is not the same. However if you can justify these claims, make them clearer.

Commented [EA39]: Rephrase, conclusion is that treatment performed better than control

Commented [EA40]: Use some other words, your results are substantive enough

Commented [EA41]: One word, correct through out your article this mistake is repeated several times

Commented [EA42]: ?

Commented [EA43]: Root length?

Commented [EA44]: These can be avoided if you try to restrict your inference to previous reports and literature search on the topic

(2013); Suleiman *et al.* (2012) also observed better rooting percentages with IBA in *Ficus benjamina* (93.90%); *Melissa officinalis* (80.00%); *Capparis spinosa* (79.39%).

Conclusion

Table1. Effect of IBA concentrations on shoot growth response of *Callistemon lanceolatus* cuttings under poly tunnel condition

Treatments	SP (%)	SL (Cm)	NL	LL (Cm)
T1	46.32±1.88 ^c	8.43±0.27 ^{de}	5.23±0.37 ^{fg}	2.43±0.00 ^g
T2	48.49±1.81 ^c	8.56±0.44 ^{de}	5.87±0.27 ^{def}	3.12±0.07 ^f
T3	48.52±1.32 ^c	9.24±0.45 ^{cde}	5.95±0.06 ^{cde}	3.78±0.06 ^{de}
T4	61.87±2.68 ^b	10.47±0.42 ^{cd}	6.52±0.22 ^c	4.89±0.09 ^b
T5	75.73±4.53 ^a	11.65±0.17 ^{ab}	7.89±0.09 ^b	4.97±0.16 ^b
T6	80.87±0.86 ^a	12.72±0.73 ^a	9.45±0.26 ^a	5.67±0.27 ^a
T7	40.14±0.42 ^{cd}	7.90±0.35 ^e	4.57±0.08 ^g	3.34±0.12 ^{ef}
T8	42.34±0.42 ^c	8.16±0.25 ^e	5.34±0.11 ^{ef}	3.58±0.22 ^{ef}
T9	57.25±0.20 ^b	8.24±0.51 ^{de}	5.65±0.10 ^{def}	4.23±0.04 ^{cd}
T10	61.98±0.69 ^b	9.64±0.11 ^{cd}	6.23±0.09 ^{cd}	4.76±0.12 ^{bc}
T11	73.13±2.89 ^a	11.13±0.52 ^c	6.43±0.32 ^c	5.74±0.24 ^a
T12	79.15±3.43 ^a	11.68±0.05 ^{ab}	8.56±0.03 ^b	5.87±0.32 ^a
T13	32.73±1.80 ^d	8.23±0.36 ^{de}	2.35±0.06 ^h	1.11±0.02 ^h
MSE	2.19	0.39	0.19	0.17
P (0.001)	0.00	0.00	0.00	0.00

Commented [EA45]: This should be included and state categorically what your finds mean and their importance to Agriculture or clonal propagation of *Callistemon lanceolatus*

Commented [EA46]: Tables need footnote for T1.... Or state what they are on the tables

Table2. Effect of IBA concentrations on rooting response of *Callistemon lanceolatus* cuttings under poly tunnel condition

Treatments	RP (%)	RL (Cm)	NR	SP (%)
T1	33.58±1.87 ^{de}	3.28±0.07 ^f	3.89±0.09 ^c	32.76±0.86 ^s

T2	34.12±0.81 ^{de}	5.87±0.15 ^{de}	3.76±0.09 ^c	48.76±3.21 ^{de}
T3	37.40±0.03 ^d	6.39±0.23 ^d	4.09±0.27 ^{bc}	50.23±0.22 ^{cd}
T4	48.47±2.39 ^c	9.25±0.39 ^c	4.86±0.08 ^b	53.76±3.44 ^{cd}
T5	50.12±1.97 ^c	10.78±0.78 ^b	5.97±0.35 ^a	61.89±0.37 ^b
T6	73.28±3.87 ^a	12.54±0.00 ^a	6.45±0.36 ^a	69.84±0.65 ^a
T7	30.26±0.75 ^e	3.11±0.10 ^f	3.89±0.17 ^c	31.98±2.09 ^g
T8	36.53±0.04 ^{de}	4.98±0.24 ^e	3.76±0.01 ^c	35.67±1.40 ^{fg}
T9	39.18±2.53 ^d	5.23±0.21 ^e	4.09±0.03 ^{bc}	42.36±0.54 ^{ef}
T10	50.32±1.00 ^c	8.76±0.22 ^c	4.86±0.19 ^b	49.67±2.68 ^{cd}
T11	57.39±0.03 ^b	10.65±0.47 ^b	5.97±0.43 ^a	56.45±4.11 ^{bc}
T12	72.27±1.94 ^a	12.01±0.29 ^a	6.45±0.03 ^a	63.54±0.48 ^{ab}
T13	20.19±1.27 ^f	1.76±0.02 ^g	1.02±0.01 ^d	20.45±0.20 ^h
MSE	1.79	0.31	0.21	2.04
P (0.001)	0.00	0.00	0.00	0.00




Figure 1. The data exhibit a normal distribution in relation to the growth parameters.

Commented [EA47]: More components needed on this chart for clarity

Commented [EA48]: This is not clear to me, is there anything that can be done, maybe a different style of presentation?




Figure2. growth response of *Callistemon lanceolatus* cuttings under poly-tunnel condition

Shoot




Figure 3. Rooting response of *Callistemon lanceolatus* cuttings under poly tunnel condition




Figure 4. The rooting percentage of both apical shoot cuttings and stem cuttings is directly correlated with the survival percentage of the cuttings.

Commented [EA49]: These clustered together so not sure of what is been depicted here, rearrange for clarity or make appropriate corrections




Figure 5. The rooting performance of the cuttings positively correlates with the shoot growth performance of the cuttings.

References

1. Sowndhararajan, K., Deepa, P., & Kim, S. (2021). A review of the chemical composition and biological activities of *Callistemon lanceolatus* (Sm.) Sweet. *Journal of Applied Pharmaceutical Science*, 11(12): 065-073.
2. Kumar, R., Gupta, A., Singh, A. K., Bishayee, A., & Pandey, A. K. (2020). The antioxidant and antihyperglycemic activities of bottlebrush plant (*Callistemon lanceolatus*) stem extracts. *Medicines*, 7(3): 11.
3. Singh, P. P., Kumar, A., & Prakash, B. (2020). Elucidation of antifungal toxicity of *Callistemon lanceolatus* essential oil encapsulated in chitosan nanogel against *Aspergillus flavus* using biochemical and in-silico approaches. *Food Additives & Contaminants: Part A*, 37(9): 1520-1530.
4. Nazreen, S., Mahboob Alam, M., Hamid, H., Ali, M., & Sarwar Alam, M. (2020). Chemical constituents with antimicrobial and antioxidant activity from the aerial parts of *Callistemon lanceolatus* (Sm.) Sweet. *Natural product research*, 34(22): 3275-3279.
5. Ahmad, K., Hafeez, Z. B., Bhat, A. R., Rizvi, M. A., Thakur, S. C., Azam, A., & Athar, F. (2018). Antioxidant and apoptotic effects of *Callistemon lanceolatus* leaves and their compounds against human cancer cells. *Biomedicine & Pharmacotherapy*, 106: 1195-1209.
6. Kumar, S., Kumar, V., & Prakash, O. (2011). Antidiabetic, hypolipidemic, and antioxidant activities of *Callistemon lanceolatus* leaves extract. *Journal of Herbs, Spices & Medicinal Plants*, 17(2): 144-153.
7. Park, S., Lim, W., Jeong, W., Bazer, F. W., Lee, D., & Song, G. (2018). Sideroxylin (*Callistemon lanceolatus*) suppressed cell proliferation and increased apoptosis in ovarian cancer cells accompanied by mitochondrial dysfunction, the generation of

reactive oxygen species, and an increase of lipid peroxidation. *Journal of Cellular Physiology*, 233(11): 8597-8604.

8. Shukla, R., Singh, P., Prakash, B., & Dubey, N. K. (2012). Antifungal, aflatoxin inhibition and antioxidant activity of *Callistemon lanceolatus* (Sm.) Sweet essential oil and its major component 1, 8-cineole against fungal isolates from chickpea seeds. *Food Control*, 25(1): 27-33.
9. Sudhakar, M., Rao, C. V., Rao, A. L., Ramesh, A., Srinivas, N., Raju, D. B., & Murthy, B. K. (2004). Antinociceptive and anti-inflammatory effects of the standardized oil of Indian *Callistemon lanceolatus* leaves in experimental animals. *East and Central African Journal of Pharmaceutical Sciences*, 7(1), 10-15.
10. Hartman HT, Kester DE, Davies FT, Genev RL (2002). *Plant Propagation, Principles and Practices*. 7th Edn. Prentice Hall, Englewood Cliffs, New Jersey, USA.
11. Shokri, S., Zarei, H., & Alizadeh, M. (2012). Evaluation of rooting response of stem cuttings and in vitro micro-cuttings of bottlebrush tree (*Callistemon viminalis*) for commercial mass propagation. *Journal of Agricultural Research*, 1(10): 424-428.
12. Singh, S. R., Singh, R., Kalia, S., Dalal, S., Dhawan, A. K., & Kalia, R. K. (2013). Limitations, progress and prospects of application of biotechnological tools in improvement of bamboo—a plant with extraordinary qualities. *Physiology and Molecular Biology of Plants*, 19, 21-41.
13. Acquaah, G. (2015). Conventional plant breeding principles and techniques. *Advances in plant breeding strategies: Breeding, biotechnology and molecular tools*, 115-158.
14. Müller-Starck, G., Ziehe, M., & Schubert, R. (2005). Genetic diversity parameters associated with viability selection, reproductive efficiency, and growth in forest tree species. In *Forest diversity and function: Temperate and boreal systems* (pp. 87-108). Berlin, Heidelberg: Springer Berlin Heidelberg.

15. Muchugi, A., Kalinganire, A., Chege, J., & Nyoka, I. (2022). Commodity tree crop planting material infrastructures in Africa.
16. Cardoso, J. C., Vilcherrez-Atoche, J. A., Iiyama, C. M., Germanà, M. A., & Vendrame, W. A. (2023): Breeding of Orchids Using Conventional and Biotechnological Methods: Advances and Future Prospects. *Advances in Orchid Biology, Biotechnology and Omics*, 27-58.
17. Gradziel, T. M. (2012). Traditional genetics and breeding. *Genetics, genomics and breeding of crop plants*, CRC Press, Boca Raton, 22-54.
18. Leakey, R. R. (2004). *Physiology of vegetative reproduction*. Academic Press.
19. Mei, G., Chen, A., Wang, Y., Li, S., Wu, M., Liu, X., & Hou, X. (2023). A simple and efficient in planta transformation method based on the active regeneration capacity of plants. *bioRxiv*, 2023-01.
20. Kantarli, M. (1993). Vegetative propagation of *Hopea odorata* by cuttings: a low-cost technology. *Vegetative propagation of Hopea odorata by cuttings: a low-cost technology.*, (16).
21. Parthiban, K. T., Palanikumar, B., & Krishnakumar, N. (2017). *Clonal Evaluation, Variability and Association Studies in Casuarina junghuhniana. Plantation and Agroforestry Pulpwood Value Chain Approach*, 194.
22. Parthiban K. T. and Seenivasan. R (2017). *A complete value chain approach*. Scientific publisher. Jodhpur, p629.
23. Ciriello, E., & Mori, E. S. (2015). Rooting of Guanandi (*Calophyllum brasiliense* CAMBESS) cuttings using Indole-butyric acid. *Cerne*, 21: 641-648.
24. Rickli, H.C., Bona, C., Wendling, I., Koehler, H.S., Zuffellato-Ribas, K.C. (2015). Origem de brotações epicórmicas e aplicação de ácido indolilbutírico no

Commented [EA50]: 2023? check

Commented [EA51]: You cant use this symbol be consistent

enraizamento de estacas de *Vochysia bifalcata* Warm. *Ciência Florestal* 25: 385-393.

Commented [EA52]: ?

25. Gonin, M., Bergougnoux, V., Nguyen, T. D., Gantet, P., & Champion, A. (2019). What makes adventitious roots? *Plants*, 8(7): 240.
26. Solgi, M., & Sahraei, F. (2022). Influence of cutting lengths and IBA on propagation of red willow ornamental-medicinal plant by stem cutting. *European Journal of Horticultural Science*, 87(2): 1-7.
27. Lamont, B. B., Pausas, J. G., He, T., Witkowski, E. T., & Hanley, M. E. (2020). Fire as a selective agent for both serotiny and nonserotiny over space and time. *Critical Reviews in Plant Sciences*, 39(2), 140-172.
28. Brophy, J. J., Craven, L. A., & Doran, J. C. (2013). *Melaleucas: their botany, essential oils and uses*. Australian Centre for International Agricultural Research (ACIAR): 415pp.
29. Schwambach, J., Ruedell, C. M., de Almeida, M. R., Penchel, R. M., de Araújo, E. F., & Fett-Neto, A. G. (2008). Adventitious rooting of *Eucalyptus globulus* × *maidennii* mini-cuttings derived from mini-stumps grown in sand bed and intermittent flooding trays: a comparative study. *New Forests*, 36: 261-271.
30. Wendling, I., Brondani, G. E., Dutra, L. F., & Hansel, F. A. (2010). Mini-cuttings technique: a new ex vitro method for clonal propagation of sweetgum. *New Forests*, 39: 343-353.
31. Dirr, M. A., & Heuser, Jr., C. W. (2006). *The reference manual of woody plant propagation: From seed to tissue culture* (2nd ed.). Cary, North Carolina: Varsity Press, Inc.
32. Hartmann HT, Kester DE, Davies Junior FT, Geneve RL (2011). *Plant propagation: principles and practices*. Prentice-Hall, New Jersey.

33. Abu-Zahra, T. R., Al-Shadaideh, A. N., Abubaker, S. M., & Qrunfleh, I. M. (2013). Influence of auxin concentrations on different ornamental plants rooting. *International Journal of Botany*, 9(2): 96-99.
34. Rahdari, P., Khosroabadi, M., Delfani, K., & Hoseini, S. M. (2014). Effect of different concentration of plant hormones (IBA and NAA) on rooting and growth factors in root and stem cuttings of *Cordyline terminalis*. *Journal of Medical and Bioengineering Vol*, 3(3).
35. Costa, E. M., Loss, A., Pereira, H. P. N., & Almeida, J. F. (2015). Rooting of *Bougainvillea spectabilis* Willd. stem cuttings using indolbutyric acid. *Acta Agronomica*, 64(3): 205-210.
36. Wen, D., Gong, B., Sun, S., Liu, S., Wang, X., Wei, M., ... & Shi, Q. (2016). Promoting roles of melatonin in adventitious root development of *Solanum lycopersicum* L. by regulating auxin and nitric oxide signaling. *Frontiers in plant science*, 7: 718.
37. Gonin, M., Bergougnoux, V., Nguyen, T. D., Gantet, P., & Champion, A. (2019). What makes adventitious roots?. *Plants*, 8(7): 240.
38. Yusnita, Y., Jamaludin, J., Agustiansyah, A., & Hapsoro, D. (2017). A combination of IBA and NAA resulted in better rooting and shoot sprouting than single auxin on Malay apple [*Syzygium malaccense* (L.) Merr. & Perry] stem cuttings. *AGRIVITA, Journal of Agricultural Science*, 40(1): 80-90.
39. Husen, A., & Pal, M. (2006). Variation in shoot anatomy and rooting behaviour of stem cuttings in relation to age of donor plants in teak (*Tectona grandis* Linn. f.). *New Forests*, 31:57-73.
40. Kumar, V., Singh, M. K., Kumar, M., Prakash, S., Kumar, A., Rao, S., & Malik, S. (2015). Effect of different doses of IBA and rooting media on rooting of stem cutting

Commented [EA53]: Repetition?

- of lemon (*Citrus limon* Burm) cv. Pant lemon-1. Journal of Plant Development Sciences, 7(7): 587-591.
41. Kaur, R., & Grewal, H. S. (2017). Effect of growth regulator treatment on bud sprouting of hardwood cutting in different ornamental plants. Journal of Applied and Natural Science, 9(4): 2068-2072.
42. Sevik, H., & Guney, K. (2013). Effects of IAA, IBA, NAA, and GA3 on rooting and morphological features of *Melissa officinalis* L. stem cuttings. The Scientific World Journal.
43. Yeshiwas, T., Alemayehu, M., & Alemayehu, G. (2015). Effects of Indole Butyric Acid (IBA) and stem cuttings on growth of stenting-Propagated rose in Bahir Dar, Ethiopia. World Journal of Agricultural Sciences, 11(4): 191-197.
44. Damar, D., Barholia, A. K., Lekhi, R., & Haldar, A. (2014). Effect of growth regulators and biofertilizers on survival of pomegranate (*Punica granatum* L.) stem cuttings. Plant Archives, 14(1): 347-350.
45. Sarkar, T., Ravindra, K. N., Doss, S. G., Kumar, P. P., & Tewary, P. (2022). In vitro regeneration of mulberry plants from seedling explants of *Morus indica* cv. G4 through direct organogenesis. Trees, 36(1): 113-125.
46. Topacoglu, O., Sevik, H., Guney, K., Unal, C., Akkuzu, E., & Sivacioglu, A. (2016). Effect of rooting hormones on the rooting capability of *Ficus benjamina* L. cuttings. Šumarski list, 140(1-2): 39-44.
47. Suleiman Awwad, M. (2012). An application of the American Customer Satisfaction Index (ACSI) in the Jordanian mobile phone sector. The TQM Journal, 24(6): 529-541.

48. Sarkar, T., Sengupta, S., Kundu, S., & Das, K. (2023). Vegetative Multiplication Of *Psidium guajava* L. Through Stem Cutting. *Journal of Survey in Fisheries Sciences*, 10(1S): 754-7056.
49. Joel, P. G., Bala, M., Kaur, A., & Dubey, R. K. (2023). Efficacy of rooting hormones on propagation and rooting potential of Japanese chrysanthemums (*Chrysanthemum morifolium* Ramat.). *Rhizosphere*, 27: 100740.

UNDER PEER REVIEW