

Effect of microbial consortiums on germination and initial growth attributes of mangosteen

Abstract

An experiment was conducted to study the influence of microbial consortiums / biofertilizers on germination and initial growth attributes of mangosteen grown in the College Orchard, Department of Fruit Science, College of Agriculture, Kerala Agricultural University, Vellanikkara, Thrissur, Kerala, India during 2023. The research work was conducted under rain shelter with five treatments, which include microbial consortiums / biofertilizers such as PGPR MIX-1 (T₁), *Piriformospora indica* (T₂), PGPR MIX-1 + *Piriformospora indica* (T₃), Arka Microbial Consortia (AMC) (T₄) and control (T₅). The various parameters of germination and initial growth attributes of mangosteen seedlings were recorded viz., time taken for germination, germination percentage, height of seedlings, girth of seedlings, number of leaves, length of leaves, breadth of leaves, length of roots, girth of roots, chlorophyll content, carotenoid content, seedling fresh weight and dry weight. Among the treatments, PGPR MIX-1 + *Piriformospora indica* (T₃) recorded significantly higher values for majority of the parameters under study over control (T₅), and *Piriformospora indica* (T₂) was found to be on par with Arka Microbial Consortia (AMC) (T₄) and PGPR MIX-1 (T₁). The recorded values were lowest under control (T₅). The seeds of mangosteen treated with PGPR MIX-1 + *Piriformospora indica* or Arka Microbial Consortia alone or PGPR MIX-1 alone @ 100 g per kg of seeds were found to be the best in overall performance of the seedlings for various growth parameters under study

Key words: Mangosteen, Microbial consortium, Biofertilizer, PGPR MIX-1, *Piriformospora indica*, Arka Microbial Consortia

Introduction

The mangosteen (*Garcinia mangostana* L.) is a tropical fruit native to Southeast Asia, known for its exquisite taste and remarkable health benefits. Often hailed as "queen of fruits," mangosteen has captured the attention of fruit enthusiasts worldwide. Here's a comprehensive overview of mangosteen fruit, its characteristics, nutritional profile, health benefits, and culinary uses. Mangosteen is not only a culinary delight, but also a nutritional powerhouse. Despite its relatively low-calorie content, fruit is rich in essential nutrients, vitamins, and antioxidants. A typical serving of mangosteen (100 g) provides calories: 76, carbohydrates: 18.4 g, fibre: 1.7 g, calcium: 9 mg, phosphorus: 14 mg, iron: 0.5 mg, copper: 0.11 g, Vitamin C: 2 mg, Vitamin B1: 0.09 mg and Vitamin B2: 0.06 mg (Anon., 2004) (2). Some potential health benefits of mangosteen are the presence of antioxidants, immune support system, anti-inflammatory properties, and improvement of skin health. Mangosteen is rich in xanthenes, which are potent antioxidants that help to combat oxidative stress and inflammation in the body, thereby reducing the risk of chronic diseases such as heart disease and

cancer. Immunity supported by vitamin C content of mangosteen helps to boost immune function, supporting body's defensive mechanism against infections and illnesses. Anti-inflammatory property compounds found in mangosteen have found to help to alleviate symptoms of inflammatory conditions such as arthritis and inflammatory bowel disease.

Mangosteen propagation remains primarily reliant on seed propagation, which offers most reliable method for cultivating this prized tropical fruit tree. Successful propagation depends on the use of good quality seeds, environmental conditions, and cultural practices. Though, grafting methods were explored, they are not commonly used due to their lower success rates. By understanding the characteristics of mangosteen seeds and providing optimal germination conditions, growers can successfully use mangosteen seedlings for orchard establishment or home cultivation. Despite its challenges, seed propagation remains the primary method for expanding mangosteen cultivation and ensuring the availability of this exquisite fruit to consumers worldwide. Continued research into seed handling innovative techniques can improve the success rates of seed germination of mangosteen and also further growth and development of seedlings which will result in ensuring the sustainable production.

As mangosteen is handicapped with low percentage of germination and slow growth of seedlings it hinders propagation and overall production of this fruit crop. Hence the use of microbial consortiums/biofertilizers containing microbes, which can mobilise nutrients through biological processes from an unusable form to a useful form is found to be beneficial (Pathak *et al.*, 2017) (7). One of the most significant scientific developments in the coming ten years will be improving our understanding and management of rhizosphere processes so as to address the global concerns of climate change and population expansion (Bora *et al.*, 2016) (4). Gaining further knowledge about these biofertilizers has become essential to preserve plant health, provide food for soil-dwelling creatures, extend soil productivity, and preserving environmental biodiversity (Morrissey *et al.*, 2004) (5). By inoculating Azotobacter, nitrogenous fertiliser input can be reduced by 10-20% (Sukhada, 1999) (12). To increase soil organic carbon and also to preserve sustainability biofertilizers should be combined with chemical fertilisers and organic manures (Pathak and Kumar, 2016) (8).

An *Azospirillum* sp. may fix up to 20-40 kg N ha⁻¹, and so when inoculated with it, yield will definitely increase and the species *Azospirillum lipoferum* found to be capable of producing gibberellic acid (GA₃), which favours the growth and development of crop plants (Pathak *et al.*, 2017). Seed treatment with microbial consortiums in mangosteen could address the above said problems, and in this context research work on effect of microbial consortiums on germination and initial seedling growth attributes of mangosteen was taken up in the College Orchard, Department of Fruit Science, College of Agriculture, Vellanikkara, Thrissur.

MATERIALS AND METHODS

Research work was conducted during 2023 under rain shelter located at college orchard, Department of Fruit Science, College of Agriculture, Kerala Agricultural University, Vellanikkara, Thrissur, Kerala, India with a longitude of 76.282351° and latitude 10.55046°. The experiment was conducted in mangosteen seeds using Completely Randomized Design (CRD) with five treatments viz., T₁- PGPR MIX-1, T₂- *Piriformospora indica*, T₃- PGPR MIX-1 + *Piriformospora indica*, T₄ - AMC- Arka Microbial Consortia and T₅- Control with four replications. The plumpy seeds of mangosteen were extracted and collected from fully ripened purple coloured fruits by removing snow-white pulp and seed meat, then dried under shade for two days. Further, seeds were treated with microbial consortiums PGPR mix-1 and *Piriformospora indica* received from department of Agricultural Microbiology, College of Agriculture, Vellayani. The PGPR mix-I contains *Azospirillum lipoferum*, *Azotobacter chroococcum*, *Bacillus megaterium* and *Bacillus sporothermodurans*. Whereas, Arka Microbial Consortia (AMC) is a mixture of *Azotobacter tropicalis* strain PAN MC1, *Bacillus aryabhatai* strain Bel 6 and *Pseudomonas taiwanensis* Mpf2 (Barman *et al.*, 2016) (3) from Indian Institute of Horticulture Research (IIHR), Hesaraghatta, Bangalore. The seeds were treated with these microbial consortiums at the rate 100 g per kg of seeds *i.e.*, 10 g per 100 g of mangosteen seeds

In each replication, forty seeds were sown in poly bags of size 6x9". Observations were recorded during germination and seedling growth stage at 15 days intervals up to 45 days. Time taken for germination (days), germination percentage (%), seedling height at 15, 30 & 45 days (cm), girth of seedling at 15, 30 & 45 days (mm), number of leaves at 15, 30 & 45 days, length of leaves at 15, 30 & 45 days (cm) and breadth of leaves at 15, 30 & 45 days (cm) were recorded. By destructive sampling method at the end of experiment *i.e.*, after 45 days, various parameters recorded were chlorophyll-a (mg g⁻¹), chlorophyll-b (mg g⁻¹), total chlorophyll (mg g⁻¹), carotenoid (mg g⁻¹), length of taproot (cm), girth of taproot (mm) (0.5 cm below collar region), no. of secondary roots on taproot, seedling fresh weight and seedling dry weight. The time taken for germination was counted as number of days taken from first day of sowing to initiation of germination of seed and germination percentage was calculated by number of seeds germinated divided by total number of seeds sown multiplied by hundred. Whereas, chlorophyll-a, chlorophyll-b, and carotenoid were estimated by extraction using 80% acetone then centrifuged at 3000 rpm for 10 minutes followed by measuring the optical density (OD) values at 480, 510, 645 and 663nm by spectrophotometer. The calculation of chlorophyll-a, chlorophyll-b, total chlorophyll, ratio of chlorophyll-a/b and carotenoids were done using following formulae.

$$\text{Chlorophyll-a} = (12.7 \times \text{OD at } 663) - (2.69 \times \text{OD at } 645) \times \frac{V}{1000 \times W}$$

$$\text{Chlorophyll-b} = (22.9 \times \text{OD at 645}) - (4.68 \times \text{OD at 663}) \times \frac{V}{1000 \times W}$$

$$\text{Total chlorophyll} = (8.02 \times \text{OD at 663}) - (20.2 \times \text{OD at 645}) \times \frac{V}{1000 \times W}$$

$$\text{Carotenoids} = (7.6 \times \text{OD at 480}) - (1.49 \times \text{OD at 510}) \times \frac{V}{1000 \times W}$$

$$\text{Chlorophyll- a and b ratio} = \frac{\text{Chlorophyll-a}}{\text{Chlorophyll-b}}$$

Girth of seedlings was recorded at 15 days interval, using digital vernier caliper, which was also used for measuring the girth of taproot at the end of 45 days. The data of various parameters collected at regular intervals during seedling stage were compiled, and statistically analysed by using one factor CRD design of OPSTAT

Results and Discussion

The induction of early germination and fastening initial growth attributes of mangosteen seedlings are very challenging for the farming community engaged in mangosteen cultivation. As there existed a possibility of managing this by the use of microbial consortium / biofertilizers to a greater extent, this possibility was tried in the abovementioned experiment on mangosteen, wherein various microbial consortiums and their combinations were tested. These consortiums include PGPR MIX-1, *Piriformospora indica*, PGPR MIX-1 + *Piriformospora indica*, AMC-Arka Microbial consortia, and control (seeds that were not subjected to any seed treatment).

Table 1. Effect of microbial consortiums / biofertilizers on germination attributes of mangosteen

Treatments	Time taken for germination (days)	Germination percentage (%)
T ₁ -PGPR MIX-1	24.11	87.50
T ₂ - <i>Piriformospora indica</i>	26.63	82.50
T ₃ -PGPR MIX-1 + <i>Piriformospora indica</i>	24.16	90.00

T ₄ -AMC- Arka Microbial Consortia	23.99	90.00
T ₅ -Control	27.11	75.00
SE(m)±	0.92	11.57
CD (p=0.05)	NA	NA

When the first set of observations like time taken for germination and germination percentage were recorded during germination period (Table 1), no significant difference could be observed among the treatments

Table 2. Influence of microbial consortiums / biofertilizers on height and girth of mangosteen seedlings.

Treatments	Height of seedlings (cm)			Girth of seedlings (mm)		
	15 days	30 days	45 days	15 days	30 days	45 days
T ₁ -PGPR MIX-1	4.66	5.01	5.76	2.23	2.53	2.89
T ₂ - <i>Piriformospora indica</i>	4.11	4.51	5.08	2.14	2.50	2.61
T ₃ -PGPR MIX-1 + <i>Piriformospora indica</i>	4.81	5.64	6.34	2.32	2.64	2.99
T ₄ -AMC- Arka Microbial Consortia	4.45	5.09	5.80	2.28	2.55	2.86
T ₅ -Control	3.63	3.91	4.63	2.08	2.31	2.44
SE(m)±	0.40	0.32	0.25	0.14	0.10	0.14
CD (p=0.05)	NA	0.97	0.77	NA	NA	NA

Height of seedlings was not found to be influenced by any of the treatments under study at 15th day after germination. At 30th and 45th day, seeds treated with PGPR MIX-1 + *Piriformospora indica* (T₃) recorded significantly higher of height of mangosteen seedlings over control (T₅) and

Piriformospora indica (T₂), and was found to be on par with Arka Microbial Consortia (T₄) and PGPR MIX-1 (T₁). The control treatment recorded the least value for seedling height throughout the period of study. Girth of mangosteen seedlings did not show any significant difference throughout study period. However, seeds treated with a combination of PGPR MIX-1 + *Piriformospora indica* recorded higher values and lower values were noted in control at 15th day after germination and the same trend was observed at 30th and 45th day (Table 2).

Table 3. Effect of microbial consortium / biofertilizers on number of leaves, length of leaves and breadth of leaves of mangosteen.

Treatments	Number of leaves			Length of leaves (cm)			Breadth of leaves (cm)		
	15 days	30 days	45 days	15 days	30 days	45 days	15 days	30 days	45 days
T ₁ -PGPR MIX-1	2.22	2.82	3.32	2.93	3.91	5.04	1.91	2.43	2.57
T ₂ - <i>Piriformospora indica</i>	2.00	2.38	3.06	2.87	3.65	4.49	1.78	2.23	2.37
T ₃ -PGPR MIX-1 + <i>Piriformospora indica</i>	2.25	3.13	3.71	3.21	4.07	5.62	2.01	2.63	2.79
T ₄ -AMC- Arka Microbial Consortia	2.33	2.87	3.49	3.35	4.02	5.16	1.96	2.41	2.56
T ₅ -Control	2.00	2.13	2.62	2.68	3.08	4.18	1.58	2.08	2.16
SE(m)±	0.12	0.18	0.18	0.20	0.21	0.32	0.08	0.10	0.13
CD (p=0.05)	NA	0.55	0.56	NA	0.65	0.98	0.24	0.31	0.38

The maximum number of leaves of mangosteen seedling were significantly influenced by the treatments and values recorded were 3.13 and 3.71 at 30 and 45 days respectively in PGPR MIX-1 + *Piriformospora indica* (T₃) over control (T₅) and *Piriformospora indica* (T₂). The T₃ (PGPR MIX-1 + *Piriformospora indica*) was found to be on par with Arka Microbial Consortia (T₄) and PGPR MIX-1 (T₁) at 30th day and 45th day. No significant difference could be observed with regard to number of leaves among the treatments at 15th day of seedlings growth. Nevertheless, when higher number of leaves were recorded in T₃ (*Piriformospora indica*) and the least number of leaves were found in T₅ (control) (Table 3). With regard to length of leaves a similar trend was observed at 15th, 30th and 45th day as in the case of number of leaves, as no notable variations was noted among the treatments. During 30th day of seed germination, the treatments PGPR MIX-1 + *Piriformospora indica* (T₃), Arka

Microbial Consortia (T₄) and PGPR MIX-1 (T₁) recorded significant difference with regard to length of leaves with values of 4.07 cm, 4.02 cm, and 3.91 cm respectively. While 45th day of seed germination also the treatments recorded significantly higher values for length of leaves in PGPR MIX-1 + *Piriformospora indica* (T₃) (5.62 cm), Arka Microbial Consortia (T₄) (5.16 cm) and PGPR MIX-1 (T₁) (5.04 cm). The minimum length of leaves was recorded *i.e.*, 4.18 cm in control (T₅) and 4.49 cm in *Piriformospora indica* (T₂) (Table 3).

The influence of microbial consortium / biofertilizers on breadth of leaves of mangosteen at 15 days interval is presented in Table 3. Statistically remarkable difference with regard to breadth of leaves was observed under T₃ (PGPR MIX-1 + *Piriformospora indica*), T₁ (PGPR MIX-1) and T₄ (Arka Microbial Consortia) over T₅ (control) and T₂ (*Piriformospora indica*) during 15th, 30th and 45th day. The control, T₅ recorded the least value for breath of leaves among the treatments, which were recorded at in 15 days interval up to 45 days.

Table 4. Effect of biofertilizers on root growth and dry weight of mangosteen seedlings

Treatments	Length of taproot at 45 th day (cm)	Girth of taproot at 45 th day (mm) (0.5 cm below collar region)	No. of secondary roots on taproot at 45 th day	Seedling fresh weight at 45 th day (g)	Seedling dry weight at 45 th day (mg)
T ₁ -PGPR MIX-1	5.53	1.65	8.50	2.17	422.50
T ₂ - <i>Piriformospora indica</i>	5.03	1.42	7.25	1.83	400.50
T ₃ -PGPR MIX-1 + <i>Piriformospora indica</i>	6.19	1.90	12.00	3.33	644.75
T ₄ -AMC- Arka Microbial Consortia	5.76	1.81	10.50	2.74	546.00
T ₅ -Control	4.10	1.32	5.75	1.62	357.50
SE(m)±	0.18	0.05	0.77	0.25	40.50
CD (p=0.05)	0.55	0.14	2.34	0.75	123.18

The length of taproot at 45th day of germination varied significantly among the treatments. The treatment T₃ (PGPR MIX-1 + *Piriformospora indica*) recorded the longest taproot 6.19 cm, which

was on par with T₄ (Arka Microbial Consortia) (5.76 cm). Treatments T₁ (PGPR MIX-1), T₂ (*Piriformospora indica*), and T₅ (control) recorded remarkably lower values, when compared to T₃. However, the shortest taproot was recorded in T₅ *i.e.*, control (4.10 cm). At 45th day, treatment, T₃ and T₄ were found to be significantly superior with regard to girth of taproot and values recorded were 1.90 mm and 1.81 mm respectively. The control (T₅) recorded remarkably lower value. The treatments T₂ (1.42 mm) and T₁ (1.65 mm) recorded similar trend as observed for length of taproot. Number of secondary roots on taproot notably differed among the treatments, however T₃ (12) and T₄ (10.5) were found to be significantly superior over other treatments and were statistically on par. The lowest number was control (5.75) (Table 3).

Seedling fresh weight was found to be significantly higher in treatments T₃ (3.33 g) and T₄ (2.74 g) than in T₁ (2.17 g), T₂ (1.83 g), and T₅ (1.62 g). However, T₃ (PGPR MIX-1 + *Piriformospora indica*) and T₄ (Arka Microbial Consortia) remained statistically on par with each other, and the control recorded the lowest value for fresh weight of seedlings (Table 4). Seedling dry weight varied from 357.50 mg to 644.75 mg and the treatments exhibited significant difference. The treatment T₃ (644.75 mg) and T₄ (546.00 mg) were found to be on par with each other with significantly superior values. The control (T₅) was recorded significantly minimum value of dry weight of seedlings (357.50 mg) which was statistically on par with T₁ and T₂.

Table 5. Effect of biofertilizers on biochemical parameters of mangosteen leaf at 45th day

Treatments	Chlorophyll-a (mg g ⁻¹)	Chlorophyll-b (mg g ⁻¹)	Total chlorophyll (mg g ⁻¹)	Ratio of chlorophyll-a/b	Carotenoid (mg g ⁻¹)
T ₁ -PGPR MIX-1	0.67	0.65	1.32	1.03	0.23
T ₂ - <i>Piriformospora indica</i>	0.57	0.51	1.08	1.13	0.23
T ₃ -PGPR MIX-1 + <i>Piriformospora indica</i>	0.75	0.71	1.46	1.05	0.25
T ₄ -AMC- Arka Microbial Consortia	0.71	0.71	1.42	1.01	0.24
T ₅ -Control	0.53	0.52	1.05	1.02	0.22

SE(m)±	0.02	0.01	0.03	0.04	0.01
CD (p=0.05)	0.07	0.04	0.09	NA	0.02

The biochemical parameters of mangosteen leaves such as chlorophyll-a, chlorophyll-b and carotenoid were also estimated after 45th day and is represented in Table 5. The level of chlorophyll-a was found to be significantly higher in seeds treated with PGPR MIX-1 + *Piriformospora indica* (T₃) and Arka Microbial Consortia (T₄) with values of 0.75 mg g⁻¹ and 0.71 mg g⁻¹ respectively. Other treatments scored 0.67 mg g⁻¹, 0.57 mg g⁻¹ and 0.53 mg g⁻¹ in T₁, T₂, and T₅ respectively. All the treatments recorded similar trends for chlorophyll-b, total chlorophyll and carotenoid as recorded in chlorophyll-a. Significantly higher chlorophyll-b amount was recorded in T₃ (0.71 mg g⁻¹) and T₄ (0.71 mg g⁻¹), when compared to T₁ and T₂ remarkably lower amount of chlorophyll-b was analysed in T₁ (0.65 mg g⁻¹), T₂ (0.51 mg g⁻¹), and T₅ (0.52 mg g⁻¹). However, PGPR MIX-1 (T₁) was found to be significantly inferior to T₃ and T₄, but significantly superior over T₂ (*Piriformospora indica*) and T₅ (control). Total chlorophyll content was notably significant in treatments T₃ and T₄, with values 1.46 mg g⁻¹ and 1.42 mg g⁻¹ respectively. The least value of total chlorophyll was observed in T₅ (1.05 mg g⁻¹), which was found to be on par with T₂ (1.08 mg g⁻¹). No considerable variation was noticed among treatments with respect to ratio of chlorophyll-a/b. Carotenoid content of leaves was found to be statistically significant in T₃ (0.25 mg g⁻¹) and T₄ (0.24 mg g⁻¹) and superior over other treatments. No remarkable variation could be observed among the treatments like T₁ (0.23 mg g⁻¹), T₂ (0.23 mg g⁻¹), and T₅ (0.22 mg g⁻¹).

The aforementioned outcomes were consistent with conclusion of Barman *et.al.*, (2016) in jamun, Sindhu *et al.* (2010) (10) in horticultural crops, Pathak *et. al.*, (2009) (6) in guava, Abdelaal *et.al.*, (2010) (1) in Washington navel orange, Ramakrishnan and Selvakumar, (2012) (9) in tomato, Singh and Banik, (2011) (11) in mango cv. Himsagar and Umar *et.al.*, (2009) in strawberry cv. Chandler.

CONCLUSION

From the experiment, it was observed that significantly no difference existed among the treatments with regard to initial time taken for germination, germination percentage and seedling growth attributes at 15th day of seed germination. But as the experiment progressed

significant difference could be observed among the treatments during 30th and 45th day. Seed treatment of mangosteen with PGPR MIX-1 + *Piriformospora indica* (T₃) recorded numerically higher values for majority of the parameters under study and was on par with Arka Microbial Consortia (T₄). For some parameters T₃ and T₄ were found to be on par with PGPR MIX-1 (T₁). The performance of mangosteen seedlings grown from the seeds treated with different microbial consortiums were found to be greatly influenced by these microbial consortiums of which the combination of PGPR MIX-1 + *Piriformospora indica* (T₃), Arka Microbial Consortia (T₄), and PGPR MIX-1 (T₁) were found to have significantly positive influence on the growth of seedlings of mangosteen. From the results, it can be concluded that based on the availability of these microbial consortiums in their area, farming community can opt for PGPR MIX-1 + *Piriformospora indica* or Arka Microbial Consortia or PGPR MIX-1 they can use this at the rate of 100 g per kg of seeds for treating seeds for speeding up the growth of mangosteen seedlings.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

1. Abdelaal S, El-Sheikh MH, Hassan HAS, Kabeil SS. Microbial bio-fertilization approaches to improve yield and quality of Washington navel Orange and reducing the survival of nematode in the soil. *J Am Sci*,2010;6(12):264–271.
2. Anon. Fruits in Thailand , Department of Agricultural Extension, Ministry of Agriculture and Cooperatives , Bangkok, Thailand. 2004.
3. Barman, P., Rekha A. and Pannerselvan, P. Effect of microbial inoculants on physiological and biochemical characteristics in jamun (*Syzygiumcumini*L. Skeels) under different propagation substrates. *Int. J. Minor Fruits, Medicinal and Aromatic Plants*. 2016;2(1): 1-5.
4. Bora L, Tripathi A, Bajeli J, Chaubey AK, Chander S. A review on microbial association :its potential and future prospects in fruit crops. *Plant Arch*.2016;16(1):1–11.
5. Morrissey J, Dow J, Mark G, Ogara F. Are microbes at the root of a solution to world foodreduction. *EMBO Rep*.2004;5(10):922–926.
6. Pathak D, Singh S, Saini RS, Sharma JR. Impact of bio-inoculants on germination and plantgrowth of guava (*Psidium guajava*). *Haryana J Horti Sci*.2009;38(1&2):26–28.
7. Pathak, D.V., Kumar, M. and Rani, K. Biofertilizer application in horticultural crops. *Microorganisms for Green Revolution: Volume 1: Microbes for Sustainable Crop Production*, 2017; pp.215-227.
8. Pathak D, Kumar M. Microbial inoculants as biofertilizers and biopesticides. In: Singh DP et al (eds) *Microbial inoculants in sustainable agricultural productivity*. Springer, New Delhi. 2016
9. Ramakrishnan K, Selvakumar G. Effect of biofertilizers on enhancement of growth and yield on tomato (*Lycopersicum esculentum* Mill.) *Int J Res Bot*.2012;2(4):20–23.
10. Sindhu SS, Verma N, Dua S, Chaudhary D. Biofertilizer application for growth stimulationof horticultural crops. *Haryana J Horti Sci* 2010;39(1&2):48–70.
11. Singh SR, Banik BC. Response of integrated nutrient management on flowering, fruit setting, yield and fruit quality in mango (*Mangifera indica* L.) cv. Himsagar. *Asian J Horti*. 2011; 6(1):151–154.
12. Sukhada M. Biofertilizers for horticultural crops. *Indian Hort*.1999;44(1):32–35.