

Review Article

**ENHANCING ANIMAL NUTRITION AND SUSTAINABILITY: THE VITAL ROLE OF
GENETICALLY MODIFIED CROPS IN ANIMAL FEEDING**

Abstract

Agriculture and livestock are fundamental to the economies of developing countries. A substantial portion of crop harvests is allocated for animal feed. Thus, more technological advancements are necessary to enhance agricultural productivity and provide affordable feed. Future biotech crops are expected to play a crucial role in this area. The goal is often to introduce traits into plants that do not naturally occur in these species. These new traits may include resistance to pests, diseases, or environmental conditions, or the production of specific nutrients or pharmaceutical agents. GM crops have indirectly benefited the livestock sector by increasing the yield of feed ingredients and improving quality traits. These crops are primarily used in livestock feed rations as energy and/or protein sources. Numerous animal feeding studies have been conducted to demonstrate that genetically modified crops are as nutritious and wholesome as compared to their conventional counterparts. No biological relevant differences in animal performance, health, or animal product (meat and milk) composition had been observed in various studies conducted. Overall, no significant differences in gain, intake, and feed conversion have been reported. Since the GM crop's composition is not different from its conventional counterpart [except for the introduced transgene(s) and expressed protein(s)] and the expressed transgenic protein is rapidly digested in the digestive system, one would not expect any unintended effects. The introduction of genetically modified (GM) crops into the market undergoes extensive testing and a rigorous approval process to ensure food, feed, and environmental safety. This process includes thorough analyses before GM crops are deemed safe for commercial use. GM livestock feed is assessed for its nutritional composition and digestibility by comparing it with conventional crops. Therefore, while GM crops have the potential to enhance the efficiency of animal agriculture by improving nutritional content, reducing pesticide use, and increasing crop yields, it is essential to approach their adoption with

caution.

Keywords: Livestock, Conventional, Agriculture, Productivity, Nutrients

Commented [st1]: Inappropriate selection of keywords, rewrite

Introduction

Agriculture and livestock are fundamental to the economies of developing countries. As populations grow, the demand for livestock products like meat, milk, and eggs increases significantly. With rising urbanization and income levels in many parts of the developing world, per capita consumption of these products is expected to increase by about 2% annually [1]. Global meat demand is projected to rise by over 55% from current levels, with most of this increase occurring in developing countries [2]. A substantial portion of crop harvests is allocated for animal feed. Compound feeds for pigs, poultry, dairy cows, and other livestock are formulated using a variety of raw materials, including soya, maize, oilseed rape, cottonseed, canola, and other grains. Approximately 90% of the world's soya production is used in animal feed [3]. Consequently, the demand for feed grains is anticipated to grow by 3% annually in developing countries and by 0.5% in developed countries. On average, producing 1 kg of livestock meat requires less than 3 kg of feed grain, and producing 1 kg of milk requires less than 1 kg of feed grain [4]. Thus, more technological advancements are necessary to enhance agricultural productivity and provide affordable feed [5]. Future biotech crops are expected to play a crucial role in this area. The goal is often to introduce traits into plants that do not naturally occur in these species. These new traits may include resistance to pests, diseases, or environmental conditions, or the production of specific nutrients or pharmaceutical agents.

Commented [st2]: Rephrase???

Commented [st3]:

Commented [st4R3]: Write in passive voice i.e., on an average the production of 1kg livestock meat requires<3kg...

Commented [st5]: Introduce the idea of GM crops and significance here

Since their introduction in 1996, genetically modified (GM) crops have been used as livestock feed. GM crops have indirectly benefited the livestock sector by increasing the yield of feed ingredients and improving quality traits. These crops are primarily used in livestock feed rations as energy and/or protein sources. Conventional crops like rapeseed and mustard oil cake can be used as protein supplements for ruminants, but the presence of glucosinolate can result in a pungent smell and bitter taste after hydrolysis by endogenous enzymes [6,7]. Considering the quality concerns of the feed and other aspects like pest control and herbicide resistance, GM crops

have an advantage in feeding trials. Genetically modified crops, also known as transgenic crops, are plants whose genetic makeup has been artificially altered by inserting new genes through recombinant DNA technology [8]. The first GMO crop, glyphosate-resistant soybean, was commercialized in 1996 by Monsanto (US). Aflatoxins, toxic secondary metabolites produced by some *Aspergillus* species, are a global agricultural and health issue, causing significant crop losses annually. Host-induced gene silencing is an effective method to eliminate aflatoxin in transgenic maize. Maize plants transformed with a kernel-specific RNA interference (RNAi) gene cassette targeting the aflC gene, which encodes an enzyme in the *Aspergillus* aflatoxin biosynthetic pathway, showed no aflatoxin in kernels from these RNAi transgenic plants after pathogen infection, while nontransgenic control kernels had high toxin loads. Transcript comparisons in developing aflatoxin-free transgenic kernels and nontransgenic kernels revealed no significant differences, demonstrating that small interfering RNA can silence aflatoxin biosynthesis in maize. This precise engineering strategy can be extended to other crops to improve food security [9].

Commented [st6]: The introduction of 1st GM crop here seems irrelevant

Commented [st7]: The paragraph needs to be rephrased to be understood clearly. Re write whole para

Agricultural plants are often cited as examples of genetically modified organisms (GMOs).

Commented [st8]: Rephrase??

The benefits of genetic engineering in agriculture include increased crop yields, reduced food or drug production costs, reduced pesticide use, enhanced nutrient composition and food quality, pest and disease resistance, greater food security, and medical benefits for the growing global population. Advances have also been made in developing crops that mature faster and tolerate environmental stressors such as aluminum, boron, salt, drought, and frost, enabling plants to grow in otherwise unfavorable conditions [10]. Other applications include producing non-protein (bioplastic) or non-industrial (ornamental plant) products. The aim of this review article is to explore the critical role of genetically modified (GM) crops in meeting the rising demand for livestock products in developing countries.

Commented [st9]: Cite reference

Commented [st10]: Elaborate on bioplastic & ornamental plants & their utility wrt use of genetic engineering

Crops have been modified for various traits, as summarized in Table 1.

Table 1: Crops modified for various traits [11]

Feed Crop	Improved Traits
Alfalfa	Herbicide tolerance, modified product quality

Canola	Herbicide tolerance, modified product quality, pollination control system
Cotton	Insect resistance, herbicide tolerance
Cowpea	Insect resistance
Flax	Herbicide tolerance
Maize/corn	Modified product quality, insect resistance, herbicide tolerance, pollination control system, abiotic stress tolerance
Rice	Insect resistance, herbicide tolerance
Safflower	Modified oil/fatty acid , Antibiotic resistance
Soybean	Modified product quality, herbicide tolerance, insect resistance, altered growth/yield
Sugarcane	Insect resistance
Wheat	Herbicide tolerance

Global Scenario of GM Crops used for Feeding Livestock

Around 90% of GM crops were grown in developing countries like China, India, the Philippines, and South Africa, where a majority of farmers operate on smaller scales compared to their counterparts in the USA, the largest producer of GM commodity crops. In 2010, the USA cultivated 66.8 million hectares of GM crops, followed by Brazil with 25.4 million hectares and Argentina with 22.9 million hectares. Other significant producers with over 1 million hectares include India, Canada, China, Paraguay, Pakistan, South Africa, and Uruguay. Soybean dominates as the leading GM crop in the Americas, accounting for more than half of global GM production by volume, followed by GM maize, which constitutes about one-third of global GM production, primarily from the Americas. Canada leads in GM oilseed rape production, while Brazil, India, and China contribute substantially to GM cotton production.

The main GM crops modified for agronomic traits include soybean (36.5 million hectares), maize (12.4 million hectares), cotton (6.8 million hectares), and canola (3.0 million hectares) [12]. These crops are typically modified for traits like herbicide tolerance and insect resistance. They are utilized in livestock production as sources of energy and/or protein in various forms such as

Commented [st11]: Ref????

Commented [st12]: Ref???

Commented [st13]: Ref???

whole crop (maize silage), specific components (maize grain), or co-products like oilseed meals. Oilseed meals, in particular, derived from GM crops, are extensively used in livestock feed. For instance, in 2002, it was estimated that the global production of soybean exceeded 150 million tonnes, with approximately 50% of the area planted with GM soybean, resulting in around 35 million tonnes of GM soybean meal being used in the livestock industry. Additionally, significant quantities of maize grain, canola, cottonseed meal, and maize silage are also integrated into livestock rations.

Table 2: Cultivation of GMP in countries with the highest portions (by FAOSTAT 2019)

Country	GMP-Area (million ha)	Most cultivated plants
USA	71.5	Soybean, Maize, Cotton, Rape seed, Sugar beet, Alfalfa
Brazil	52.8	Soybean, Maize, Cotton, Sugar cane
Argentina	24.0	Soybean, Maize, Cotton, Alfalfa
Canada	12.5	Rape seed, Maize, Soybean, Sugar beet, Alfalfa,
India	11.9	Cotton
Paraguay	4.1	Soybean, Maize, Cotton
China	3.2	Cotton, Papaya
South Africa	2.7	Maize, Soybean, Cotton
Pakistan	2.5	Cotton

Commented [st14]: Center align???

GM Crops as Feed Ingredients for Farm Animals in India

In India, which boasts 20% of the world's livestock population growing at an annual rate of 0.66%, competition for land between human food production and animal fodder is intensifying. Currently, only 4% of India's total cultivable area is dedicated to fodder production, leading to significant deficits in green fodder (35.6%), dry crop residues (10.5%), and concentrate feed ingredients (44%)[13]. Efforts to address these challenges have seen recognition since the Earth

Commented [st15]: Rephrase

Summit in 1992, acknowledging that modern agricultural biotechnology alone cannot fully solve the complexities of future food security.

The adoption of GM technology in India began with initiatives like Monsanto's offer in the early 1990s to transfer insect-resistant Bt cotton technology. This move aimed to combat significant losses faced by smallholder cotton farmers due to pests like the bollworm complex. Despite concerns raised by environmental activists and scientists about the suitability of GM crops for human consumption and their potential negative impacts on health and soil quality, GM adoption has been steadily progressing. Recently, the Indian government approved the import of 5.5 lakh tonnes of GM soymeal for poultry feed, signaling a potential shift towards wider acceptance of GM crops in livestock feed [14, 15, 16]. In Karnataka, stakeholders including the Compound Livestock Feed Manufacturers' Association of India (CLFMAI) and Karnataka Poultry Farmers and Breeders Association (KPFBA) have advocated for GM feed, emphasizing its potential benefits for meat production in India amidst declining availability of conventional soybean and maize. Numerous studies on ruminant feeding have indicated that GM crops, such as corn, soybeans, cottonseed, and alfalfa, modified for traits like insect resistance and herbicide tolerance, show no significant differences in animal performance, health, or the composition of animal products like meat and milk compared to conventional crops [17,18]. These findings support the argument that GM crops, with their specific transgenic traits, do not pose unintended effects when digested by animals.

As discussions continue around the adoption of GM soybean and maize in India's livestock feed industry, stakeholders are optimistic about convincing policymakers of the benefits of GM technology for enhancing agricultural productivity and ensuring food security in the country.

Effect of Feeding of GM crops in Ruminants

Numerous ruminant feeding studies have been conducted to demonstrate that genetically modified crops are as nutritious and wholesome as compared to their conventional counterparts [17,18]. No biological relevant differences in animal performance, health, or animal product (meat and milk) composition had been observed in various studies conducted. —Table 3 provides a description of studies where GM crops were fed to ruminants. The GM crops included corn, corn silage, soybeans, soybean meal, cottonseed, fodder beets and alfalfa. The GM traits included a

Commented [st16]: Ref??

Commented [st17]: Elaborate more on benefits of Bt cotton and then move to reiterate drawbacks & issues faced by GM crops

Commented [st18]: The studies reported no significant diff. in productivity parameters so why have the associations emphasized on benefits in meat production. There is contradiction in stating reports. Do you mean the composition is not altered but the production of products is increased??? Kindly explain well, the paragraph cannot be clearly understood

Commented [st19]: The paragraph explains that no ill effects of GM was observed, but what is the actual significance to use these crops instead of conventional feed?

variety of insect-protected and herbicide-tolerant traits or a combination of them. Overall, no significant differences in gain, intake, and feed conversion were reported. Since the GM crop's composition is not different from its conventional counterpart [except for the introduced transgene(s) and expressed protein(s)] and the expressed transgenic protein is rapidly digested in the digestive system, one would not expect any unintended effects.

Table 3: Effect of Feeding of GM crops in Ruminants

Commented [st20]: Introduce separate column for Series, change the heading, since no effect is noted so the heading cannot be Effects of feeding....

GM crop	Animal	Duration of study and Experimental design	Studied health parameters	Results	Reference
1. Bt maize (MON810)	Lactating cows	25 months GM-maize group Control group	Body condition score, Immune response	No adverse effect on performance	[19]
2. GM maize and GM soybean	10 days old bull	90 days Non-GM maize and SBM Non-GM maize and GM SBM GM maize and non-GM SBM GM maize and GM SBM	Performance parameters	No adverse effects of GM components	[20]
3. Bt cotton	Twenty crossbred (KS and KF) multiparous cow	1. Control (n=10) 2. Treatment (n=10)	Milk yield, voluntary feed intake, milk composition	No effect on mean voluntary DM intake, Improved body weight in Bt group No effect on average milk yield, milk composition	[21]

4. Bt maize (MON810) and RR soybean (MON-40-30-2)	1-week old calves	<p>12 weeks</p> <p>Group I – control, conventional soybean meal and conventional maize</p> <p>Group II – GM soybean meal and conventional maize</p> <p>Group III – conventional soybean meal and GM maize</p> <p>Group IV – GM soybean meal and GM maize</p>	Haematology, Immune response	No effect of GM crops	[22]
5. Bt cottonseed	lactating Murrah buffaloes	<p>35 days</p> <p>20 Murrah buffaloes in mid-lactation</p> <p>1. 39.5% non-transgenic cottonseed in concentrate mixture</p> <p>2. 39.5% Bt cottonseed in concentrate mixture</p>	DMI, performance, blood biochemical constituents	No significant difference in the DMI, Body weight, TEC, Hb content and PCV similar. Plasma glucose, serum total proteins, albumin, globulin, triglycerides and high density lipoprotein similar	[23]
6. Bt maize (MON810) and RR soybean	10-day old calves	80 days	Histopathology	No effect of GM feed	[24]

(MON-40-3-2)	<p>Group 1-control, conventional SBM and conventional maize</p> <p>Group 2-GM SBM and conventional maize</p> <p>Group III – conventional SBM and GM maize</p> <p>Group IV – GM SBM and GM maize</p>			
--------------	---	--	--	--

Effect of Feeding of GM crops in Pigs

Table 4 summarises the effects of feeding various genetically modified crops to different age groups of pigs. No adverse effects had been observed on the performance of the animals.

Table 4: Effect of Feeding of GM crops in Pigs

GM crop	Animal	Duration of study and Experimental design	Studied parameters	Result	Reference
1. Bt maize (MON810)	35-day old male pigs	31 days GM maize group (n=16) Control (n=16)	Immune response, histopathology, serum biochemistry, organ weight	No visual effects observed on histology or serum biochemistry	[25]
2. Bt maize (MON810) and RR soybean (MON-40-30-2)	Fattening pigs	GM-maize group (n=6) GM-soybean groups (n=6)	Haematology	No effect of GM feed observed	[22]

		GM-maize AND GM-soybean group(n=6) Control (n=6)			
3. Btmaize(MON810) and RR soybean (MON-40-3-2)	Pregnant sows and offspring	Gestation and lactation GM-maize group(n=6sows, 65piglets) GM-soybean group(n=6sows, 66piglets) GM-maize & GM-soybean group(n=6sows, 69 piglets) Control (n=6sows, 65 piglets)	Haematology	No effect of GM feed observed	[26]
4. Bt maize (MON810)	Offspring and sows	Gestation and lactation GM-maize group (n = 12 sows, 160 piglets) Control (n = 12 sows, 140 piglets)	Haematology, immune response, serum biochemistry, gastrointestinal microbiota, organ weight (offspring only)	Lower granulocyte count and percentage at birth in offspring from GM-fed animals	[27]

Table 5: Effect of Feeding of GM crops in Poultry

GM crop	Animal	Duration of study and Experimental design	Studied parameters	Result	Reference
1. Bt maize (Cry1Ab)	10 days Japanese quails	49 days GM maize grp Non GM grp	body weight, hematology, serum chemistry, relative organ weight	No effect of GM feed observed	[28]
2. Bt sugarcane (Cry1Ac)	2 week old broilers	120 days GM-sugarcane grp (n=10) Control (n=10)	Serum biochemistry, histopathology, organ weight	No effect of GM feed observed	[29]
3. Bt maize (MON810) and RR soybean (MON-40-30-2)	25-week old laying hens	30 weeks GM-maize grp (n=24) GM-soybean grp (n=24) GM-maize & GM-soybean grp (n=24) Control (n=24)	Histopathology	No effect of GM feed observed	[24]
4. Bt maize	Newly hatched Japanese quail	22 weeks GM-maize grp (n=120) Control (n=120)	Immune response	Higher serum zinc concentrations in GM-maize fed animals	[30]
6. Bt maize (MON810) and RR soybean (MON-40-30-2)	6-week old Japanese quail	10 weeks per gen. GM-maize gen. GM-soybean gen. Control gen.	Clinical examination, histopathology, organ weight	Higher relative weight of breast muscle and gizzard in GM-maize fed animals	[31]

Hence, feeding trials conducted to examine the safety and efficacy of GM feeds for farm livestock indicated that there was no evidence of significantly altered nutritional composition, deleterious effects (Table 5). Animals perform in comparable manner when fed biotechnological feed ingredients.

Negative effects of feeding GM crops

Feeding genetically modified (GM) crops to animals raises concerns about potential negative effects across several fronts. These crops may introduce allergens or toxins not present in their non-modified counterparts, posing risks of allergic reactions or toxicity to animals. Modifications can also alter the nutritional content of crops, potentially impacting animal health and growth. Moreover, GM crops often include antibiotic-resistant genes, which could transfer to gut bacteria in animals, contributing to antibiotic resistance. The environmental impact is also a concern, as the cultivation of GM crops, particularly herbicide-resistant varieties, may increase herbicide use, affecting biodiversity and ecosystems crucial for animal habitats. Despite rigorous testing, the long-term effects of consuming GM crops on animal health and the environment remain uncertain, necessitating continued research and monitoring to assess these potential risks thoroughly. Studies have been conducted which shows the negative effects of feeding GM crops to animals (Table 6).

Commented [st21]: It is evident that no ill effects are noted, however the significance of GM over conventional feed is still not emphasized or clear in the whole manuscript. kindly emphasize more on significance of GM

Commented [st22]: Ref????

Commented [st23]: Ref???

Commented [st24]: Rephrase line and introduce this elsewhere

Table 6: Negative effects of feeding GM crops

GM crop	Animal and experiment. Pd.	Exp. Grp.	Parameters studied	Results	Reference
Bt maize (MON810)	Lactating cows (25 months)	GM-maize group (n=18) Control group (n=18)	Body condition score, immune response	Lower body condition	[19]
Bt maize (NK603, MON863, MON810)	Weaned piglets (22.7 weeks)	GM-fed maize& GM soybean group (n=84) Control group (n=84)	Serum biochemistry, histopathology, organ weight	Severe inflammation in stomach	[32]

and RR soybean					
Bt maize (CBH 351 Starlink)	3-month old pigs	GM-maize group (n=20) Control group (n=20)	Till end of fattening period Haematology, serum biochemistry, histopathology	Higher BUN Lower glucose	[33]
Bt maize (MON810)	Nulliparous sows and offspring	Gestation and lactatio n GM-maize group (n =12 sows, 160 piglets) Control (n=12 sows, 140 piglets)	Haematology, immune response, serum biochemistry, gastrointestinal microbiota, organ weight (offspring only)	Lower granulocyte percentage on day 110 of gestation in GM- fed animals as well as in offsprings Higher serum creatinine	[34]

Safety Assessment of GM Feeds in Livestock

The introduction of genetically modified (GM) crops into the market undergoes extensive testing and a rigorous approval process to ensure food, feed, and environmental safety. This process includes thorough analyses before GM crops are deemed safe for commercial use. GM livestock feed is assessed for its nutritional composition and digestibility by comparing it with conventional crops [35]. Agronomic, phenotypic, and compositional analyses are crucial for establishing substantial equivalence between GM and conventional crops. While these comparisons do not constitute a safety assessment *per se*, they help identify any differences and similarities.

Critical safety assessments also involve studying the properties of proteins produced by introduced genes in GM crops, particularly focusing on their potential toxicity and allergenicity when used as animal feed [36]. Studies are conducted to examine the effects of feeding GM crops on animals and on animal products such as meat, milk, and eggs (Table 7). The presence of foreign DNA from transgenic crops in animal tissues and derived products like milk, meat, and eggs is also evaluated. Studies have shown that transgenic DNA is not detectable in these food products derived from animals fed with GM crops, reaffirming safety standards.

The World Health Organization (WHO) has concluded that consuming DNA, including that from GM crops, poses no inherent risk, as mammals regularly consume DNA from various sources such as plants, animals, bacteria, parasites, and viruses. Scientific evidence supports that transgenic DNA and proteins expressed in GM crops are rapidly degraded in the animal digestive system and during feed processing. Furthermore, studies on various animals, including beef cattle, swine, sheep, fish, dairy cows, and chickens, have consistently shown no adverse effects on animal performance when fed with GM crops.

Commented [st25]: Ref???

Commented [st26]: Ref????

Detection of transgenic DNA in animal tissues plays a crucial role in assessing the impact of GM crops on animal health and the environment. Techniques such as Polymerase Chain Reaction (PCR), quantitative PCR (qPCR), Southern blotting, and fluorescent in situ hybridization (FISH) are employed to detect and quantify transgenic DNA. These methods provide insights into the persistence and potential integration of transgenes in animal genomes, as well as the possibility of horizontal gene transfer to gut microorganisms or other species. Continued advancements in molecular biology techniques contribute to ongoing research and regulatory assessments aimed at ensuring the safety and environmental sustainability of GM crop adoption in agriculture.

Table 7: Effects of Feeding GM Crops on Animal Products

TYPE OF STUDY	ANIMAL SPECIES	TISSUES SAMPLED	DETECTION OF DNA		REFERENCE
			Transgene	Positive tissue	

Feeding GM maize & soybean Bt& herbicide tolerant genes	Dairy cow	Milk	No		[37]
Feeding Phytase transgenic corn	laying hens	digestive tract, blood, heart, liver, spleen, kidney, breast muscle, eggs	Yes	gizzard	[38]
Feeding Bt maize, glyphosate tolerant soya	pig	blood	No		[26]
Feeding Bt maize	cow (dairy)	blood	No		[39]
Feeding glyphosate tolerant canola	pig	GIT content, and GIT tissues kidney, liver, spleen	Yes	entire GIT content, and GIT tissues	[40]
Feeding Bt cottonseed glyphosate tolerant cottonseed	cow (dairy)	milk	No		[41]
Feeding Bt maize	chicken (broiler)	blood, GIT content, heart, kidney, liver, muscle spleen, thymus	No		[42]
Feeding Bt maize	chicken (broiler)	kidney, liver, muscle, spleen	No		[43]
Feeding Bt maize, glyphosate tolerant soya	cow (dairy)	blood, feces, intestinal and rumen content,	Yes	Rumen	[44]

		milk			
--	--	------	--	--	--

From several studies, it has been concluded that, transgenic DNA occasionally detected in animal fluid and organ samples. Also, no transgene DNA has been reported in animal-derived edible products such as milk or eggs.

Coordinated regulatory framework on GM crops in India

The National Biotechnology Board, established in 1982, issued safety guidelines in 1983 for conducting biotechnological research in laboratories and contained settings. In 1986, it was elevated to the Department of Biotechnology (DBT) under the Ministry of Science and Technology (MOST) [45]. Initially, DBT focused on global biotechnology developments, formulated safety guidelines, and promoted the use of relevant biotechnologies in India. Recognizing the need for robust biosafety assessments, biodiversity protection, and environmental risk evaluations [46], DBT transferred oversight of genetically modified organisms (GMOs), hazardous

Commented [st27]: Rephrase the sentence in passive voice

Commented [st28]: Rephrase the sentence in passive voice




Fig 1 : Regulatory framework on GM crops in India

The Government of India, under the Allocation of Business Rules 1961, assigned responsibilities for biodiversity conservation and environmental protection to MOEF (Government of India, 1961). Consequently, MOEF began regulating GMOs and their products under the Environment Protection Act (EPA) 1986. The EPA Rules 1989 were introduced under the hazardous substances section of EPA 1986, categorizing GM crops and GMOs as potentially harmful substances akin to hazardous materials impacting human, animal, or environmental health.

The regulatory framework established a tiered system with six competent authorities: Policy Advisory Committees such as the Recombinant DNA Advisory Committee (RDAC); Regulating and Approval Committees like the Institutional Biosafety Committee (IBSC), Review Committee on Genetic Manipulation (RCGM), and Genetic Engineering Approval Committee (GEAC); and Post-Monitoring Committees including the State Biotechnology Coordination Committee (SBCC) and District Level Committee (DLC). These committees were mandated to oversee different aspects of GMO regulation as outlined in the EPA Rules 1989. While DBT played a role in biosafety assessments under these rules, MOEF primarily managed post-monitoring activities at the state level.

However, the roles of the Ministry of Health and Ministry of Agriculture, crucial for regulating seed quality and human health in relation to GM crops, were not clearly defined under the EPA Rules 1989. Over time, the biosafety regulatory system evolved into a comprehensive framework involving multiple ministries, each administering specific legislative acts like the Environment Protection Act 1986 by MOEF, Seed Act 1966 and Seeds (Control) Order by Ministry of Agriculture (MOA), and Food Safety and Standards Act 2006 by Ministry of Health and Family Welfare (MOH&FW). The EPA Rules 1989 remained central to biosafety regulations for GM crops, while other laws focused on food safety and seed quality.

Commented [st29]: Plagiarism???

Guidelines for safety in biotechnology, including specific guidelines for GM crops, were developed by DBT starting with the Recombinant DNA Safety Guidelines in 1990, revised in 1994, and the comprehensive Guidelines for Research in Transgenic Plants and Guidelines for Toxicity and Allergenicity. Evaluation of Transgenic Seeds, Plants, and Plant Parts in 1998. These guidelines emphasized case-by-case biosafety evaluations, risk assessments for environmental impacts, and agronomic performance tailored to specific crops, traits, and agro-ecological systems[47]. They also addressed safety protocols for imported GM materials.

Commented [st30]: Plagiarism???

Concerns about GM crops encompass health risks, environmental hazards, and economic issues. These include potential toxicity from unexpected mutations, allergenicity risks, environmental impacts like non-target insect effects from Bt toxins, and economic disparities due to high seed costs. Despite potential benefits like enhanced agricultural efficiency and reduced pesticide use, the adoption of GM crops necessitates careful consideration of these complex and multifaceted issues.

Conclusion

Rising population in developing countries, and certainly in India, requires extraordinary steps to intensify agriculture production to meet the growing demand by substantially increasing crop yield for utilization of both humans and animals. It has now been recognized that neither the current agriculture production system nor the modern biotechnology alone can solve the complex challenges of feeding the world of tomorrow. The use of genetically modified crops in animal feeding is hence a complex and multifaceted topic with both potential benefits and concerns. While GM crops have the potential to enhance the efficiency of animal agriculture by improving nutritional content, reducing pesticide use, and increasing crop yields, it is essential to approach their adoption with caution.

Commented [st31]: Repetition??? Re write

References

1. Delgado CL, Rosegrant MW, Steinfeld H, Ehui SK, Courbois C. Livestock to 2020: The next food revolution. Intl Food Policy Res Inst; 1999.
2. Rosegrant MW, Paisner MS, Meijer S, Witcover J. 2020 Global food outlook: Trends, alternatives, and choices. Intl Food Policy Res Inst; 2001.
3. Brown L. The United States and China, the soybean connection. Worldwatch Institute, November. 1999 Nov;9:1999.
4. Schmitz JA. Department of Veterinary and Biomedical Sciences: 2001 Annual Report.
5. FAO W. Principles and methods for the risk assessment of chemicals in food. Environmental Health Criteria. 2009; 240.
6. Pailan G H & Singhal K K, Effect of dietary glucosinolates on nutrient utilization, milk yield and blood constituents of lactating goats, *Small Ruminants Res*, 71 (1997)31-37.
7. Papas A, Ingalls JR, Campbell LD. Studies on the effects of rapeseed meal on thyroid status of cattle, glucosinolate and iodine content of milk and other parameters. *The Journal of nutrition*. 1979 Jul 1;109(7):1129-39.
8. Chandler S, Tanaka Y. Genetic modification in floriculture. *Critical Reviews in Plant Sciences*. 2007 Aug 6;26(4):169-97.
9. Thakare D, Zhang J, Wing RA, Cotty PJ, Schmidt MA. Aflatoxin-free transgenic maize using host-induced gene silencing. *Science Advances*. 2017 Mar 10;3(3):e1602382.
10. Takeda, S., & Matsuoka, M. Genetic approaches to crop improvement: Responding to environmental and population changes. *Nature Reviews Genetics* 9, 444–457.
11. Briefs IS. Global status of commercialized biotech/GM crops in 2017: Biotech crop adoption surges as economic benefits accumulate in 22 years. ISAAA brief. 2017;53:25-6.
12. James C. ISAAA Briefs brief 49 Global Status of Commercialized Biotech/GM Crops: 2014.
13. Department of animal husbandry dairying, ministry of fisheries, animal husbandry and dairying. Government of India: New Delhi, India. 2019.
14. Ghosh PK. Bt-cotton: government procedures. *Current science*. 2001; 80(3):323-324.
15. Karihaloo JL, Kumar PA. Bt Cotton in India. A status report. 2009:1-58.

16. Kranthi KR. Bt cotton-questions and answers. Indian Society for Cotton Improvement (ISCI), Mumbai; 2012 Jan 1.
17. Clark JH, Ipharraguerre IR. Biotechnology crops as feeds for livestock. 2003; 866: 177-198.
18. Flachowsky G, Chesson A, Aulrich K. Animal nutrition with feeds from genetically modified plants. Archives of Animal Nutrition. 2005 Feb 1;59(1):1-40.
19. Guertler P, Brandl C, Meyer HH, Tichopad A. Feeding genetically modified maize (MON810) to dairy cows: comparison of gene expression pattern of markers for apoptosis, inflammation and cell cycle. Journal für Verbraucherschutz und Lebensmittelsicherheit. 2012 Sep;7:195-202.
20. Furgał-Dierżuk I, Strzetelski J, Kwiatek K, Twardowska M, Mazur M, Sieradzki Z, Kozaczyński W, Reichert M. The effect of genetically modified maize (MON 810) and soyabean meal (Roundup Ready) on rearing performance and transfer of transgenic DNA to calf tissues. Journal of Animal and Feed Sciences. 2014 Mar 10;23(1):13-22.
21. Singhal KK, Tyagi AK, Rajput YS, Prashad S. Effect of feeding cottonseed produced from cotton hybrids carrying Cry1C gene on feed intake, milk production and composition in lactating crossbred dairy cows. Report submitted to Metahelix Life Sciences Ltd, Bangalore. National Dairy research Institute, Karnal. 2008.
22. Bednarek D, Dudek K, Kwiatek K, Świątkiewicz M, Świątkiewicz S, Strzetelski J. Effect of a diet composed of genetically modified feed components on the selected immune parameters in pigs, cattle, and poultry. Journal of Veterinary Research. 2013 Jan 1;57(2):209-17.
23. Singh M, Tiwari DP, Kumar A. Effect of transgenic cottonseed in lactating Murrah buffaloes. Buffalo Journal. 2003;19(1):117-26.
24. Reichert M, Kozaczyński W, Karpińska TA, Bocian Ł, Jasik A, Kycko A, Świątkiewicz M, Świątkiewicz S, Furgał-Dierżuk I, Arczewska-Włosek A, Strzetelski J. Histopathology of internal organs of farm animals fed genetically modified corn and soybean meal. Journal of Veterinary Research. 2012 Jan 1;56(4):617-22.
25. Walsh MC, Buzoianu SG, Gardiner GE, Rea MC, Ross RP, Cassidy JP, Lawlor PG. Effects of short-term feeding of Bt MON810 maize on growth performance, organ morphology and function in pigs. British Journal of Nutrition. 2012 Feb;107(3):364-71.

26. Świątkiewicz M, Bednarek D, Markowski J, Hanczakowska E, Kwiate K. Effect of feeding genetically modified maize and soybean meal to sows on their reproductive traits, haematological indices and offspring performance. *Journal of Veterinary Research*. 2013 Oct 30;57(3):413-8.
27. Walsh MC, Buzoianu SG, Gardiner GE, Rea MC, O'Donovan O, Ross RP, Lawlor PG. Effects of feeding Bt MON810 maize to sows during first gestation and lactation on maternal and offspring health indicators. *British journal of nutrition*. 2013 Mar;109(5):873-81.
28. Zhang L, Shen W, Fang Z, Liu B. Effects of genetically modified maize expressing Cry1Ab and EPSPS proteins on Japanese quail. *Poultry Science*. 2021 Feb 1;100(2):1068-75.
29. Hameed A, Nasir IA, Tabassum B, Qamar Z, Zameer M, Younus M, Rao A, Rashid B, Tariq M, Khan GA, Ali M. Biosafety assessment of locally developed transgenic sugarcane. *JAPS: Journal of Animal & Plant Sciences*. 2016 Aug 1;26(4).
30. Scholtz ND, Halle I, Dänicke S, Hartmann G, Zur B, Sauerwein H. Effects of an active immunization on the immune response of laying Japanese quail (*Coturnix coturnix japonica*) fed with or without genetically modified *Bacillus thuringiensis*-maize. *Poultry science*. 2010 Jun 1;89(6):1122-8.
31. Korwin-Kossakowska A, Sartowska K, Tomczyk G, Prusak B, Sender G. Health status and potential uptake of transgenic DNA by Japanese quail fed diets containing genetically modified plant ingredients over 10 generations. *British Poultry Science*. 2016 May 3;57(3):415-23.
32. Carman, Judy A., Howard R. Vlieger, Larry J. Ver Steeg, Verlyn E. Sneller, Garth W. Robinson, Catherine A. Clinch-Jones, Julie I. Haynes, and John W. Edwards. "A long-term toxicology study on pigs fed a combined genetically modified (GM) soy and GM maize diet." 2013: 38-54.
33. Yonemochi C, Suga K, Harada C, Hanazumi M. Tevaluation of transgenic event CBH 351 (StarLink) corn in pig. *Animal science journal*. 2010 Feb;81(1):94-101.
34. Buzoianu SG, Walsh MC, Rea MC, Quigley L, O'Sullivan O, Cotter PD, Ross RP, Gardiner GE, Lawlor PG. Sequence-based analysis of the intestinal microbiota of sows and their offspring fed genetically modified maize expressing a truncated form of *Bacillus*

- thuringiensis Cry1Ab protein (Bt maize). *Applied and Environmental Microbiology*. 2013 Dec 15;79(24):7735-44.
35. Novak WK, Haslberger AG. Substantial equivalence of antinutrients and inherent plant toxins in genetically modified novel foods. *Food and Chemical Toxicology*. 2000 Jun 1;38(6):473-83.
 36. Kuiper HA, Kleter GA, Noteborn HP, Kok EJ. Assessment of the food safety issues related to genetically modified foods. *The plant journals*. 2001 Sep;27(6):503-28.
 37. De Giacomo M, Di Domenicantonio C, De Santis B, Debegnach F, Onori R, Brera C. Carry-over of DNA from genetically modified soyabean and maize to cow's milk. *Journal of Animal and Feed Sciences*. 2016 May 19;25(2):109-15.
 38. Ma Q, Gao C, Zhang J, Zhao L, Hao W, Ji C. Detection of transgenic and endogenous plant DNA fragments and proteins in the digesta, blood, tissues, and eggs of laying hens fed with phytase transgenic corn. *PLoS One*. 2013 Apr 8;8(4):e61138.
 39. Bertheau Y, Helbling JC, Fortabat MN, Makhzami S, Sotinel I, Audeon C, Nignol AC, Kobilinsky A, Petit L, Fach P, Brunschwig P. Persistence of plant DNA sequences in the blood of dairy cows fed with genetically modified (Bt176) and conventional corn silage. *Journal of agricultural and food chemistry*. 2009 Jan 28;57(2):509-16.
 40. Sharma R, Damgaard D, Alexander TW, Dugan ME, Aalhus JL, Stanford K, McAllister TA. Detection of transgenic and endogenous plant DNA in digesta and tissues of sheep and pigs fed Roundup Ready canola meal. *Journal of Agricultural and Food Chemistry*. 2006 Mar 8;54(5):1699-709.
 41. Castillo AR, Gallardo MR, Maciel M, Giordano JM, Conti GA, Gaggiotti MC, Quaino O, Gianni C, Hartnell GF. Effects of feeding rations with genetically modified whole cottonseed to lactating Holstein cows. *Journal of Dairy Science*. 2004 Jun 1;87(6):1778-85.
 42. Marzok MA. Detection of genetically modified soybeans and maize in Egypt as well as comparative nutritional safety investigations of isogenic and transgenic (Bt) maize in broiler nutrition: broiler performance, degradation and metabolic fate of maize DNA in some tissues and organs. 2004 Sep 8.
 43. Jennings JC, Albee LD, Kolwyck DC, Surber JB, Taylor ML, Hartnell GF, Lirette RP, Glenn KC. Attempts to detect transgenic and endogenous plant DNA and transgenic

- protein in muscle from broilers fed YieldGard Corn Borer Corn. *Poultry Science*. 2003 Mar 1;82(3):371-80.
44. Phipps RH, Deaville ER, Maddison BC. Detection of transgenic and endogenous plant DNA in rumen fluid, duodenal digesta, milk, blood, and feces of lactating dairy cows. *Journal of Dairy Science*. 2003 Dec 1;86(12):4070-8.
45. Sharma M, Charak KS, Ramanaiah TV. Agricultural biotechnology research in India: Status and policies. *Current Science*. 2003 Feb 10;84(3):297-302.
46. Ghosh PK. Transgenic plants and biosafety concerns in India. *Current science*. 1997 Feb 10;72(3):172-9.
47. Tripathi KK, Behera UN, VibhaDhawan. Agri-biotechnology in India: Biosafety Issues, Experiences and Expectations. *Agriculture for Food Security and Rural Growth*. 2008.