

Effect Of Different Liquid Organic Manures On The Nutrient Release Pattern In Radish Grown Soil

ABSTRACT

The study of nutrient release patterns from various organic manures is critical for ensuring enough nutrient availability to crop plants at the right time and in the right quantity. To assess the nutrient release pattern from several organic manures, including farm yard manure, panchagavya and jeevamrutha, a field experiment was conducted at organic farming block of Zonal Agricultural Research Station, V. C. Farm, Mandya during late Kharif 2020. The experiment consists of 9 treatments including different rates of recommended dose of nitrogen was applied through FYM and one and two times application of panchagavya and jeevamrutha in different combinations. The experiment was laid out in a Randomized Block Design with three replications. The release pattern of available nitrogen, phosphorus, potassium, calcium, magnesium, sulphur and micronutrients in soil applied with various organic manures at various days (15, 30, 45 DAS and at harvest). The study revealed that application of various organic manures showed substantial increase in primary (N, P and K) and secondary nutrient (S) release and there was no significant variations found in Ca, Mg, Fe, Mn, Cu and Zn release pattern. Based on the release rate of nutrients, it was found that there was consistent and stable release of N, P, K and S from 75% RDN through FYM + two time application of Panchagavya followed by 75% RDN through FYM + two time application of Jeevamrutha, compared to other treatments.

KEYWORDS

Panchagavya, Jeevamrutha, Radish, Nitrogen, Phosphorus, Potassium and Sulphur release rate

1 INTRODUCTION

Radish is root vegetable crop and it is grown for its young tuberous roots where even shoots are used as vegetable and consumed as either cooked or raw as salad. Different colored radishes are available in the market. The pink color is due to the presence of anthocyanin pigment. It is relished for its pungent flavor and is considered as an appetizer (Kumar *et al.*, 2014). Radish is a good source of vitamin-C and supplies variety of minerals like calcium, potassium and phosphorous. The pink skin radish is generally rich in ascorbic acid than white skin radish (Singh and Bhandari, 2015). The characteristic pungent flavor of radish is due to isothiocyanate (Kushwah, 2016). It is used for neurological headaches, chronic diarrhea, urinary complaints, sleeplessness, and piles (Singh and Bhandari, 2015).

After the onset of the Green Revolution in India, farmers began utilizing chemical fertilizers to boost the production and productivity of different crops. The inadequate use of chemical fertilizers has adversely affected the soil fertility, quality, biodiversity, human and soil health. Furthermore, it also increased soil acidity, deteriorated soil physical condition, decreased organic matter and created micronutrient deficiencies (Mali *et al.*, 2018). To solve these problems, the future of agriculture should be averted to organic farming. Because organic manure not only provides vital nutrients (both macro and micro nutrients), but it also improves the physical properties like bulk density, water holding capacity, infiltration rate, soil aeration and biological properties like improving microbial population and enzyme activity (Diacono and Mon-temurro, 2010) . However, single organic source of nutrient supplementation may not cope up with the nutrient demand of crops. Integration of different organic nutrient sources and/or liquid organic manures help to solve dual problem of supplementation of sufficient nutrients besides synchronized nutrient availability as per crop demand associated with variable nutrient release pattern among different organic manures. Hence, the study was conducted to assess the effect of different liquid organic manures on the nutrient release pattern in radish grown soil.

2 MATERIALS AND METHODS

Present investigation was carried out at at organic farming block of Zonal Agricultural Research Station (ZARS) V. C. Farm, Mandya, University of Agricultural Sciences, Bangalore during late *Kharif* 2020. The experiment was laid out in the Randomized Block Design with nine treatments which were replicated three times. The treatment employed were, T₁ (100% RDN through FYM), T₂ (50% RDN through FYM + one time application of Jeevamrutha), T₃ (50%

RDN through FYM + two time application of Jeevamrutha), T₄ (75% RDN through FYM + one time application of Jeevamrutha), T₅ (75% RDN through FYM + two time application of Jeevamrutha), T₆ (50% RDN through FYM + one time application of Panchagavya), T₇ (50% RDN through FYM + two time application of Panchagavya), T₈ (75% RDN through FYM + one time application of Panchagavya) and T₉ (75% RDN through FYM + two time application of Panchagavya). Recommended dose of Nitrogen (RDN- 50 kg ha⁻¹) was supplied through FYM in equal proportion on N content basis prior to 15 days of sowing. Two liquid formulations (Panchagavya and Jeevamrutha) were applied one at the time of sowing and second at 30 days aftersowing at the rate of 500 L ha⁻¹.

A composite soil sample was collected from the experimental site (0-20 cm) before the experiment and was analyzed for chemical properties. Experimental soil was sandy loam in texture. The soil was neutral in reaction (pH 7.30) and medium in soluble salts (0.29 dS m⁻¹) and medium in organic carbon (5.24 g kg⁻¹) content. Available nitrogen content of soil was low (302.24 kg ha⁻¹), medium in available P₂O₅ (43.30 kg ha⁻¹), available K₂O (210.66 kg ha⁻¹) and available sulphur (15.12 mg kg⁻¹). The exchangeable calcium and magnesium content of soil was 7.01 and 2.21 cmol (p⁺) kg⁻¹, respectively. DTPA extractable iron, manganese, zinc and copper were 19.85, 10.69, 0.49 and 1.26 mg kg⁻¹, respectively. The FYM, Panchagavya and Jeevamrutha were analyzed with respect to their chemical composition by adopting standard procedures and analytical data are presented in Table 1.

Chemical properties of the soil such pH, EC, OC, available N, P₂O₅, K₂O, Ca, Mg, S, Fe, Mn, Zn, and Cu were determined as per the standard methods. Statistical analysis of the data recorded was done as per the method suggested by Gomez and Gomez (1984). The significance of variation among the treatments was observed by applying ANOVA. The level of significance used in “F” was P = 0.05. Critical difference (CD) values were calculated for the P = 0.05 whenever “F” test was found significant.

Table 1: Chemical characters of Panchagavya, Jeevamruth and FYM

Parameter	Panchagavya	Jeevamrutha	FYM
Colour	Light brown	Light green	-
Odour	Fermented odour	Mild foul odour	-
pH	6.02	7.17	9.4
EC (dS m ⁻¹)	3.40	1.48	7.2

OC (%)	0.88	0.69	13.98
Nitrogen (mg L⁻¹)	1240	910	0.49
Phosphorus (mg L⁻¹)	196	156	0.17
Potassium (mg L⁻¹)	888	632	0.46
Calcium (mg L⁻¹)	156	178	0.82
Magnesium (mg L⁻¹)	46	20	0.49
Sulphur (mg L⁻¹)	566	543	0.31
Zinc (mg L⁻¹)	1.19	3.91	13.80
Copper (mg L⁻¹)	0.34	1.82	2.4
Iron (mg L⁻¹)	27.87	29.80	142.24
Manganese (mg L⁻¹)	1.69	10.40	63.58

3 Results and discussion:

Table 2: Soil pH, electrical conductivity and organic carbon content as influenced by application of liquid organic manures at different growth stages of radish

Treatment	Available N (kg ha ⁻¹)				Available P ₂ O ₅ (kg ha ⁻¹)				Available K ₂ O (kg ha ⁻¹)			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T ₁	7.38	7.43	7.45	7.49	0.33	0.35	0.38	0.39	5.23	5.79	5.88	5.97
T ₂	7.42	7.47	7.49	7.52	0.32	0.35	0.36	0.37	5.05	5.35	5.52	5.64
T ₃	7.43	7.48	7.50	7.53	0.37	0.39	0.42	0.44	5.11	5.71	6.10	6.17
T ₄	7.44	7.51	7.51	7.56	0.36	0.38	0.39	0.41	5.15	5.75	5.83	5.89
T ₅	7.46	7.55	7.59	7.62	0.38	0.40	0.42	0.46	5.30	5.84	6.21	6.33
T ₆	7.41	7.45	7.48	7.51	0.31	0.33	0.34	0.35	5.13	5.73	5.86	5.94
T ₇	7.42	7.47	7.49	7.52	0.34	0.37	0.39	0.42	5.26	5.86	6.13	6.28
T ₈	7.40	7.43	7.46	7.48	0.33	0.34	0.35	0.36	5.18	5.78	5.89	5.91
T ₉	7.39	7.45	7.50	7.51	0.35	0.36	0.37	0.40	5.43	5.93	6.34	6.52
S.Em ±	0.33	0.34	0.34	0.34	0.02	0.02	0.02	0.03	0.23	0.26	0.27	0.27
CD@ 5%	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

The soil chemical properties *viz.*, pH, EC and OC of soil at 15, 30, 45 DAS and at harvest as influenced application of liquid organic manures are presented in table 1. The data of chemical properties *viz.*, pH, EC and OC of soil at 15, 30, 45 DAS and at harvest of radish crop

is given in table 1 which indicated that soil chemical properties at different intervals were not affected significantly due to different treatments applied in this investigation.

Slight variation in pH was observed compare to initial data in all the treatments which might be due to increase in bases by active degradation of organic matter solution and the beneficial influence of liquid organic manures that provide favourable environment for nutrient availability the results are in accordance with Ali *et al.* (2011), Kumawat *et al.* (2013) and Narkhede *et al.* (2017). Slight variation in EC was observed it might be due to faster release of bases and soluble organic fractions to the soil system by mineralization. This is similar to the findings of Ali *et al.* (2011), Kumawat *et al.* (2013) and Narkhede *et al.* (2017). Variations in organic carbon content in soil compare to initial soil data this was due to, Patel *et al.* (2018) reported that the addition of organic matter to soil increased the root biomass production which in turn increases the carbon content in soil. Narkhede *et al.* (2017) reported that addition of farm waste and organic manures increased the status of organic carbon and available NPK of the soil.

Table 3: Available N, P₂O₅ and K₂O content of soil as influenced by application liquid organic manures at different growth stages of radish

Treatment	Available N (kg ha ⁻¹)				Available P ₂ O ₅ (kg ha ⁻¹)				Available K ₂ O (kg ha ⁻¹)			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T ₁	313.02	316.89	319.22	322.22	46.73	49.06	51.61	52.94	218.66	222.26	224.66	228.33
T ₂	315.88	318.76	320.76	323.76	47.15	50.02	53.00	55.64	222.14	225.81	228.81	231.48
T ₃	316.77	319.75	323.08	326.42	48.01	50.35	55.14	58.03	224.18	226.78	231.82	234.82
T ₄	320.67	323.55	326.88	328.22	50.36	52.69	54.35	55.39	226.97	229.04	230.53	232.53
T ₅	322.65	325.53	330.53	334.20	51.56	54.48	57.16	59.39	227.44	230.84	234.17	236.84
T ₆	317.39	320.78	322.44	325.11	48.68	51.75	55.17	56.84	223.92	225.58	227.25	230.92
T ₇	318.20	321.08	325.08	329.41	49.97	52.12	56.91	58.42	224.06	226.72	229.72	233.72
T ₈	324.12	326.33	328.00	331.33	51.87	53.87	55.82	57.62	230.48	232.15	233.93	234.26
T ₉	326.58	328.46	332.13	336.46	52.11	56.11	59.65	61.92	231.13	233.46	236.13	239.46
S.Em±	2.39	1.95	1.43	1.78	1.18	1.40	1.20	1.23	2.42	1.91	1.82	1.47
CD@ 5%	7.16	5.83	4.30	5.34	3.52	4.19	3.58	3.68	7.26	5.72	5.45	4.42

Available nitrogen, phosphorus and potassium content at 15, 30, 45 DAS and at harvest of radish has been presented in table 3. The results revealed that higher soil N (326.58, 328.46, 332.13 and 336.46 kg ha⁻¹), P₂O₅ (52.11, 56.11, 59.65 and 61.92 kg ha⁻¹) and K₂O

(231.13, 233.46, 236.13 and 239.46 kg ha⁻¹) content in soil was recorded in treatment T₉ (75% RDN through FYM + two time application of Panchagavya). Significant increase in available nitrogen, phosphorus and potassium content of soil was due to the increased multiplication of microbes which mineralize the nitrogen, phosphorus and potassium contained in the soil and applied organic manures. Application of FYM also contributing in supply of nitrogen, phosphorus and potassium to soil. Characterization study of liquid organic manures indicated that among the liquid organic manures, Panchagavya registered the highest N, P and K content and nitrogen fixers, PSB population which helps in solubilizing the nitrogen and phosphorous present in soil. Several reports supporting this observation had been made by Pradeep (2013) in groundnut, Vajantha *et al.* (2013), Amareswar and Sujathamma, (2015), Yogananda *et al.* (2015), Patel *et al.* (2018) and Gangadhar *et al.* (2019).

Table 4: Exchangeable calcium and magnesium content of soil as influenced by application of organic manures at different growth stages of radish

Treatment	Exch. Ca [cmol (p ⁺) kg ⁻¹]				Exch. Mg [cmol (p ⁺) kg ⁻¹]			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T ₁	7.18	7.27	7.36	7.62	2.53	2.60	2.67	2.83
T ₂	7.25	7.46	7.81	8.09	2.59	2.70	2.81	2.90
T ₃	7.30	7.52	7.95	8.24	2.60	2.74	3.03	3.13
T ₄	7.41	7.61	7.87	8.16	2.77	2.83	2.95	3.06
T ₅	7.45	7.64	7.96	8.27	2.80	2.89	3.18	3.33
T ₆	7.28	7.40	7.74	8.01	2.62	2.76	2.97	3.09
T ₇	7.32	7.45	7.98	8.26	2.65	2.82	3.00	3.16
T ₈	7.48	7.56	8.08	8.18	2.82	3.05	3.07	3.10
T ₉	7.52	7.59	8.14	8.33	2.91	3.17	3.25	3.42
S.Em±	0.33	0.33	0.35	0.36	0.12	0.12	0.14	0.14
CD@ 5%	NS	NS	NS	NS	NS	NS	NS	NS

Exchangeable calcium and magnesium content in soil as influenced by application of liquid organic manures at 15, 30, 45 DAS and at harvest are presented in table 4. The results revealed that calcium and magnesium content in soil did not vary significantly due to application of liquid organic manures at all the stages of crop growth. According to Haynes (1986), liquid organic manures has the property of binding mineral particles like calcium and magnesium in the form of colloids of humus and clay, facilitating stable aggregates of soil particles for desired

porosity to sustain plant growth. Manjunatha *et al.* (2009) noticed that application of liquid organic manures maintains soil health and productivity by improving physical, chemical and biological properties of soil.




Fig. 1: Available sulphur content in soil as influenced by application of liquid organic manures at different growth stages of radish

Available sulphur content in soil as influenced by application of liquid organic manures at 15, 30, 45 DAS and at harvest are presented in Fig.1. Highest amount of sulphur (19.84, 23.94, 26.76 and 28.57 mg kg⁻¹, respectively) in the soil was recorded in T₉ (75% RDN through FYM + two times application of Panchagavya) and the lowest amount of sulphur in the soil (15.69, 17.11, 19.79 and 21.15 mg kg⁻¹, respectively) was recorded in T₁ (100% RDN through FYM). Increase in sulphur content might be ascribed to adsorption of S on organic matter and thereby reducing the leaching losses of sulphur. Microbial sulphur oxidation is whole beneficial to soil fertility, resulting in the formation of sulphate, the major S-ion used by plants, this was similar with the findings of Jat and Ahlawat (2010).




Fig 2: Micronutrient status of soil as influenced by application liquid organic manures at different growth stages of radish

DTPA extractable micronutrient such as, iron, manganese, zinc and copper content of soil at 15, 30, 45 DAS and at harvest of radish crop did not vary significantly due to application of liquid organic manures (Fig 2.). DTPA-Fe content ranged from 20.31, 21.59, 23.55 and 25.91 to 23.34, 25.72, 27.92 and 29.93 mg kg⁻¹, DTPA-Mn content ranged from 10.51, 14.65, 15.64 and 17.08 to 12.19, 16.63, 19.77 and 20.94 mg kg⁻¹, DTPA-Zn content ranged from 0.50, 0.54, 0.64 and 0.67 to 0.58, 0.63, 0.73 and 0.76 mg kg⁻¹ and DTPA-Cu 1.25, 1.32, 1.41 and 1.50 to 1.59, 1.65, 1.75 and 1.84 mg kg⁻¹ at 15, 30, 45 DAS and at harvest, respectively in T₆ (50% RDN through FYM + one time application of Panchagavya) and T₅ (75% RDN through FYM + two time application of Jeevamrutha).

The effect of application of liquid organic manures on soil micronutrient status did not vary significantly. However slight increase in micronutrient content was observed. The difference was only because of varied levels of micronutrient composition of Panchagavya and Jeevamrutha. Jeevamrutha showed more micronutrient content hence more increase of micronutrients was observed in 75% RDN through FYM + two times application of Jeevamrutha. The increase in available Fe, Mn, Zn and Cu upon addition of organic manures might be due to intensified microbial population and reduction in pH of soil and also formation

of stable complexes with organic ligands. This might have decreased the susceptibility of micronutrients to adsorption, fixation or precipitation reaction in soil resulting in greater availability. The increased soil micronutrient status with addition of organic manures to the soil was reported by Kumawat *et al.* (2013) and Jain *et al.* (2014).

Conclusion:

Application of organic manures including liquid organic manures like Panchagavya and Jeevamrutha provided a significant improvement on soil chemical properties and contributed to the availability of plant nutrient. The prominent increase in N, P₂O₅, K₂O, and is a benefit from the application of the different liquid organic manures, even though pH, EC, OC, Ca, Mg and micronutrients were not affected by the application of liquid organic manures. Clearly, liquid organic manures can be a useful tool to improve soil fertility and nutrient status in soil.

REFERENCE:

- ALI, M. N., GHATAK, S. AND RAGUL, T., 2011, Biochemical analysis of panchagavya and sanjibani and their effect in crop yield and soil health. *J. Crop Weed.*, **7** (2): 84-86.
- AMARESWARI, P. U. AND SUJATHAMMA, P., 2015, Effect of vermicompost, jeevamrutha, panchagavya and mulching on the cost benefit ratio in french bean. *Hort. Flora Res. Spectr.*, **4** (4): 333-336.
- DIACONO, M. AND MONTEMURRO, F., 2010, Long term effects of organic amendments on soil fertility. A review. *Agron. Sustain. Dev.*, **30**: 401–422.
- GANGADHAR, K., DEVAKUMAR, N., VISHWAJITH AND LAVANYA, G., 2019, Growth, yield and quality parameters of chilli (*Capsicum annuum* L.) as influenced by Application of different organic manures and decomposers. *Int. J. Chem. Stud.*, **8** (1): 473-482.
- GOMEZ, K. A. AND GOMEZ, A. A., 1984, Statistical procedures for agricultural research. 2nd Ed. *John Wiley Sons*, New York.

- HAYNES, R. J., 1986, The decomposition process mineralization, immobilization, humus formation and degradation. In: Haynes, R. J. (ed.), Mineral nitrogen in the plant-soil system. Academic press, New York, pp. 75-98.
- JAIN, P., SHARMA, R. C., BHATTACHARYYA, P. AND BANILE, P., 2014, Effect of new organic supplemented (panchagavya) on seed germination and soil quality Environ. *Monit. Assess*, **186** (4): 1999-2001.
- JAT, R. A. AND AHLAWAT, I. P. S., 2010, Effect of organic manure and sulphur fertilization in pigeonpea (*Cajanus cajan*) + groundnut (*Arachis hypogaea*) intercropping system. *Indian J. Agron.*, **55** (4): 276-281.
- KUMAR, S., SUTANU, M., KUMAR, S. AND SINGH, H. D., 2014, Efficacy of organic manures on growth and yield of radish (*Raphanus sativus* L.) cv. Japanese White. *Int. J. Plant Sci.*, **9** (1): 57-60.
- KUMAWAT, R. N., MAHAJAN, S. S. AND SANTRA, P., 2013, Effect of panchgavya on soil chemical properties of groundnut (*Arachis hypogaea*) rhizosphere and crop productivity in western Rajasthan. *J. Food Legumes*, **26** (2): 39-43.
- KUSHWAH, L., 2016, Effect of organic manures, inorganic fertilizers and their combinations on growth, yield and quality of radish (*Raphanus sativus* L.). M. Sc. (Ag) thesis. Rajmata Vijayaraje Scindia Krishi Vishwa Vidhyalaya, Gwalior, India.
- MALI, D. L., SINGH, V., SAROLIA, D. K., TELI, S. K., CHITTORA, A. AND DHAKAR, R., 2018, Effect of organic manures and bio-fertilizers on growth and yield of radish (*Raphanus sativus* L.) cv. Japanese white. *Int. J. Chem. Stud.*, **6** (2): 1095-1098
- MANJUNATHA, G. S., UPPERI, S. N., PUJARI, B. T., YELEDHALLI, N. A. AND KULIGOD, A. V. B., 2009, Effect of farm yard manure treated with jeevamrutha on yield attributes and economics of sunflower (*Helianthus annuus* L.). *Karnataka J. Agric. Sci.*, **22** (1): 342-348.
- NARKHEDE, W. N., KHANDARE, S. K., NAYAK AND KHAZI, G. S., 2017, Effect of organic and inorganic nutrient sources and their integrated nutrient management on

productivity, economic efficiency, available N, P, K and microbial population in soybean (*Glycine max*) – onion (*Allium cepa*) cropping sequence. *Indian J. Agron.*, **62** (2): 147–154

PATEL, D. M., PATEL, I. M., PATEL, B. T., SINGH, N. K. AND PATEL, C. K., 2018, Effect of panchgavya and jivamrut on yield, chemical and biological properties of soil and nutrients uptake by *Kharif* groundnut (*Arachis hypogaea* L.). *Int. J. Chem. Stud.*, **6** (3): 804-809.

PRADEEP GOPAKKALI, 2013, Development of liquid organic manures and their evaluation for organic production of Chilli and Onion. Ph.D. Thesis, University of Agricultural Sciences, Bangalore.

SINGH, K. P. AND BHANDARI, R. R., 2015, Vegetable crops production technology. First edition Bagbazar, Nepal. Pp. 308.

VAJANTHA, B., UMADEVI, M., PATNAIK, M. C. AND RAJKUMAR, M., 2013, Micronutrient status of soils under ashwagandha grown farmer fields in Andhra Pradesh. *Asian J. Soil Sci.*, **8** (1): 143-147.

YOGANANDA, S. B., DEVAKUMAR, N., SHRUTI, M. K. AND NINGARAJU, 2015, Growth and yield of cowpea as influenced by different sources of organic manures, *Nat. Symp. Org. Agric. Sust. Food Sec. Challenges Opp.*, Tamilnadu, India, p. 113.