

Exploring Natural Solutions: Botanical Approaches for Controlling Early Blight in Tomato Plants under Laboratory Conditions

In vitro influence of some plant extracts against mycelial growth of *Alternaria solani* which causes Late Blight of Tomatoe

Formatted: Font: (Default) Arial, 18 pt, Bold, English (United States)

ABSTRACT:

Early blight of Tomato caused by *Alternaria solani* is one of most serious and destructive disease worldwide. In the present study, the concentration-treatment viz. 10% of each plant extract was tested by poison food technique to manage the mycelial growth of the *A. solani*. All the tested extracts were significantly inhibited the mycelial growth of pathogen when compared with control. Among the tested plant studied extracts, garlic clove extract was found expressively most efficacious and indicated maximum percentage inhibition that was 90.54% at 10% concentration followed by Neem leaf extract (87.83%), Ginger (83.78%), Eucalyptus (75.67%), Bael (71.67%), Onion bulb extract (68.67%) and Ashwagandha (63.51%) respectively. While, Lantana (52.70%), was the least effective extract with 35 mm mycelial growth diameter. Mancozeb was the most effective fungicide against early blight of tomato percentage inhibition was 93.24% at 0.25% concentration, which gave the best control over against mycelial development of *Alternaria solani* after seven days of incubation at 24°C. Presently many Many farmers are using chemicals chemical fungicides for control of to control early blight of tomato however, they which cause health many hazards especially for human health. Therefore, this investigation will be helpful to the farmers for selection of suitable plants extract, used to use as control chemical fungicides alternatives of for tomatoe's early blight of tomato. control.

KEYWORDS: Early blight of Tomato, *Alternaria solani*, Plant extracts, Natural fungicides, biological control, Garlic clove extract, Neem leaf extract

Introduction

Tomato is the an important vegetable crop, and grown in world wide. It's fruits are is very good important source of many vitamins like A, B,C and some minerals viz. Iron, calcium, potassium, phosphorus and sulphur. Tomato crop is highly susceptible to plant pathogens viz. bacteria, fungi, and viruses which deteriorate the yield and quality of tomato food crop [18, 19]. Among the fungal diseases Tomaro's early blight is caused by *Alternaria solani* is important fungal disease (Ellis and Martin) (Biradar et al., 2021) [1,7]. During all stages of plant development, *A. solani* causes disease (leaf blight, stem rot, fruit lesions) and results in severe damage across the country [4, 18]. Early blight is most catastrophic disease incurring loss under field and post-harvest stages causing 50 to 86% reduction in fruit yield [19, 9].

The This disease is managed by the use of several conventional fungicides Neha et al. (2021) [8, 14, 22], but due to the development of resistance in most common pathogenic fungi against fungicides and also the factor of exposure risks fungicide residues and human health hazards have given a push for obtaining alternatives to control *A. solani*. It is very essential to determine the efficacy of botanicals agents and their integrations against early blight of tomato (Reference). In addition, given the increasing self-awareness on the sides of producers and consumers and the increasing increase tendency towards organic products, it is obvious that botanical plants will gain much more importance in the context of integrated pathogen control management [5, 15, 20, 21]. In this study, it was aimed to determine the availability of botanical plant extracts which are eco-friendly and unharmed to health of human and other living creatures as botanical control against early blight disease pathogen *A. solani* which causes significant yield losses in tomato.

Formatted: Font: Not Italic

MATERIALS AND METHODS

The experiment was conducted in vitro at the Department of Plant Pathology and Agricultural microbiology, College of Agriculture, Pune, during 2017-2018, in order to evaluate the efficacy of different plants extracts against early blight of tomato.

Isolation of pathogen

Infected tomato plant showing symptoms of early blight was collected from the field and washed thoroughly with water then after placed in between blotting paper to remove excess moisture. The infected leaf was cut aseptically into small pieces and surface sterilized with mercuric chloride (0.1%) for 10 sec. followed by two to three washed-washing with sterilized distilled water and then placed aseptically on solidified potato dextrose agar (PDA) in petri plates (90 mm diameter??). Inoculated petri plates were incubated in BOD incubator at 25°C 24??°C for mycelial growth of pathogen. The pure culture of this fungus was sub cultured and maintained for further investigation.

Preparation of plants Extract

The plants parts (which part of every plant) were grinded with sterilized water and filtered through a muslin cloth to produce a 100% crude extract. The required quantity of each plant extract was added separately so as to get a requisite concentration by using potato dextrose agar which was used as nutrient medium. The plant extracts were carefully mixed by stirring and about 10 ml extract contained medium was poured to each of the 9 cm petri dishes and allowed for solidification. Potato dextrose agar plate without the plant extract was used as control. Inoculated plates were incubated at 26.25±1 °C in BOD Incubator for twelve days and the colony diameter was recorded and the percent inhibition was calculated using the formula(reference?).

$$I = \frac{C-T}{C} \times 100$$

I = percent inhibition of mycelial growth, C = radial growth of fungus in control,
T = radial growth of fungus in treatment

RESULTS AND DISCUSSION

Methods: In-vitro assessment of plant extracts and fungicide against Alternariasolani was done by poisoned food technique. The treatments evaluated after seven days of colony growth by taking average mycelial growth and inhibition percentage. In present study, all the All of the tested tested extracts at 10% concentration were significantly ($P \leq 0.01$) reduced linear growth and increased inhibition percentage compared to control. Among the tested plant extracts, garlic clove extract was found expressively most efficacious and indicated maximum percentage inhibition that was 90.54% at 10% concentration followed by Neem leaf extract (87.83%), Ginger (83.78%), Eucalyptus (75.67%), Bael (71.67%), Onion bulb extract (68.67%) and Ashwagandha (63.51%) respectively. While, Lantana (52.70%), was the least effective extract with 35 mm mycelial growth. Mancozeb was most effective fungicide against early blight of tomato percentage inhibition was 93.24% at 0.25% conc. gave the best control over mycelial development.



Fig 1. In vitro evaluation of botanicals and fungicide against *Alternariasolani*
You must explain the Fig.

In case of plant extracts, garlic was found to be significant and gave maximum percentage inhibition i.e. 90.54 at 10%. While neem was the second most effective extract followed by ginger, onion and eucalyptus at 10% concentration which was found to be at par with each other. The

Formatted: Line spacing: single

Formatted: Tab stops: 0.9", Left

Formatted: Font: Italic

results revealed that the concentrations of all the botanicals compressed the growth of *Alternariasolani*, however, the garlic application at 10% concentration revealed determined control over the mycelial development of *Alternariasolani* after seven days of incubation at 24°C. The inhibitory effect of these extracts may be due to their direct lethal effect on the pathogen growth or antimicrobial activity against fungal pathogens under *invitro*.

The result of present study is in accordance to the finding of Deshmukh (2020)[6, 7], Naik et al (2020) [13, 14] and Biradar et al. (2021) [3] tested plant oils and plants products clove extract of *Allium sativum* (10%) recorded the highest reduction of the mycelial growth of 100% followed by leaf extract (10%) of Daturametel (68.44%) and oil (3%) of *Azadirachtaindica* (59.88%) Biradar et al. (2021) and Sahu *et al.* (2014) [2,10] tested plant extracts produced some antifungal activities, whereas *Azadirachtaindica* (neem), *Daturastramonium* (datura) and *Withaniasomnifera* (ashwagandha) showed significant antifungal activities. The leaf extract of *W. somnifera* was the most effective in inhibiting the mycelial growth of *A. solani* (62.56%) followed by *D. stramonium* (34.65%) and *A. indica* (25.27%) [11].

Table1. *In vitro* evaluation of plant leaf extracts and fungicide influence against *Alternariasolani*

* = Mean of four replications, Figure in parenthesis are Arcsine transfer values(Arcsine transfer values are not necessary.

Tr.No.	Treatments	Mean colony diameter* (mm)	Inhibition over control (%)
1	Ginger	12.02 (20.69)	83.78
2	Garlic	6.9 (15.21)	90.54
3	Ashwgandha	27.1 (31.42)	63.51
4	Lantana	35 (36.22)	52.70
5	Neem	8.93 (17.43)	87.83
6	Onion	23 (28.73)	68.91
7	Eucalyptus	18.3 (24.98)	75.67
8	Bael	21.1 (27.49)	71.67
9	Mancozeb	5 (12.91)	93.24
10	Control	74.25 (59.5)	00
SE(m)±		0.44	
CD (5%)		1.29	
<u>must be at 1% in vitro</u>			

Formatted Table




Fig. 2 Inhibition of colony growth of *Alternariasolani* over control

CONCLUSION

Early blight of tomato is very worst disease and yield and quality losses due to this disease. In the view of summarized experimental finding, it may conclude that, antifungal properties are present in the plant extracts. Suitable plants extracts can be used as antifungal against myciliial growth of *Alternariasolani*. The use of fungicide (harmful chemicals) can lead health hazards for farmers as well as consumers and it also harms the plant and fruits. Plant extract of garlic and neem were highly effective against pathogen. There are different types of plants available in the nature with medicinal properties. However, some plant extracts are effective and some are not effective against the studied pathogen. Therefore, this investigation will be helpful to farmers for selection of suitable and effective plant extract for managing Early blight disease of tomato.

REFERENCES

- Biradar PD, Suryawanshi AP, Patait NN. In vitro bio efficacy of bioagents against *Alternariasolani* and *Colletotrichumcapsici*, causing tomato fruit rot. The Pharma Innovation Journal. (2021);10(12):1773-1776.
- Biradar PD, Suryawanshi AP, Patait NN. In vitro bio-efficacy of phyto extracts against *Alternariasolani* and *Colletotrichumcapsici*, causing tomato fruit rots. The Pharma Innovation journal. (2021);10(12):1784-1787.
- Biradar PD, Suryawanshi AP, Giri VV. In vitro bio efficacy of essential oils against *Alternariasolani* and *Colletotrichumcapsici*, causing tomato fruit rots. ThePharma Innovation Journal. (2021);10(12):1671-1674.
- Jones LR, Grout AJ. Notes on two species of *Alternaria*. Bulletin of the Torrey Botanical Society. 1897; 24:254- 258.
- Kadam NP, Mulekar V. G., Biradar Pratiksha D.* and Suryawanshi A. P. In vitro Efficacy of Aqueous and Solvent Phytoextracts against *Ralstoniasolanacearum* (Poster Presentation) National symposium on Probing beneficial microorganism for next green revolution, Akola, India (2021).
https://www.researchgate.net/publication/370108833_Thematic_Area_Novel_approaches_for_the_management_of_plant_diseases_In_vitro_Efficacy_of_Aqueous_and_Solvent_Phytoextracts_against_Ralstonia_solanacearum
- Agrios GN. Plant Pathology. 5th edition, Alsevier Academic Press, New Yark, 2005, 665.
- Mathur K, Shekhawat KS. Chemical control of early blight in Kharif sown tomato. Indian Journal of Mycology Plant Pathology. 1986; 16:235-238.
- Patil MJ, Ukeyand SP, Raut BT. Evaluation of Fungicides and Botenicals for the management of Early Blight (*Alternariasolani*) of tomato. PKV Research Journal. 2002; 25(1):49-51.

9. Nivedha M, Ebenezer EG, Kalpana K, Kumar R. *In vitro* antifungal evaluation of various plant extracts against leaf blight disease of *Jasminum grandiflorum* caused by *Alternaria alternata* (Fr.) Keissler, Journal of Pharmacognosy and Phytochemistry. 2019; 8(3):2143- 2147.
10. Sahu DK, Khare CP, Patel R. Eco friendly management of early blight of tomato using botanical plant extracts. Journal of Industrial Pollution Control. 2014; 30(2):215- 218.
11. Arunakumara, K.T. (2006). Studies on *Alternariasolani* (Ellis and Martin) Jones and Grout causing early blight of tomato. M. Sc (Ag) thesis, University of Agricultural Sciences, Dwarwad.
12. Chourasia, P. K., A. A. Lal and S. Simon, (2013.) Effect of fungicides and botanicals against early blight of tomato caused by *Alternariasolani*. *International Journal of Agricultural Science Research* 3(3), 151-155.
13. Nashwa, and M.A. Sallam, (2012). Control of tomato early blight disease by certain aqueous plant extracts. *Plant Pathology Journal* 10(4): 187-191.
14. Neha NP, AP Suryawanshi, PD Biradar, VR Wadhawe. Efficacy of Fungicides against *Phyllosticta zingiberi*, causing *Phyllosticta* leaf spot of ginger in *In vitro* conditions. *Pharma Innovation Journal* 2021;10(12):1788-1792.
15. Neha NP, AP Suryawanshi, PD Biradar, VR Wadhawe, MR Kharade. Bioefficacy of Bioagents and Phytoextracts against *Phyllosticta zingiberi*, causing *Phyllosticta* leaf spot of ginger in *In vitro* conditions. *Pharma Innovation Journal* 2021;10(12):1804-1809.
16. Singh, P. C. and D. Singh. (2006). *In vitro* evaluation of fungicides against *Alternaria alternata*. *Ann. Pl. Protec. Sci.*, 14 (2): 500-502.
17. Deshmukh HV, Deokar CD, Khaire PB and Brahmane PR. Efficacy of different botanicals against the *Alternariasolani* under *in vitro* conditions. *Journal of Pharmacognosy and Phytochemistry*. (2020) 9(6): 1986-1989.
18. Khaire PB, Mane SS and Pawar SV. Identification and Management of Fungal Diseases of Tomato-A review. 1 (1) *Agri meet Multidisciplinary e-Magazine*. (2021) 1-18. https://www.researchgate.net/publication/348322121_IDENTIFICATION_AND_MANAGEMENT_OF_FUNGAL_DISEASES_OF_TOMATO-A_REVIEW.
19. Khaire PB and Hake LG. Some Important Post Harvest Diseases of Tomato and Their Management. *Popular Kheti*. (2018) 6 (3) 80-86. https://www.researchgate.net/publication/333203668_Some_Important_Post_Harvest_Diseases_of_Tomato_and_Their_Management.
20. Naik SC, Narute TK, Narute TT and Khaire PB. Effects of selected biological control agents and plant extracts on early blight disease management and yield of cherry tomato. *Journal of Pharmacognosy and Phytochemistry*. (2020) 9(5): 587-592. https://www.researchgate.net/publication/348884009_Effects_of_selected_biological_control_agents_and_plant_extract_on_early_blight_disease_management_and_yield_of_cherry_tomato.
21. Naik SC, Narute TK, Narute TT and Khaire PB. *In vitro* efficacy of plant extract (botanicals) against *Alternariasolani* (early blight of tomato). *Journal of Pharmacognosy and Phytochemistry*. (2020) 9(5): 614-617. https://www.researchgate.net/publication/348884330_In_vitro_efficacy_of_plant_extract_botanicals_against_Alternaria_solani_early_blight_of_tomato.
22. Deshmukh HV, Deokar CD, Raghuvanshi KS, Khaire PB and Brahmane PR. Efficacy of different fungicides against the *Alternariasolani* under *in vitro* conditions. *J Pharmacogn Phytochem* 2020;9(6):1957-1960.

Insert Page number