

## Original Research Article

### Yield and yield-attributing traits of some bitter gourd genotypes ~~ASSESSMENT OF BITTER GOURD GENOTYPES FOR YIELD AND YIELD ATTRIBUTING TRAITS~~

#### ABSTRACT

The present study was undertaken at the Department of Horticulture, University of Agricultural Sciences, GKVK Bengaluru, India during the *Rabi* season of 2022 using a Randomized Complete Block Design (~~RCBD~~)—with three replicationsto evaluate ~~eighteen~~ 18 bitter gourd genotypes, including the check Pusa Hybrid 6, for various yield and yield attributing traits, highlighting significant variations among them. The analysis of variance revealed significant differences among the genotypes for the studiedtraits. Genotype Pusa Purvi produced the maximum number of fruits per vine (46.67), whereas, Pusa Rasdar produced the least number of fruits (12.33) per vine. The longest fruits were recorded in Pant Karela 4 (24 cm) followed by Phule Green Gold (19.37 cm) and Priya (19 cm), whereas, the smallest fruits were found in Pusa Purvi (5.07 cm).The genotype Pusa Rasdar exhibited the maximum fruit diameter (6.53 cm) and highest fruit weight (148.14 g), while Pusa Purvi exhibited the minimum diameter (2.72 cm). The genotype Konkan Tara(1.96 kg) followed by Pusa Rasdar (1.79 kg) and Priya (1.60 kg) exhibited the maximum fruit yield per vine. These genotypes may be selected as parents in further breeding programme to improve the crop in terms of yield traits.

**Keyword:** bitter gourd; variability; evaluation; yield; traits

**Comment [R1]:** Don't use words that are already there in the title as key words.

#### 1. INTRODUCTION

Bitter gourd (*Momordica charantia* L.) holds a prominent position as a cucurbitaceous vegetable cultivated throughout India during the warm season for its tender fruits, which is commonly called as Balsam pear or bitter cucumber or bitter melon (English), Hagalakayi (Kannada), Karela (Hindi), etc. The genus *Momordica* encompasses several other species including *M. balsamina*, *M. cochinchinesis*, *M. dioca*, *M. denudate*, *M. macrocarpa*, *M. subangulata*, and *M. tuberosa*. Bitter gourd is diploid having chromosome number  $2n=22$  and has its origin in the Indo-Burma region [14-17].

**Comment [R2]:** Add author citations for the binomials.

**Formatted:** Font: Not Italic

Leading countries in bitter gourd production include India, Indonesia, Malaysia, China, and Tropical Africa. Major bitter gourd cultivating states in India are Madhya Pradesh, Chhattisgarh, Tamil Nadu, Andhra Pradesh, Odisha, and Bihar. Its cultivation is widespread in India's tropical and sub-tropical regions, with a total area of 109.51 thousand hectares, production of 1375.82 thousand tonnes and a productivity of 12.56 MT (Anon., 2022).

Bitter gourd is known for its various health benefits, including its reported wormicidal effect and its role as a laxative and digestion stimulant. The bitter taste of bitter gourd is attributed to certain compounds, including the cucurbitacin-like alkaloid momordicine and

**Comment [R3]:** Cite suitable references for these benefits.

triterpene glycosides such as momordicoside K and L (Jeffrey, 1980; and Okabe *et al.*, 1982). The antidiabetic properties of bitter melon have garnered significant attention in recent years. Extensive research over the past decade has revealed the presence of a hypoglycemic compound called "charantin" (Yeh *et al.*, 2003). In terms of vitamins, it is a source of 126.0 mg of vitamin A and 96.0 mg of vitamin C (Gopalan *et al.*, 1993).

Bitter melon, with its monoecious nature and high degree of heterozygosity, primarily undergoes cross-pollination. Its medicinal and nutritional value have increased its commercial significance, leading to consistent consumer demand throughout the year. Given its wide variability, monoecious nature, conspicuous and accessible flower architecture for crossbreeding, and high seed production per fruit, bitter melon can serve as a potent material for exploiting heterosis on a commercial scale (Rani, 2014). Therefore, the objectives of the study were to evaluate the bitter melon genotypes for their yield and yield attributing traits to identify the best genotypes that can be utilized for further breeding programme.

## 2. MATERIALS AND METHODS

The experiment was conducted in the eastern dry zone of Karnataka at the Department of Horticulture, University of Agricultural Sciences, Bengaluru, India. The experimental site was located at 12° 58' North latitude and 77° 35' East longitude with an altitude of 830 m above Mean Sea Level (MSL). The experimental site majorly consisted of red sandy loam soil with pH of 6.56 and the average rainfall in this area is 679.1 to 888.9 mm. Eighteen bitter melon genotypes including check Pusa Hybrid 6 collected from various sources (Table 1-) for the experiment were evaluated during the Rabi season of 2022 using a Randomized Complete Block Design (RCBD) with three replications. Bitter melon seeds which were pre-soaked for about 12 hrs were sowed in beds at a spacing of 2.0 x 2.0 meters under open field condition by adopting recommended package of practices (Plate Fig. 1). Data recorded on five randomly selected plants from each genotype for yield traits viz., fruit length (cm), fruit diameter (cm), average fruit weight (g), number of fruits per vine, and yield per vine were statistically analysed by following the method given by Panse and Sukhatme (1954).

**Table 1. List of bitter melon genotypes along with their source of collection**

Sl.No.	Genotypes	Source of collection
1	Punjab 14	NSC, New Delhi
2	Hirkani	MPKV, Rahuri
3	Pusa Rasdar	IARI, New Delhi
4	Pusa Purvi	IARI, New Delhi
5	Priya	KAU, Thrissur
6	Konkan Tara	KKV, Dapoli
7	Konkan Karali	KKV, Dapoli

**Comment [R4]:** Delete the column serial number.

**Comment [R5]:** Spell out the acronyms in the foot note.

8	Pant Karela 3	GBPUAT, Pantnagar
9	Pant Karela 4	GBPUAT, Pantnagar
10	CO 1	TNAU, Coimbatore
11	Preethi	KAU, Thrissur
12	Phule Green Gold	MPKV, Rahuri
13	Pusa Do Mausami	IARI, New Delhi
14	Pusa Vishesh	IARI, New Delhi
15	Pant Karela 1	GBPUAT, Pantnagar
16	Pusa Hybrid 4	IARI, New Delhi
17	Pusa Hybrid 5	IARI, New Delhi
18	Pusa Hybrid 6 (Check)	IARI, New Delhi



**Plate-Figure1:** General view of the experiment plot

### 3. RESULTS AND DISCUSSION

The mean values for all genotypes, including the check Pusa Hybrid 6, across yield attributing traits are summarized in Table2. Significant variations were observed among the genotypes for all the traits studied (Plate-Fig.2). Genotype Pant Karela 4 (~~24.00 cm~~) exhibited the longest fruits, followed by Phule Green Gold (~~19.37 cm~~), Priya (~~19.00 cm~~), and Pusa Do Mausami (~~18.43 cm~~), all surpassing the check Pusa Hybrid 6 (~~18.33 cm~~). Conversely, Pusa Purvi (~~5.07 cm~~) produced the smallest fruits among the evaluated genotypes. As the length of the fruit increases, the total yield is also expected to increase in a positive direction. Similar findings were reported by Ullah *et al.* (2012), Ranjan *et al.* (2015) in cucumber and Hossain *et al.* (2016), Yadav *et al.* (2018) in bitter gourd.

The maximum fruit diameter was recorded in Pusa Rasdar (~~6.53 cm~~), followed by Konkan Tara (~~5.24 cm~~) and Pusa Vishesh (~~4.68 cm~~), with percentage reductions over the control of 43.20%, 14.19%, and 2.63%, respectively. Whereas, Pusa Purvi (~~2.72 cm~~) exhibited the minimum fruit diameter with a reduction of -40.35% over the control. These findings align closely with those reported by Rani (2014) and Khan *et al.* (2015) in bitter gourd.

Among the genotypes evaluated for average fruit weight, Pusa Rasdar succeeded in achieving a maximum fruit weight (~~148.14 g~~) which was significantly higher than all the

**Comment [R6]:** Don't repeat the values that are already there in the table again in the results.

other genotypes. Additionally, average fruit weight was observed in Pant Karela 4 (~~79.93 g~~), Konkan Tara (~~74.70 g~~), Phule Green Gold (~~70.42 g~~), Priya (~~66.69 g~~), and Konkan Karali (~~68.83 g~~). The genotypes such as Punjab 14 (~~66.10 g~~), Hirkani (~~64.62 g~~), CO 1 (~~65.00 g~~), Preethi (~~65.00 g~~), Pusa Vishesh (~~60.00 g~~), and Pant Karela 1 (~~58.00 g~~) were on par with the control Pusa Hybrid 6 (~~61.40 g~~). In contrast Pusa Purvi (~~15.33 g~~) exhibited the smallest fruit weight, with a substantial reduction of -75.03% compared to the control. The significant variations in fruit weight might have been due to fruit length, fruit width, and number of fruits per vine. Similar findings are reported by Mallikarjunarao *et al.* (2020) in bitter gourd.

The highest number of fruits per vine was observed in Pusa Purvi (~~46.67~~) followed by Konkan Tara (~~26.33~~) and Priya (~~24.33~~) whereas, the lowest was recorded in Pusa Rasdar (~~12.33~~). The results revealed that the genotypes with a smaller average fruit weight recorded highest number of fruits per vine and the variation in number of fruits per vine might be due to fruit set percentage, sex ratio, and vine length.

The fruit yield per vine ranged from 0.62 kg to 1.96 kg, with an average of 1.15 kg per vine across the genotypes. Konkan Tara exhibited the highest yield per vine (~~1.96 kg~~), followed by Pusa Rasdar (~~1.79 kg~~), Priya (~~1.60 kg~~), and Punjab 14 (~~1.54 kg~~). Genotypes such as Hirkani (~~1.26 kg~~), Konkan Karali (~~1.35 kg~~) and Pant Karela 4 (~~1.29 kg~~) showed yields per vine similar to that of control Pusa Hybrid 6 (~~1.26 kg~~). In contrast, Pusa Hybrid 5 recorded the lowest yield per vine (~~0.62 kg~~). The significant variation in yield per vine might be due to fruit set percentage, sex-ratio, fruit length, number of fruits per vine, fruit weight and fruit width. These findings were supported by the observations of Yadav *et al.* (2004) in bitter gourd.

Yield components such as fruit length, fruit diameter, fruit weight and number of fruits per vine are critical factors that significantly influence the total fruit yield per vine in bitter gourd. The evaluated bitter gourd genotypes showed significant phenotypic variability for fruit weight, fruit length, fruit diameter, number of fruits per vine and yield per vine. The variations in yield and yield-related traits among genotypes can be attributed to their distinct morphological and physiological features, which influence the distribution of carbohydrates from the source to the sink, resulting in differing yields.

**Table 2. Mean performance of bitter gourd genotypes for yield and yield attributing traits**

Sl. No.	Genotypes	Fruit length (cm)	Percent reduction over control	Fruit Diameter (cm)	Percent reduction over control
1	Punjab 14	14.60	-20.35	3.97	-12.94
2	Hirkani	14.00	-23.62	4.31	-5.48
3	Pusa Rasdar	13.50	-26.35	6.53	43.20
4	Pusa Purvi	5.07	-72.34	2.72	-40.35
5	Priya	19.00	3.66	4.16	-8.77
6	Konkan Tara	15.83	-13.64	5.24	14.91
7	Konkan Karali	12.83	-30.01	2.74	-39.91

Comment [R7]: Are there any previous observations in this line?

Comment [R8]: Cite studies in support of this statement.

Comment [R9]: Delete the column serial number.


8	Pant Karela 3	17.33	-5.46	3.26	-28.51
9	Pant Karela 4	24.00	30.93	3.26	-28.51
10	CO 1	10.33	-43.64	3.24	-28.95
11	Preethi	13.17	-28.15	3.97	-12.94
12	Phule Green Gold	19.37	5.67	3.98	-12.72
13	Pusa Do Mausami	18.43	0.55	3.01	-33.99
14	Pusa Vishesh	16.43	-10.37	4.68	2.63
15	Pant Karela 1	15.34	-16.31	3.53	-22.59
16	Pusa Hybrid 4	8.52	-53.52	3.62	-20.61
17	Pusa Hybrid 5	11.52	-37.15	4.20	-7.89
18	Pusa Hybrid 6 ( <del>Cheek</del> Control)	18.33	0.00	4.56	0.00
	<b>C.V</b>	<b>3.72</b>	-	<b>2.98</b>	-
	<b>SE(m)</b>	<b>0.32</b>	-	<b>0.07</b>	-
	<b>C.D @ 5%</b>	<b>0.92</b>	-	<b>0.20</b>	-

**Table 2. (contd.)Mean performance of bitter gourd genotypes for yield and yield attributing traits**

Sl No.	Genotypes	Average fruit weight (g)	Percent reduction over control	No. of fruits per vine	Percent reduction over control	Yield per vine (kg)	Percent reduction over control
1	Punjab 14	66.10	7.65	23.33	14.76	1.54	22.22
2	Hirkani	64.62	5.24	19.33	-4.92	1.26	0.00
3	Pusa Rasdar	148.14	141.27	12.33	-39.35	1.79	42.06
4	Pusa Purvi	15.33	-75.03	46.67	129.56	0.70	-44.44
5	Priya	66.69	8.62	24.33	19.68	1.60	26.98
6	Konkan Tara	74.70	21.66	26.33	29.51	1.96	55.56
7	Konkan Karali	68.83	12.10	19.67	-3.25	1.35	7.14
8	Pant Karela 3	55.82	-9.09	16.33	-19.68	0.91	-27.78
9	Pant Karela 4	79.93	30.18	16.33	-19.68	1.29	2.38
10	CO 1	65.00	5.86	15.67	-22.92	1.00	-20.63
11	Preethi	65.00	5.86	14.67	-27.84	0.94	-25.40
12	Phule Green Gold	70.42	14.69	15.33	-24.59	1.05	-16.67
13	Pusa Do Mausami	57.50	-6.35	16.00	-21.30	0.94	-25.40
14	Pusa Vishesh	60.00	-2.28	13.67	-32.76	0.82	-34.92
15	Pant Karela 1	58.00	-5.54	14.33	-29.51	0.80	-36.51
16	Pusa Hybrid 4	40.20	-34.53	22.67	11.51	0.91	-27.78
17	Pusa Hybrid 5	45.70	-25.57	13.67	-32.76	0.62	-50.79
18	Pusa Hybrid 6	61.40	0.00	20.33	0.00	1.26	0.00

Comment [R10]: Delete the column.

	(CheckControl)						
	<b>C.V</b>	<b>4.91</b>	-	<b>5.08</b>	-	<b>4.12</b>	-
	<b>SE(m)</b>	<b>1.83</b>	-	<b>0.57</b>	-	<b>0.03</b>	-
	<b>C.D @ 5%</b>	<b>5.26</b>	-	<b>1.64</b>	-	<b>0.08</b>	-



**Plate Figure 2: Phenotypic variability among bitter melon genotypes**

#### 4. CONCLUSION

Identifying genotypes that excel in yield attributing components is crucial for future breeding programs and enhancing bitter melon production. Genotypes like Konkan Tara, Pusa Rasdar, Priya, Phule Green Gold, and Pant Karela 4 showed superior traits such as larger fruit length, fruit diameter and higher yields, making them promising for cultivation. Conversely, Pusa Purvi, despite its smaller fruit size, produced the highest number of fruits per vine, indicating potential balance between fruit size and yield quantity. These findings contribute to the understanding of bitter melon genetics and offer insights for breeding programs aimed at developing varieties with improved agronomic traits suited to diverse agricultural conditions.

#### Disclaimer (Artificial intelligence)

Option 1: YES

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2: NO

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

## REFERENCES

1. ANONYMOUS, 2022, Indiastatagri, [www.indiastatagri.com/data/agriculture/bittergourd/data-year/all-years](http://www.indiastatagri.com/data/agriculture/bittergourd/data-year/all-years)
2. Jeffrey C. A review of the Cucurbitaceae. The Botanical Journal of the Linnean Society. 1980;81(3):233-247.
3. Okabe H, Miyahara Y, Yamauchi T. Studies on the constituents of *Momordica charantia* L. IV. Characterization of the new cucurbitacin glycosides of the immature fruits. (2) Structures of the bitter glycosides, momordicosides K and L. Chemical and Pharmaceutical Bulletin. 1982;30(12):4334-4340.
4. Yeh GY, Eisenberg DM, Kaptchuk TJ, Phillips RS. Systematic review of herbs and dietary supplements for glycemic control in diabetes. Diabetes Care. 2003;26(4):1277-1294.
5. Gopalan C, Rama Sastri BV, Balalsubramanian SC. Nutritive value of Indian foods. 2<sup>nd</sup> Ed. Hyderabad. National Institute of Nutrition, IGMR. 1993.
6. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. Statistical methods for agricultural workers. 1954: 347.
7. Yadav, M., Singh, D.B., Chaudhary, R. and Singh, D., 2008. Genetic variability in bitter gourd (*Momordica charantia* L.). *Journal of Horticultural Sciences*, 3(1), pp.35-38.
8. Rani, K.R., 2014. Performance of bitter gourd genotypes for yield and earliness. *Annals of Plant and Soil Research*, 16(4), pp.330-333.
9. Mallikarjunarao, K., Pradhan, R. and Hari Ram, K.B., 2020. Varietal Evaluation of Bitter Gourd (*Momordica charantia* L.) In Paralakhemundi, Gajapati District. *Indian Journal of Natural Sciences*, 10(60), pp.24152-24155.
10. Yadav, M., Chaudhary, R., Chandra, A. and Mehta, A. K. 2004. Genetic variability of different genotypes of bitter gourd (*Momordica charantia* L.). *The Allahabad Farmer*, 2: 70-76.
11. Ranjan, P., Gangopadhyay, K. K., Bag, M. K., Roy, A., Srivastava, R., Bhardwaj, R. and Dutta, M., 2015, Evaluation of cucumber (*Cucumis sativus* L.) germplasm for agronomic traits and disease resistance and estimation of genetic variability. *Indian Journal of Agricultural Sciences*, 85:234-239.
12. Ullah, M.Z., Hasan, M.J., Chowdhury, A.Z.M.K.A., Saki, A.I. and Rahman, A.H.M.A., 2012. Genetic variability and correlation in exotic cucumber (*Cucumis sativus* L.) varieties. *Bangladesh Journal of Plant Breeding and Genetics*, 25(1): 17-23.

**Comment [R11]:** Delete numbering in the reference list as numbering system is not used for citing references in the text.

**Formatted:** Font: Italic

**Formatted:** Font: Italic

**Formatted:** Font: Italic

**Formatted:** Font: Italic

13. Khan, M.H., Bhuiyan, S.R., Saha, K.C., Bhuyin, M.R. and Ali, A.S.M.Y., 2015. Variability, correlation and path co-efficient analysis of bitter gourd (*Momordica charantia* L.). Bangladesh Journal of Agricultural Research, 40(4): 607-618,
14. Ankitha , M. O., M. R. Bindu, S. Bhaskar Reddy, and S. Anand. 2023. "Divergence Analysis of Bitter Gourd Genotypes for Yield and Yield Attributes". International Journal of Plant & Soil Science 35 (13):95-101. <https://doi.org/10.9734/ijpss/2023/v35i132992>.
15. Bhati , Sachin, Satya Prakash, Vipin Kumar, Bijendra Singh, Manoj Kumar Singh, L. K. Gangwar, Deepak Kumar, Vishal Gangwar, and Avdhesh Kumar. 2023. "Genetic Variability, Heritability and Genetic Advance for Yield and Yield Contributing Characters in Bitter Gourd (*Momordica Charantia* L.) Genotypes". International Journal of Environment and Climate Change 13 (9):1867-75. <https://doi.org/10.9734/ijecc/2023/v13i92418>.
16. Mallikarjuna, K. N., Tomar, B. S., Mangal, M., Singh, N., Singh, D., Kumar, S., ... & Jat, G. S. (2024). Qualitative and quantitative genetic variations in bitter gourd (*Momordica charantia* L.). Genetic Resources and Crop Evolution, 1-19.
17. Hazra, P., Hazra, S., Acharya, B., Dutta, S., Saha, S., Mahapatra, P., ... & Ghosh, S. K. (2022). Diversity of nutrient and nutraceutical contents in the fruits and its relationship to morphological traits in bitter gourd (*Momordica charantia* L.). Scientia Horticulturae, 305, 111414.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic