

1
2
3
4
5
6
7
8
9
10

Varietal response to organic and inorganic fertilizers on bulb yield, storability, and production economics of onion (*Allium cepa* L.)

ABSTRACT

The field experiment was conducted in the Rabi season of 2021-22 in the research farm of the Regional Research Center, Raiya (Jhajjar) of Maharana Pratap Horticultural University, Karnal, Haryana to examine the impact of organic manures and inorganic fertilizers on the growth, bulb yield and self-life of two indigenous onion varieties was examined. The treatment combinations comprised seven treatments along with one control that was arranged in a randomized block design (factorial) with four replications and two varieties namely Hisar Onion-2 and Hisar Onion-4. The results of the study showed that bulb yield-related parameters such as average fresh weight of bulbs (82.39 g), yield per plot (26.74 kg), and yield per hectare (29.71 t) were recorded maximum in the treatment T6. However, the better storage life of bulbs was estimated in treatment control which was at par with treatment T3 i.e., physiological loss in weight (20.23%), sprouting (3.14%), and rotting (11.43%) were observed minimum after 120 days of storage period. Hisar onion-4 was found superior in all the bulb yield and shelf-life parameters in comparison to Hisar Onion-2. Treatment consisting of 50% recommended dose of NPK + 50% RDN supplied through Vermi-compost recorded maximum net return (Rs.1, 97,942) and benefit-cost ratio (2.25). Therefore, it can be inferred that the combined use of organic manures and inorganic fertilizers contributes to the improved shelf life of onion bulbs and enhances the long-term production potential of the soil.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

Keywords: Onion, organic, inorganic, manures, fertilizers, yield and variety)

1. INTRODUCTION

The onion (*Allium cepa* L.) belongs to the Amaryllidaceae family and is one of the most valuable commercial vegetable crops grown in India and around the world. It is a vital component in every household as a spice, vegetable, and salad, and hence fetches a high price in the domestic market. Onions are valued for their flavor and pungency due to the presence of the volatile sulfur compound allyl propyl disulfide. In India, the use of inorganic fertilizers, agrochemicals, and the extension of onion growing into non-traditional regions all contributed to an increase in onion production in the past (DOGR, 2013). India's average onion production per hectare is 16.41 t ha⁻¹, which is significantly less than the global average of 20.61 t ha⁻¹ (Anonymous, 2020-21). Instead of adding more agricultural inputs like inorganic fertilizers, productivity needs to be boosted with the help of the natural resources that are already available[16-19]. This can be done by cultivating improved

28 cultivars and improving nutrient usage efficiency (Tilman et al., 2002; Zhang et al., 2011)
29 because the widespread use of agrochemicals and synthetic fertilizers has led to various
30 undesirable environmental effects. The exclusive and continuous application of inorganic
31 fertilizers led to a decrease in the soil's organic matter content and diminished its
32 micronutrient content. The decline in soil organic matter adversely affected the soil's water-
33 holding capacity, soil structure, and water infiltration and resulting in increased soil
34 compaction. Simultaneously, the depletion of soil micronutrients resulted in deficiencies of
35 these essential elements (Dutta et al., 2003).

36 **According to Doran and Parkin, 1994, soil organic matter serves as an indicator of the**
37 **overall quality of the soil by influencing its biological, physical, and chemical aspects.**
38 Additionally, it serves as a reservoir for plant nutrients and serves as a substrate for soil
39 microbes (Dutta et al., 2003). The addition of organic manure is essential for revitalizing soil
40 health and sustaining prolonged onion production in tropical and subtropical climates.
41 However, while organic manure application alone enhances soil health, there is a decrease
42 in bulb yield in organic farming systems, ranging from 22% to 45% compared to inorganic
43 systems **in the early years, according to these authors** (Lawande et al., 2009). Practices
44 for managing nutrients that involve a combination of chemical fertilizers organic manures
45 and bio-fertilizers thus become critical for enhancing bulb output and maintaining the vitality
46 of the soil. The current investigation indicates that the farmers are applying excess chemical
47 fertilizers as per crop requirements. The integrated nutrient management approach also
48 improved both the quality and post-harvest storage life of the onion crop. Considering this,
49 we conducted a field experiment to explore the impact of organic manures and inorganic
50 fertilizers on both bulb yield and the storage life of onions.

51

52 **2. MATERIAL AND METHODS**

53

54 The research work was carried out during the Rabi season 2021-22 in the research farm of
55 the Regional Research Center, Raiya (Jhajjar) of Maharana Pratap Horticultural University,
56 Karnal, Haryana. **It is situated about 8 km towards South from District Jhajjar on**
57 **Jhajjar – Kosali road with an elevation of 222 m above the mean sea level.** It
58 experiences a typical semi-arid climate characterized by hot and dry summers and extremely
59 cold winters, with an annual rainfall of 456 mm. The experimental site's topography was
60 uniform, the field was leveled flat, and the soil was sandy loam with a medium water
61 retention capacity. Experimental material consists of eight treatments of different nutrient
62 status and two different varieties Hisar Onion-2 and Hisar Onion-4. The experiment was
63 arranged in a factorial randomized block design with four replications including one absolute
64 control and one recommended dose of fertilizers (RDF) (125:50:25 kg NPK/ha). At the time
65 of transplanting, half of the required nitrogen, as well as the entire quantity of the
66 phosphorus and potash were applied as basal dose. The remaining nitrogen dose was
67 administered in two equal split doses, 30 and 45 days after transplanting, while organic
68 manures, including farmyard manure, vermicompost, and poultry manure, were thoroughly
69 mixed in the top 15 cm of soil as per treatment during field preparation. The various
70 treatment combinations along with their symbols are presented in Table 1. Eight-week-old
71 seedlings were dipped into the solution of phosphorous-solubilizing bacteria for 30 minutes
72 as per treatment except for control and then transplanted in flat beds of 3 m x 3 m size at a
73 spacing of 15 cm x 10 cm.

74 The observations were collected on the bulb yield characteristics for randomly selected five
75 plants per treatment per plot in each replication and the observations recorded for yield
76 parameters were the average weight of fresh bulb (g), number of bulbs per kg, yield per plot
77 (kg), yield per hectare (tonnes). For a collection of data on self- life of onions, three
78 kilograms of onion bulbs from each treatment and replication were stored in ventilated

79 wooden cages at room temperature and observations were recorded at 40, 80, and 120
 80 days after storage for sprouting, rotting, and physiological loss in weight (%). To calculate
 81 the cost of bulb cultivation, all operations including nursery and land preparation, soil
 82 reclamation, planting materials, intercultural operation, irrigation, plant protection, harvesting,
 83 and cost of various manures and fertilizers were taken into consideration. The gross and net
 84 incomes were calculated along with the benefit-cost ratio. The data about various
 85 parameters were subjected to statistical analysis using the Analysis of Variance (ANOVA)
 86 technique as outlined by Panse and Sukhatme (1987).

87
 88
 89
 90


Table: 1 Treatments detail various organic and inorganic fertilizers and their combination

Treatment symbols	Treatment details
T1	Recommended dose of fertilizer (125:50:25 kg/ha)
T2	Recommended dose of nitrogen supplied through farmyard manure
T3	Recommended dose of N supplied through vermicompost
T4	Recommended dose of N supplied through poultry manure
T5	50% Recommended dose of NPK + 50% Recommended dose of N supplied through Farm Yard Manure
T6	50% Recommended dose of NPK + 50% Recommended dose of N supplied through vermicompost
T7	50% Recommended dose of NPK + 50% Recommended dose of N supplied through poultry manure
T8	Control (Without any fertilizers)

91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108

3. RESULTS AND DISCUSSION

The maximum average fresh weight of the bulb (82.39 g) was recorded when the nutrient was applied with 50% RDF through NPK + 50% RDN through VC, which was observed statistically at par with the application of 50% RDF through NPK + 50% RDN through PM (80.38 g) and RDF (79.98 g). Whereas the minimum average fresh weight of the bulb (54.63 g) was reported in the control treatment. Within the onion varieties, Hisar Onion-4 (26.68 t) produced significantly more bulb yield per plot and yield per hectare than Hisar Onion-2 (23.08 kg and 25.46 t. The increase in bulb yield contributing characters as mentioned in Figure 1. is because organic manures are advantageous to plants because they are a source of many crucial macro and micronutrients. They are rich in organic matter and boost the availability and uptake of nitrogen, phosphorous and potassium, which stimulates photosynthesis and metabolic activities and has a beneficial impact on plant cell division and elongation, which leads to an increase in size and fresh weight of bulb ultimately results in an overall improvement in the yield of onion bulbs. Verma et al. (2016), Singh et al. (2016), and Dhaker et al. (2018) also have similar findings.



109
110
111

112 Hisar Onion-4 (73.60 g) had produced significantly higher fresh weight of bulb than Hisar
113 Onion-2 (71.78 g). Lesser number of bulbs per kg (12.15) was recorded in treatment T6
114 (50% RDF through NPK + 50% RDN through VC) followed by treatment having 50% RDF
115 through NPK + 50% RDN through PM (12.46) and RDF (12.52), whereas, the maximum
116 number of bulbs per kg (18.37) was observed in T8 (control). In case of onion varieties the
117 minimum number of bulbs per kg was noted in Hisar Onion-4 (13.82) in comparison to Hisar
118 Onion-2 (14.19). Application of 50% RDF through NPK + 50% RDN through VC produced
119 significantly higher bulb yield per plot (26.74 kg) and yield per hectare (29.71 t), which was
120 observed statistically at par with the application of 50% RDF through NPK + 50% RDN
121 through PM (26.20 kg and 29.11 t) and RDF (25.57 kg and 28.42 t). The control treatment
122 produced the minimum bulb yield per plot and yield per hectare (16.95 kg and 17.23 t).
123 Among the onion varieties, Hisar Onion-4 (24.18 kg and 26.68 t) produced significantly more
124 yield per plot and yield per hectare than Hisar Onion-2 (23.08 kg and 25.46 t).

125 As regards the storage studies of 40 days after storage, the rotting percentage was
126 observed least (0.34%) in control (T8), whereas the maximum rotting percentage (0.75%)
127 was observed in treatment consisting of 100% RDF (T1). However, after 80 and 120 days of
128 storage, it was observed minimum (2.45%) and (11.18%) in control (T8) which was
129 statistically at par with treatment T3 – 100% RDN through VC (2.63%) and (11.43%)
130 respectively, whereas maximum rotting percentage at 80 days after storage (3.72%) and 120
131 days after storage (16.03%) was observed in treatment consisting 100% RDF (T1) as
132 depicted in table 2. This could be attributed to the elevated nitrogen application levels, which
133 may prompt the plants to produce larger bulbs with soft and succulent tissues. This condition
134 renders them more susceptible to bacterial infections, leading to the development of bulbs
135 with thick necks that hinder proper drying and consequently result in rotting. Diaz-Perez et
136 al. (2003) concluded that excessive application of nitrogen fertilizer should be avoided
137 although it minimizes the risk of bolting it tends to elevate bulb decay.

138 **Table: 2 Effect of organic and inorganic fertilizers on rotting losses (%) of different**
139 **onion varieties during storage**

	Rotting losses (%) at 40 DAS	Rotting losses (%) at 80 DAS	Rotting losses (%) at 120 DAS

Treatments	Varieties		Mean	Varieties		Mean	Varieties		Mean
	HO-2	HO-4		HO-2	HO-4		HO-2	HO-4	
T ₁	0.79	0.71	0.75	3.89	3.55	3.72	16.42	15.64	16.03
T ₂	0.62	0.53	0.57	2.97	2.72	2.84	12.61	11.63	12.12
T ₃	0.53	0.48	0.5	2.69	2.56	2.63	11.58	11.22	11.40
T ₄	0.59	0.52	0.56	2.81	2.64	2.72	12.24	11.56	11.90
T ₅	0.63	0.56	0.59	3.15	2.96	3.05	14.24	13.44	13.84
T ₆	0.64	0.61	0.63	3.26	3.05	3.15	13.67	13.12	13.40
T ₇	0.66	0.64	0.65	3.33	3.1	3.21	14.14	13.46	13.80
T ₈	0.3	0.39	0.34	2.51	2.39	2.45	11.27	11.08	11.18
Mean	0.59	0.55		3.08	2.87		13.28	12.64	
Factors	Factor (T)	Factor (V)	Factor (T X V)	Factor (T)	Factor (V)	Factor (T X V)	Factor (T)	Factor (V)	Factor (T X V)
C.D. at 5%	0.06	0.03	NS	0.22	0.11	NS	0.24	0.12	NS

140
141
142
143

T₁: RDF, T₂: RDN through FYM, T₃: RDN through VC, T₄: RDN through PM, T₅: 50% RDF through NPK + 50% RDN through FYM, T₆: 50% RDF through NPK + 50% RDN through VC, T₇: 50% RDF through NPK + 50% RDN through PM, T₈: Control

144
145
146
147
148
149
150
151
152

The data is presented in Table. 3 indicates that the sprouting percentage was noted as negligible up to a period of 40 days after storage. After 80 and 120 days of storage, the sprouting percentage varied from 1.10% to 5.06%. Same as rotting (%), significantly, least sprouting was observed in treatment T₈ (control) at 80 (1.10%) and 120 (3.08%) days of storage, which was statistically at par with T₃ (1.18%) and (3.14%). The maximum sprouting (2.07% & 5.06%) was observed in treatment T₁ (100% RDF) after 80 and 120 days of storage. The current study related to the findings of Eze and Orkwor (2010) who observed that higher sprouting was exhibited when inorganic fertilizers and organic manure were applied compared to the control.

153
154
155
156
157
158
159
160
161
162
163
164
165
166
167

At 40 days after storage, the minimum physiological loss in weight was found in the treatment control (T₈), i.e., (4.89%) which was significantly at par with the application of 100% RDN through VC, i.e., (5.03%), whereas, the maximum physiological loss in weight (8%) was observed in treatment T₁ (100% RDF) as recorded in table 4. The minimum physiological loss in weight was recorded in the treatment where no nutrient source was applied, i.e., control (9%) which was significantly lesser and at par with the application of 100% RDN through VC, i.e., T₃ (9.36%), whereas, the maximum physiological loss in weight (12.91% & 26.78%) was found in the treatment (T₁) having 100% RDF application after 80 and 120 days of storage period. The beneficial effect of organic manure in reducing the post-harvest losses of horticultural crops has been reported by Sankar et al. (2009) and Ahmad (2016).

Table: 3 Effect of organic and inorganic fertilizers on sprouting losses (%) of different onion varieties during storage

	Sprouting losses (%) at 40 DAS	Sprouting losses (%) at 80 DAS	Sprouting losses (%) at 120 DAS
--	--------------------------------	--------------------------------	---------------------------------

Treatments	Varieties		Mean	Varieties		Mean	Varieties		Mean
	HO-2	HO-4		HO-2	HO-4		HO-2	HO-4	
T ₁	0	0	0	2.19	1.94	2.07	5.12	5.01	5.06
T ₂	0	0	0	1.46	1.34	1.4	3.54	3.23	3.38
T ₃	0	0	0	1.21	1.15	1.18	3.11	3.16	3.14
T ₄	0	0	0	1.29	1.2	1.24	3.33	3.19	3.26
T ₅	0	0	0	1.64	1.45	1.55	4.23	4.17	4.20
T ₆	0	0	0	1.74	1.51	1.63	4.16	3.93	4.05
T ₇	0	0	0	1.72	1.61	1.66	4.04	4.12	4.08
T ₈	0	0	0	1.13	1.07	1.1	3.09	3.07	3.08
Mean	0	0		1.55	1.41		3.83	3.73	
Factors	Factor (T)	Factor (V)	Factor (T X V)	Factor (T)	Factor (V)	Factor (T X V)	Factor (T)	Factor (V)	Factor (T X V)
C.D. at 5%	0	0	NS	0.12	0.06	NS	0.16	0.08	NS

168 T1: RDF, T2: RDN through FYM, T3: RDN through VC, T4: RDN through PM, T5: 50% RDF through NPK +
169 50% RDN through FYM, T6: 50% RDF through NPK + 50% RDN through VC, T7: 50% RDF through NPK +
170 50% RDN through PM, T8: Control

171 **Table: 4 Effect of organic and inorganic fertilizers on physiological losses of weight**
172 **(%) of different onion varieties during storage**

Treatments	Physiological losses of weight(%) at 40 DAS			Physiological losses of weight(%) at 80 DAS			physiological losses of weight (%)at 120 DAS		
	Varieties		Mean	Varieties		Mean	Varieties		Mean
	HO-2	HO-4		HO-2	HO-4		HO-2	HO-4	
T ₁	8.32	7.68	8.00	13.12	12.71	12.91	27.13	26.42	26.78
T ₂	5.37	5.17	5.27	9.65	9.49	9.57	22.34	21.44	21.89
T ₃	5.12	4.95	5.03	9.44	9.29	9.36	20.73	19.53	20.13
T ₄	5.18	5.02	5.10	9.55	9.41	9.48	21.56	20.55	21.06
T ₅	6.53	6.30	6.42	11.23	11.08	11.15	26.24	25.29	25.76
T ₆	6.68	6.43	6.56	11.37	11.28	11.32	24.68	23.63	24.16
T ₇	6.73	6.49	6.61	11.45	11.32	11.38	25.28	24.31	24.80
T ₈	5.03	4.74	4.89	9.07	8.94	9.00	19.98	19.23	19.61
Mean	6.12	5.85		10.61	10.44		23.51	22.55	
Factors	Factor (T)	Factor (V)	Factor (T X V)	Factor (T)	Factor (V)	Factor (T X V)	Factor (T)	Factor(V)	Factor (T X V)
C.D. at 5%	0.20	0.10	NS	0.37	0.14	NS	0.53	0.26	NS

173 T1: RDF, T2: RDN through FYM, T3: RDN through VC, T4: RDN through PM, T5: 50% RDF through NPK + 50%
 174 RDN through FYM, T6: 50% RDF through NPK + 50% RDN through VC, T7: 50% RDF through NPK + 50% RDN
 175 through PM, T8: Control

176
 177 Table 5. illustrates the economic data associated with various treatments. It is evident from
 178 the data obtained that a significant maximum bulb yield of 29.71 t/ha was obtained under the
 179 treatment T6 (50% RDF through NPK + 50% RDN through VC) with a maximum net return
 180 of Rs.197942.5/ha and benefit-cost ratio of 2.25. However, the minimum bulb yield of 17.23
 181 t/ha, with a minimum net return of Rs.72260/ha and a benefit-cost ratio of 1.54 was noted in
 182 treatment T8 (Control).

183
 184 **Table: 5 Effect of organic and inorganic fertilizers on the economics of onion**
 185 **production (Rs/ha)**

Treatments	Cost of cultivation	Gross income	Net income	BCR
T1	172655	341040	168385	1.98
T2	170500	298440	127940	1.75
T3	174500	311520	137020	1.79
T4	170500	304320	133820	1.78
T5	156577.5	335280	178702.5	2.14
T6	158577.5	356520	197942.5	2.25
T7	156577.5	349320	192742.5	2.23
T8	134500	206760	72260	1.54

186 T1: RDF, T2: RDN through FYM, T3: RDN through VC, T4: RDN through PM, T5: 50% RDF through NPK + 50%
 187 RDN through FYM, T6: 50% RDF through NPK + 50% RDN through VC, T7: 50% RDF through NPK + 50% RDN
 188 through PM, T8: Control

189 **4. CONCLUSION**

190
 191 Based on the one-year study, it can be concluded that the maximum bulb yield (29.71 t) and
 192 cost-benefit ratio (2.25) were observed when the onion varieties were applied with 50% of
 193 the recommended dose of NPK + 50% of the recommended dose of nitrogen supplied
 194 through vermicompost. The storage life of onion was recorded higher in treatment, whereas
 195 100% of the recommended dose of nitrogen was supplied through vermicompost. **The**
 196 **treatment T2, T3 and T4 are the most agroecological treatments that do not depend on**
 197 **external fertilizers and pollute less soil and water.**

198 199 **ACKNOWLEDGEMENTS**

200
 201 I would like to express my sincere gratitude to Dr. Satya Pal Singh (Assistant Scientist,
 202 Vegetable Science, MHU, Karnal) for their invaluable guidance, unwavering support, and
 203 insightful feedback throughout the course of this research. Their expertise and
 204 encouragement have been instrumental in shaping the direction of this study. I am also

205 thankful to Dr. D.S. Duhan, Assistant scientist, Department of Vegetable Science, HAU,
206 Hisar for their valuable input and constructive guidance, which significantly contributed to the
207 refinement of this research and special thanks go to Sidharth, Ankit Saini, Renu Fundan,
208 Nilesh Sagwal for their collaboration, insightful discussions, and moral support. Their diverse
209 perspectives have enriched the depth and quality of this research.

210
211 Disclaimer (Artificial intelligence)

212 Option 1:

213 Author(s) hereby declare that NO generative AI technologies such as Large Language
214 Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during
215 writing or editing of manuscripts.

216 Option 2:

217 Author(s) hereby declare that generative AI technologies such as Large Language Models,
218 etc have been used during writing or editing of manuscripts. This explanation will include
219 the name, version, model, and source of the generative AI technology and as well as all
220 input prompts provided to the generative AI technology

221 Details of the AI usage are given below:

222 1.

223 2.

224 3.

225

226

227 REFERENCES

228

229 1. Ahmad, H.A.F. Impact of different fertilizer sources on vegetative growth, yield, quality and
230 storability of onion. *International Invention Journal of Agricultural and Soil Science*. 2016:
231 4(1): 1-8.

232 2. Anonymous. Area and production of horticulture Crops: Food and Agriculture
233 Organization Statistics. 2020-21, <http://faostat.fao.org/default.aspx>

234 3. Dhaker, B., Sharma, R.K., Chhipa, B.G. and Rathore, R.S. Effect of different organic
235 manures on yield and quality of onion (*Allium cepa* L.). *International Journal of Current*
236 *Microbiology Applied Science*. 2018: 6(11): 3412-3417.

237 4. Diaz-Perez, J.C., Purvis, A.C. and Paulk, J.T. Bolting, yield and bulb decay of sweet onion
238 as affected by nitrogen fertilization. *Journal of the American Society for Horticultural Science*.
239 2003: 128(1): 144-149.

240 5. DOGR. Directorate of Onion and Garlic Research Vision 2050, *Directorate of Onion and*
241 *Garlic Research, Rajgurunagar, Pune*. (2013).

242 6. Doran, J. W., & Parkin, T. B. Defining and assessing soil quality. *Defining soil quality for a*
243 *sustainable environment*. 1994: 35, 1-21.

- 244 7. Dutta, S., Pal, R., Chakerabarty, A. and Chakrabarti, K. Influence of integrated plant
245 nutrient supply system on soil quality restoration in a red and laterite soil. *Archives of*
246 *Agronomy and Soil Science*. 2003: 49: 631-37.
- 247 8. Eze S.C. and Orkwor, G.C. Studies on effect of mineral fertilizer, organic manure and
248 cultivar on the yield and storability of yam (*Dioscorea rotundata* Poir). *African Journal of*
249 *Food Agricultural, Nutritional and Development*. 2010: 10(6): 2755-2771.
- 250 9. Lawande, K.E., Anil Khar, Mahajan, V., Srinivas, P.S., Sankar, V. and Singh, R.P. Onion
251 and garlic research in India. *Journal of Horticultural Sciences*. 2009: 4: 91-119.
- 252 10. Panse V G and Sukhatme P V. Statistical Methods for Agricultural Workers. Indian
253 Council of Agricultural Research, New Delhi. (1985).
- 254 11. Sankar, V., Veeraragavathatham, D. and Kannan, M. Studies on organic farming in
255 onion (*Allium cepa* L.) for the production of export quality bulbs. *Asian Journal of*
256 *Horticulture*. 2009: 4(1): 65-69.
- 257 12. Singh, S.K., Mahto, D.K., Mane, P.D., Chaudhary, S.K., Kumar, M. and Yadav, L.M.
258 Effect of INM in onion with respect to growth and yield under North Bihar condition. *An*
259 *International Journal Print*. 2016: 11: 6515-6518.
- 260 13. Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R. and Polasky, S. Agricultural
261 sustainability and intensive production practices. *Nature*. 2002: 418: 671-77.
- 262 14. Verma, A.B. and Singh, H.M. Effect of organic manures and inorganic fertilizers on
263 growth and yield of onion (*Allium cepa* L). *Hortflora Research Spectrum*. 2016: 2(5): 173-
264 174.
- 265 15. Zhang, F.S., Cui, Z.L., Fan, M.S., Zhang, W.F., Chen, X.P. and Jiang, R.F. Integrated
266 soil-crop system management: reducing environmental risk while increasing crop
267 productivity and improving nutrient use efficiency in China. *Journal of Environmental Quality*.
268 2011: 40: 1051-57.
- 269
- 270 16. Raa, Vikas, Satya Pal Singh, D.S. Duhan, V.S. Mor, Sidharth, and Ankit Saini. 2024.
271 "Varietal Response to Organic and Inorganic Fertilizers on Growth, Bulb Yield and Quality of
272 Onion (*Allium Cepa* L.)". *International Journal of Plant & Soil Science* 36 (5):755-62.
273 <https://doi.org/10.9734/ijpss/2024/v36i54573>.
- 274
- 275
- 276 17. Hussain S, Kerketta A. Effect of Organic Manure and Inorganic Fertilizer on Growth and
277 Root Yield of Beetroot (*Beta vulgaris* L.). *Int. J. Environ. Clim. Change*. [Internet]. 2023 Jun.
278 17 [cited 2024 Jun. 11];13(8):1866-70. Available from:
279 <https://journalijecc.com/index.php/IJECC/article/view/2141>
- 280
- 281
- 282 18. Heeb A, Lundegårdh B, Savage G, Ericsson T. Impact of organic and inorganic fertilizers
283 on yield, taste, and nutritional quality of tomatoes. *Journal of plant nutrition and soil science*.
284 2006 Aug;169(4):535-41.
- 285
- 286

287 19.Moe K, Moh SM, Htwe AZ, Kajihara Y, Yamakawa T. Effects of integrated organic and
288 inorganic fertilizers on yield and growth parameters of rice varieties. *Rice Science*. 2019 Sep
289 1;26(5):309-18.
290