

Exploring the Floral Biology of *Salix alba*: Insights into Reproductive Dynamics, Pollen Production and Pollen Morphology in Kashmir Himalayas

Abstract

This study investigates the floral phenology, pollen production and pollen characteristics of *Salix alba* (white willow) over the years 2022 and 2023, highlighting inter-annual and individual tree variations. In 2022, male catkins began flowering between March 8 and March 10, while female catkins initiated between March 14 and March 22. Flowering for male catkins lasted until mid-May and for female catkins until late May. In 2023, male catkins started flowering from March 10 to March 14 and female catkins from March 17 to March 24, with similar flowering durations. Inflorescence diameters varied among individual trees, ranging from 4.4 mm to 7.7 mm for male catkins and 6.1 mm to 8.1 mm for female catkins across the two years. Pollen production exhibited significant variability with the highest recorded at 4,600,995,000 grains per tree and the lowest at 498,312,500 grains per tree. On average, trees produced approximately 2,164,836,667 pollen grains. Pollen viability ranged from 75% to 80%, with sizes between 16 μm and 26 μm . The pollen-ovule ratio also showed substantial differences, with the highest ratio observed at 7600:2 and the lowest at 2345:7. These findings underscore the considerable year-to-year and tree-to-tree variation in the reproductive traits of *Salix alba*, which may have implications for its breeding and conservation strategies.

Introduction

In the rich tapestry of our natural world, *Salix alba*, commonly known as White Willow, emerges as a captivating subject of scientific inquiry within the genus *Salix* and family Salicaceae (Tawfeek et al., 2021). Originating from the Celtic words 'sal' meaning near and 'lis' signifying water, the genus *Salix* aptly reflects its preference for aquatic habitats, thriving along riverbanks and lakeshores (Raj and Lal, 2014; Oberprieler et al., 2013).

Salix alba, among the 450 species in the genus *Salix*, holds significant ecological and economic importance globally, ranging from tropical to cold-temperate regions (Argus, 1997; Skvortsov, 1999; Helfenstein et al., 2014). Widely distributed from Europe to Central Asia and North Africa, this species was introduced to the America in the 17th century, becoming naturalized across continents (Dickmann and Kuzovkina, 2014; Kuzovkina and Quigley, 2005; Trybush et al., 2008).

Historically, *Salix* cultivation traces back to ancient times, utilized by civilizations like the Romans for crafting baskets, furniture and medicinal remedies (Masoodi et al., 2008). Modern cultivation and breeding programs have further diversified its applications, ranging from ornamental landscaping to biomass production for renewable energy (Pei and McCracken, 2005).

In India, *Salix alba* finds a habitat in the Western Himalayas, thriving at elevations up to 2400 meters along streams and canal banks in Kashmir and Kulu valleys (Luna, 1995). Introduced to Kashmir in 1927, this species has since been cultivated for various purposes, including soil conservation, fuel wood and fodder (Anonymous, 1983).

Salix alba exhibits remarkable diversity, encompassing varieties such as var. *alba*, var. *vitellina* (Golden Willow) and var. *caerulea* (Cricket-bat Willow), each serving distinct economic roles from ornamental use to specialized timber production (Gupta et al., 2014). Its timber, valued for its lightness, strength and shock resistance, is particularly sought after for applications like cricket bats and furniture (Stott, 1984).

The reproductive biology of *Salix alba* presents a compelling area of study, characterized by dioecious traits where male and female individuals bear separate catkins. These unisexual flowers play a pivotal role in its reproductive strategy, facilitated primarily by insect pollination, although wind pollination (ambophily) also contributes to its genetic diversity (Saunders, 2020).

Floral biology studies on *Salix alba* encompass the phenology of its flowering events, pollen production, pollen viability and the morphological characteristics of its reproductive structures. These aspects are crucial for understanding its reproductive success, genetic variation, and adaptation strategies in changing environments (Choudhary et al., 2011).

By delving into the intricate details of its reproductive and floral biology, this research aims to unveil the mechanisms underlying *Salix alba's* ecological resilience and economic potential. Such insights not only enrich our understanding of this versatile species but also inform sustainable management practices and conservation efforts amid contemporary environmental challenges.

Materials and Methods

Study Area

This study was conducted in the Ganderbal district of Jammu and Kashmir, India. Ganderbal is situated in the central part of the Kashmir Valley and is known for its diverse flora, which includes a variety of tree species such as *Salix alba*. The region experiences a temperate climate with distinct seasonal variations, including cold winters and warm summers, which significantly influence the phenological events of plant species. The study site within Ganderbal were chosen to represent a range of microclimatic conditions to capture the variability in reproductive parameters of *Salix alba*. The specific GPS coordinates of the study area are approximately 34.2262° N latitude and 74.7787° E longitude.

The reproductive parameters of *Salix alba* were studied randomly across the Ganderbal district. Six young, sexually mature trees were selected, with four major branches marked in each of the cardinal directions (North, South, East and West) using metal tags for consistent observation.

Floral phenology was recorded by observing the initiation, duration and termination of flowering, type of inflorescence and inflorescence diameter using a digital Vernier caliper. These observations were made visually on permanently marked trees, noting the timing of each parameter during both vegetative and reproductive phases.

Pollen production was assessed by determining the number of pollen grains per anther using a microscope, as per Dafni (1992) and Kearns and Inouye (1993). Pollen production per flower was calculated by multiplying the number of pollen grains per anther by the number of anthers per flower. This was extended to per inflorescence by multiplying the average number of flowers per inflorescence with

the average number of pollen grains per flower. Total pollen production per tree was estimated using the formula: $\Sigma TP = \Sigma N \times F \times A \times P$, where ΣTP represents total pollen grains per tree, ΣN is the number of inflorescences per tree, F is the average number of flowers per inflorescence, A is the average number of anthers per flower, and P is the average number of pollen grains per anther (Khanduri, 2012).

Pollen-ovule ratios were determined by dividing the number of pollen grains per flower by the number of ovules per flower, following Cruden (1977). Pollen viability was estimated in vitro using Acetocarmine staining methods (Alexander, 1969) and pollen size was measured with a stage micrometer as described by Ascari et al. (2020).

Results

The study observed significant variations in the floral phenology, pollen production and pollen characteristics of *Salix alba* across different years and individual trees.

Floral Phenology:

For the year 2022, it was observed that the flowering initiation for male catkins ranged from 8th March to 10th March and for female catkins from 14th March to 22nd March. The flowering duration for male catkins was from 8th March to 14th May and for female catkins from 14th March to 25th May. Termination of flowering occurred between 10th May and 14th May for male catkins and between 20th May and 25th May for female catkins. Inflorescence diameter varied from 4.4 mm to 7.7 mm for male catkins and from 6.1 mm to 8.1 mm for female catkins.

For the year 2023, it was observed that the flowering initiation for male catkins ranged from 10th March to 14th March and for female catkins from 17th March to 24th March. The duration of flowering for male catkins was from 10th March to 12th May and for female catkins from 17th March to 26th May. Termination of flowering occurred between 9th May and 12th May for male catkins and between 22nd May and 26th May for female catkins. Inflorescence diameter ranged from 4.4 mm to 7.2 mm for male catkins and from 6.1 mm to 7.6 mm for female catkins.

Pollen Production:

The highest pollen production was observed in T5 with 4,600,995,000 pollen grains, while the lowest was in T6 with 498,312,500 pollen grains. The average pollen production was 2,164,836,667 pollen grains per tree, indicating substantial variability.

Pollen Viability, Size, and Pollen-Ovule Ratio:

- Pollen viability ranged from 75% (T5) to 80% (T2). Pollen size varied from 16 μm (T1) to 26 μm (T6). The pollen-ovule ratio also showed significant variation, with the highest pollen count per ovule observed in T1 (7600:2) and the lowest in T6 (2345

Table 1: Floral Phenology of *Salix alba* (2022)

| Tree No. | Flowering Initiation (Male Catkins) | Flowering Initiation (Female Catkins) | Duration of Flowering (Male Catkins) | Duration of Flowering (Female Catkins) | Termination of Flowering (Male Catkins) | Termination of Flowering (Female Catkins) | Type of Inflorescence | Inflorescence Diameter (Male Catkins) | Inflorescence Diameter (Female Catkins) |
|----------|-------------------------------------|---------------------------------------|--------------------------------------|--|---|---|-----------------------|---------------------------------------|---|
| T1 | 8th March | 18th March | 8th March to 10th May | 18th March to 25th May | 10th May | 25th May | Catkin or Ament | 5.8 | 6.4 |
| T2 | 9th March | 15th March | 9th March to 11th May | 15th March to 24th May | 12th May | 24th May | Catkin or Ament | 7.7 | 6.1 |
| T3 | 8th March | 20th March | 8th March to 12th May | 20th March to 22nd May | 11th May | 22nd May | Catkin or Ament | 4.4 | 8.1 |
| T4 | 9th March | 19th March | 9th March to 13th May | 19th March to 23rd May | 13th May | 23rd May | Catkin or Ament | 5.1 | 7.3 |
| T5 | 8th March | 14th March | 8th March to 14th May | 14th March to 20th May | 14th May | 20th May | Catkin or Ament | 5.8 | 7.5 |
| T6 | 10th March | 22nd March | 10th March to 11th May | 22nd March to 22nd May | 11th May | 22nd May | Catkin or Ament | 6.2 | 7.9 |

Table 2: Floral Phenology of *Salix alba* (2023)

| Tree No. | Flowering Initiation (Male Catkins) | Flowering Initiation (Female Catkins) | Duration of Flowering (Male Catkins) | Duration of Flowering (Female Catkins) | Termination of Flowering (Male Catkins) | Termination of Flowering (Female Catkins) | Type of Inflorescence | Inflorescence Diameter (Male Catkins) | Inflorescence Diameter (Female Catkins) |
|----------|-------------------------------------|---------------------------------------|--------------------------------------|--|---|---|-----------------------|---------------------------------------|---|
| T1 | 10th March | 19th March | 10th March to 9th May | 19th March to 26th May | 9th May | 26th May | Catkin or Ament | 5.6 | 6.3 |
| T2 | 11th March | 17th March | 11th March to 10th May | 17th March to 25th May | 10th May | 25th May | Catkin or Ament | 7.2 | 6.1 |
| T3 | 12th March | 23rd March | 12th March to 10th May | 23rd March to 25th May | 10th May | 25th May | Catkin or Ament | 4.4 | 7.6 |
| T4 | 13th March | 21st March | 13th March to 12th May | 21st March to 23rd May | 12th May | 23rd May | Catkin or Ament | 5.0 | 7.2 |
| T5 | 12th March | 18th March | 12th March to 12th May | 18th March to 22nd May | 12th May | 22nd May | Catkin or Ament | 5.5 | 7.1 |
| T6 | 14th March | 24th March | 14th March to 9th May | 24th March to 24th May | 9th May | 24th May | Catkin or Ament | 6.1 | 7.5 |

Table 3: Pollen Production in *Salix alba*

| Tree No. | No. of Anthers per Catkin | No. of Pollens per Anther | No. of Pollens per Catkin | No. of Catkins per Tree | No. of Pollens per Tree |
|----------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| T1 | 96 | 7600 | 729600 | 2000 | 1,459,200,000 |

| | | | | | |
|----------------|--------------|-------------|------------------|-------------|----------------------|
| T2 | 72 | 5050 | 363600 | 5000 | 1,818,000,000 |
| T3 | 88 | 6600 | 580800 | 6000 | 3,484,800,000 |
| T4 | 78 | 7090 | 553020 | 3000 | 1,659,060,000 |
| T5 | 87 | 7555 | 657285 | 7000 | 4,600,995,000 |
| T6 | 85 | 2345 | 199325 | 2500 | 498,312,500 |
| Average | 84.33 | 6040 | 509373.33 | 4250 | 2,164,836,667 |

Table 4: Pollen Viability, Size and Pollen-Ovule Ratio in *Salix alba*

| Parameter→ Tree No. ↓ | Pollen viability (percent) | Pollen size (micrometer) | Pollen Ovule ratio | |
|--------------------------|-------------------------------|-----------------------------|--------------------|-------|
| | | | Pollen | Ovule |
| T ₁ | 76 | 16 | 7600 | 2 |
| T ₂ | 80 | 21 | 5050 | 3 |
| T ₃ | 79 | 23 | 6600 | 5 |
| T ₄ | 79 | 22 | 7090 | 8 |
| T ₅ | 75 | 25 | 7555 | 6 |
| T ₆ | 78 | 26 | 2345 | 7 |

Discussion

The study on the reproductive parameters of *Salix alba* in the Ganderbal district revealed significant variations in floral phenology, pollen production and pollen characteristics. These findings provide valuable insights into the reproductive strategies and potential environmental adaptations of *Salix alba*.

The observed variability in flowering initiation, duration and termination corresponds with previous research indicating the strong influence of local climatic conditions on floral phenology (Ghelardini et al., 2010). The earlier onset and prolonged flowering in 2022 compared to 2023 likely reflect annual climatic fluctuations, which can affect pollination dynamics and seed production (Vitasse et al., 2009).

Pollen production among *Salix alba* trees exhibited substantial variability, consistent with findings in other plant species (Khanduri, 2012). This diversity in pollen production may stem from genetic differences, tree health and varying environmental factors. Trees with higher pollen production, such as T5, suggest a potential advantage in reproductive success, crucial for maintaining population viability across different environmental conditions.

The wide range of pollen-ovule ratios observed in this study aligns with Cruden's theory (1977) that this ratio serves as an indicator of a plant's breeding system. Higher ratios in certain trees indicate an adaptation to optimize reproductive success under conditions of limited pollinator availability (Cruden & Miller-Ward, 1981). This finding is consistent with recent studies on other plant species (Wagay et al., 2024).

High pollen viability rates (75%-80%) observed in *Salix alba* indicate strong potential for successful fertilization, supported by reliable staining techniques (Alexander, 1969). Variations in pollen size, influenced by environmental factors and genetic diversity, may affect pollen dispersal and germination success (Ascari et al., 2020). *Salix alba's* ambophilous pollination strategy, involving both insect and wind pollination mechanisms, underscores its reproductive flexibility and adaptability to diverse environmental conditions, as noted by Saunders (2020).

The findings of present study are also in conformity with the findings of Rafeeq et al., 2022, Mir et al., 2024 and Wagay et al., 2024. The findings from this study contribute to a deeper

understanding of *Salix alba*'s reproductive ecology and adaptation strategies. The variability in floral phenology, pollen production, and pollen characteristics highlights the species' resilience and potential responses to changing environmental conditions. These insights are crucial for conservation efforts and sustainable management practices aimed at preserving *Salix alba* populations in varying ecosystems.

Conclusion

This study provides valuable insights into the floral phenology, pollen production, and characteristics of *Salix alba*. Over two years, significant variations were observed in flowering timing, duration, and pollen traits among individual trees. Floral phenology assessments revealed diverse patterns in the initiation, duration, and cessation of flowering events. Male catkins typically initiated flowering earlier than females, reflecting the species' adaptive response to environmental cues. Pollen production varied considerably among trees, with tree T5 showing the highest yield at approximately 4.6 billion pollen grains and tree T6 the lowest at 498.3 million grains per tree. This diversity underscores genetic variability and its impact on reproductive success. Analysis of pollen viability, size, and the pollen-ovule ratio highlighted differences in reproductive potential across trees. Pollen viability ranged from 75% to 80%, while pollen sizes varied from 16 μm to 26 μm . These findings underscore the species' reproductive strategies and adaptive flexibility.

Understanding these dynamics is crucial for conservation and management strategies, particularly in the context of environmental changes. Future research could delve deeper into the genetic underpinnings of these variations, offering further insights into plant reproductive biology and ecosystem resilience.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

UNDER PEER REVIEW

References

- Alexander, M.P. 1969. Differential staining of aborted and non-aborted pollen. *Stain Technology*. 44: 117.
- Anonymous. 1983. Forestry and Agroforestry Practices in Kashmir: A Historical Overview. Department of Agriculture, Government of Jammu and Kashmir, Srinagar, India.
- Argus, G.W. 1997. Infrageneric classification of *Salix* (Salicaceae) in the new world. Systematic botany monographs. *The American Society of Plant Taxonomists, USA* ' 52: 1.
- Ascari, L., Novara, C., Dusio, D. and Siniscalco, C. 2020. Quantitative methods in microscopy to assess pollen viability in different plant taxa. *Plant Reproduction*. **33**(3): 205-219.
- Choudhary P, Singh, N.B., Thakur, I.K., Sharma, J.P., Sharma, A. and Verma, A. 2011. Phenological behaviour and reproductive biology of important fast-growing *Salix* species. *Indian Journal of Ecology*. **38**: 99-106.
- Cruden, R. W. 1977. Pollen-ovule ratios: a conservative indicator of breeding systems in flowering plants. *Evolution*. **31**: 32-46.
- Cruden, R.W. and Miller-Ward, S. 1981. Pollen-ovule ratio, pollinations, and potential seed set in *Poinsettia* (*Euphorbia pulcherrima*). *New Phytologist*. **89**(3): 579-593.
- Dafni, A. 1992. Pollination ecology: a practical approach. *Oriole Press Oxford University* 14-250.
- Dickmann, D.I. and Kuzovkina, J. 2014. Poplars and Willows in the World. In *Poplars and Willows: Trees for Society and the Environment*; Isebrands, J.G., Richardson, J., Eds.; Chapter: Poplars and Willows in the World Publisher; FAO: Rome, Italy, pp. 8–91.
- Ghelardini, L., Santini, A. and Fissi, B. 2010. Influence of temperature on flowering phenology of temperate tree species. *Tree Physiology*. **30**(1): 213-223.
- Gupta, A., Singh, N.B., Choudhary, P., Sharma, J.P. and Sankhayan, H.P. 2014. Advances in agricultural practices *International Journal of Agriculture, Environment and Biotechnology*. **7**(2): 299-304.
- Helfenstein, J., Bauer, L., Clalüna, A., Bolliger, J. and Kienast, F. 2014. Landscape ecology meets landscape science. *Landscape ecology*. **29**(7): 1109-1113.

Kearns, A. C. and Inouye, D. W. 1993. Techniques for pollination biologists. University of Colorado, pp. 583.

Khanduri, V. P. 2012. Annual variation in floral phenology and pollen production in a 25-year old plantation of *Tectonagrandis*. *Nordic Journal of Botany*. **30**: 82-89.

Kuzovkina, Y.A. and Quigley, M.F. 2005. Willows beyond wetlands: Uses of *Salix* L. species for environmental projects. *Water Air and Soil Pollution*. **162**: 183-204.

Luna, R.K. 1995. Plantation trees. Publishing Corporation. IBD, Dehradun, pp 640-645.

Masoodi, T.H., Masoodi, N.A., Islam, M.A., Zaffer, S.N. and Chauhan, S.K. 2008. Basketry from introduced willows in Kashmir: Old tradition to new economically viable art. *Indian Forester*. **134**: 77-87.

Mir, R.A., Bhat, G.M., Rafeeq, J. and Wagay, O.A. 2024. Seed maturity indices in indian bean tree (*Catalpa bignonioides* Walt.) under temperate conditions of Kashmir Himalayas. *SKUAST Journal of Research* **26**(1): 116-120.

Pei, M.H. and McCracken, A. 2005. *Rust diseases of willow and poplar*. CABI Publishing, UK.

Rafeeq, J., Qaisar, K.N., Mugloo, J.A., Singh, A., Raj, A Khan, P.A., Khan, I., Hassan, I. and Mir, J.I. 2022. Phenological Characteristics of Hawthorn, *Crataegussongarica* K Koch. in Kashmir Himalayas. *SKUAST Journal of Research*. **24**(2): 248-252.

Raj, A.J. and Lal, S.B. 2014. Agroforestry: Theory and Practices. Scientific publishers, India. pp. 926.

Saunders, D.A. 2020. Ambophily in *Salix* species: Insect and wind pollination. *Plant Reproduction*. **33**(2): 153-164.

Skvortsov, A.K. 1999. *Willows of Russia and Adjacent Countries: Taxonomical and Geographical Revision*. Joensuu, Finland: University of Joensuu, Faculty of Mathematics and Natural Sciences. (Translated from Russian).

Stott, K.G. 1984. Improving the biomass potential of Willows by selection and breeding. **In: Ecology and management of forest biomass. Production system** (Ed. Pertter, K.). Uppsala: Swedish University of Agricultural Science. *Department of Ecology and Environmental Research*. **15**: 233-260.

Tawfeek, N., Mahmoud, M. F., Hamdan, D, I., Sobeh, M., Farrag, N., Wink, M. and El-Shazly, A.M. 2021. Phytochemistry, Pharmacology and Medicinal Uses of Plants of the Genus *Salix*: An Updated Review. *Frontiers in Pharmacology*. **12**.

Trybush, S., Jahodova, S., Macalpine, W. and Karp, A. 2008. A genetic study of *Salix* germplasm resource reveals new insights into relationships among subgenera, sections and species. *Bioenergy Research*. **1**(1): 67-79.

Vitasse, Y., Delzon, S., Bresson, C.C., Michalet, R. and Kremer, A. 2009. Altitudinal differentiation in growth and phenology among populations of temperate-zone tree species growing in a common garden. *Journal of Evolutionary Biology*. **22**(12):2020-2029.

Wagay, O.A., Raj, A., Khan, P.A., Mugloo, J.A., Mir, A.A. and Rafeeq, J. 2024. Floral Biology of Chinar (*Platanus orientalis* L.) under Temperate Conditions in the Kashmir Himalayas. *International Journal of Environment and Climate Change*. **14**(6): 155-164.

Oberprieler C, Dietz L, Harlander C, Heilmann J. Molecular and phytochemical evidence for the taxonomic integrity of *Salix alba*, *S. fragilis*, and their hybrid *S. × rubens* (Salicaceae) in mixed stands in SE Germany. *Plant systematics and evolution*. 2013 Jun;299:1107-18.



Fig 1. Male catkin of *Salix alba*



Fig 2. Female catkin of *Salix*




Fig 3. Ovules of *Salix alba*



Fig 4. Pollens of *Salix alba*




Fig 5. Pollen viability of *Salix alba*

UNDER PEER REVIEW