

## Heterosis and Combining Ability Analysis in Tomato using Line $\times$ Tester Model

### ABSTRACT:

The experiment entitled “Heterosis and combining ability studies for yield and yield attributing trait in Tomato (*Lycopersicon esculentum*) in Bundelkhand Region” was carried out during Rabi 2021-22 and 2022-23 was carried at the Experimental, Organic Research farm Kargunwa ji, Jhansi, Department of Horticultural Sciences, Institute of Agricultural Sciences, Bundelkhand University Jhansi (Uttar Pradesh). The present experiment was design under Randomized block design with three replication with plot size - 3 $\times$ 3m and number of rows per plant – 5 rows per plant accommodating spacing (60 $\times$ 60) cm. All F<sub>1s</sub> (50) along with parents were raised in RBD in 3 replications and population size were 30 plants per treatment with standard spacing. All 17 parents (5 lines and 12 testers) were raised in separate plots along with a check grown in RBD in three replications to study the combining ability, to estimate the magnitude or percentage of heterosis in the crosses, for identification of the best F<sub>1</sub> hybrid and correlation among different traits. Since, Line  $\times$  tester analysis provides information of potential parents the hybrid line. The three best combinations for line parents viz., H-88-78-5, Kashi Aman, H-88-78-5 and their F<sub>1</sub> hybrid viz., H-88-78-1 $\times$  Kashi Chaya, VRT-67  $\times$  Kashi Chaya, H-88-78-5 $\times$ VRT-50 performed significantly better amongst rest of the other treatment.

**Key words:** Heterosis, Line, Tomato, (*Lycopersicon esculentum* L.), tester.

### 1.INTRODUCTION

The solanaceae family include tomato *Solanum lycopersicum* L., which is asexually reproduce annual herbaceous vegetable crops. One of the most extensively grown vegetable crops it has 2n=2x=24 chromosomes is a diploid species. According to [1], the tomato is said to have originated in Peru Ecuador. The Portugese introduce it to the Indian *pennelli*, *L.hirsutum*, *L.chinese* and *L.peruvianum* species. According to [2] the genus *Lycopersicon* has 9 closely related species, including *L.esculentum*, *L. pimpinefolium*, *L.cheesmaniae*, *L.perviflorum*, *L.chnielewskii* and L. species. The word tomato in English derived from the Tomate which itself derived from the Mexican word *Tomatal*. Tomato output is second only to the potato production world wide [3]. India is the world second largest producer of the tomatoes, right after China.

According to India 3<sup>rd</sup> estimate of production figure, 778000 hectares produced over 19397000 Metric Tonnes in 2018-19 (*nhb.gov.in*) the total tomato production in India is produced in the major tomato growing states of Karnataka, Odisha, Madhya Pradesh, West Bengal, Chhattisgrah, Gujrat and Telangana, Bihar Uttar Pradesh, Tamil Nadu. The output of tomato fruits in the years 2022-23 is anticipated to be 3.8% greater than in prior year, according to the first advance estimate. After maize, wheat and cotton, it ranks fifth in terms of crop value at the moment [4].

According to [5], the line x Tester mating design is essentially a important for development of top cross analysis in that multiple testers are used as opposed to just one in top cross. Together and individually, they all contribute a shared genetic background that the inbreds' genotype is measured against. A line is tested due to the utilization of several testers in vegetable crops. The first design that simultaneously gives both full sibling (FS) and half sibling (HS) relationships is the line x tester mating design (top cross and poly cross design only supply HS). Half-siblings are people with one parent in common and a different parent. Therefore, a group of half-siblings is one person's offspring. Therefore, a group of half-sibs is the offspring of one person who mated with a random group of the other person and had one child by each mate. Full siblings, on the other hand, share both parents, and as a result, the mean genotypic value of a full sibling group is equal to the average breeding value of two parents [6]. Thus, FS/HS analysis is another name for line x tester.

## **2.MATERIALS AND METHODS**

The experiment entitled “Heterosis and combining ability studies for yield and yield attributing trait in Tomato (*Lycopersicon esculentum*) in Bundelkhand Region” was carried out during *Rabi* 2021-22 and 2022-23 was carried at the Experimental, Organic Research farm Kargunwa ji, Jhansi, Department of Horticultural Sciences, Institute of Agricultural Sciences, Bundekhand University Jhansi (Uttar Pradesh). The present experiment was conducted in a Randomized block design (RBD) with three replication to appraise the performance of 17 parents (5 lines and 12 testers) which were selected based upon their performance for various traits. The characteristics features of the parents involved in the study are given below in Randomized block design with three replication.

### **2.1Hybridization program**

Each of the 5 lines were crossed with 12 testers to produce 60 hybrids and  $F_{1S}$  was allow to selfing to produce  $F_{2S}$ . The healthy flower buds from new flush, which were due to open next day, were selected emasculation and pollination. The selected buds were emasculated by hand using forceps in the evening hours between 4:00pm to 5:30 pm. Emasculated flowers were covered with cotton to avoid contamination by foreign pollen. Pollination of the emasculated flowers was done by next day after the anthesis time (8:30am to 9:00 am). Well opened flowers with dehisced anthers were collected from male parents, the cotton were removed carefully and the stigma was touched with dehisced anthers of the male flowers. The female flowers were covered with white colour butter paper bag immediately for easy identification and further avoiding the contamination from other pollen. The pedicel of each pollinated flowers was tied

with the label, bearing information of female and male parents and date of crossing for identification.

### **3. RESULTS**

#### **Mean performance of parents**

Mean performance of parents with respect to plant height as presented in (Table 1) in F<sub>1</sub> hybrids Line 97.34 cm (H-88-78-5), tester 90.26 cm (Kashi Aman) with 98.11cm (H-88-78-1x Kashi Chaya). The number of fruit per cluster as in F<sub>1</sub> hybrids Line 13.18 (ToLcv-15), tester 11.54 (Kashi Aman) with 98.11cm (H-88-78-1x Kashi Chaya). The number of branches per plant as in F<sub>1</sub> hybrids Line 18.89 (H-88-78-1), tester 18.93 (Kashi Chaya) with 98.11cm (H-88-78-1x Kashi Chaya). The number of fruits per plant as in F<sub>1</sub> hybrids Line 15.16 (H-88-78-5), tester 14.06 (VRT-30) with 98.11cm (H-88-78-1x Kashi Chaya). 50% flowering was as in F<sub>1</sub> hybrids Line 54.23 (TOLcv-32), tester 49.10 (VRT-30) with 98.11cm (H-88-78-1x Kashi Chaya). For number of cluster per plant was as in F<sub>1</sub> hybrids Line 7.37 (TOLcv-32), tester 49.10 (VRT-30) with 98.11cm (H-88-78-1x Kashi Chaya). For number of flower per plant was as in F<sub>1</sub> hybrids Line 7.77 (H-88-78-1), tester 7.40 (VRT-30) with 98.11cm (H-88-78-1x Kashi Chaya). For number of fruit per cluster was as in F<sub>1</sub> hybrids Line 6.75 (H-88-78-5), tester 6.40 (Vaibhav) with 98.11cm (H-88-78-1x Kashi Chaya). For days to first fruit harvest was as in F<sub>1</sub> hybrids Line 55.95 (H-88-78-1), tester 54.92 (VRT-50) with 98.11cm (H-88-78-1x Kashi Chaya). For fruiting duration was as in F<sub>1</sub> hybrids Line 51.80 days (H-88-78-5), tester 52.63 days (Kashi Chaya) with 98.11cm (H-88-78-1x Kashi Chaya). For fruit length was as in F<sub>1</sub> hybrids Line 7.69 (H-88-78-5), tester 7.37 (Kashi Chaya) with 98.11cm (H-88-78-1x Kashi Chaya). For fruit width was as in F<sub>1</sub> hybrids Line 8.27 (H-88-78-5), tester 7.33 (Kashi Aman) with 98.11cm (H-88-78-1x Kashi Chaya). For Lycopene content was as in F<sub>1</sub> hybrids Line 11.27 (H-88-78-1), tester 10.51 (Kashi Chaya) with 98.11cm (H-88-78-1x Kashi Chaya). For number of yield per plant (g) was as in F<sub>1</sub> hybrids Line 113.57 (H-88-78-1), tester 106.10 (Kashi Chaya) with 98.11cm (H-88-78-1x Kashi Chaya).

**Table 1: Mean performance of F<sub>1</sub>-hybrids of Genotypes**

S.No	Parents	Plant height (cm)	Number of fruit per cluster	Number of branches per plant	Number of fruits per plant	50% flowering g	Number of cluster per plant	Number of flower per cluster	Number of fruit per cluster	Days to first fruit harvest	Fruiting duration	Fruitlen gth	Fruit width	Lycopene content	Yield per plant (g)
	<b>Line</b>														
1	H-88-78-5	97.34	12.48	17.80	15.16	44.37	7.30	7.77	6.75	63.67	51.80	7.47	8.27	11.21	103.82
2	H-88-78-4	94.65	12.20	17.07	14.13	48.70	7.10	7.10	6.68	59.57	53.53	6.73	7.93	10.56	108.65
3	VRT-67	95.11	12.30	16.32	14.88	44.73	7.17	7.47	6.72	56.28	51.13	6.77	7.97	11.31	110.69
4	TOLcV-15	93.79	13.18	17.00	14.06	50.43	7.13	7.27	6.59	60.08	53.83	6.89	7.37	10.34	105.94
5	VRT-16-1	92.88	12.87	17.90	14.03	49.33	7.17	7.20	6.29	60.54	54.63	6.80	7.33	10.30	103.82
6	VRT-06	93.40	12.29	16.97	13.14	50.50	7.20	7.40	6.29	58.40	55.37	6.66	7.47	10.38	102.23
7	VRT-19	92.80	12.14	17.94	13.34	52.58	7.23	7.30	6.59	60.11	54.30	6.66	7.37	10.18	100.24
8	H-88-78-1	95.09	11.82	18.89	12.97	47.03	7.37	7.40	6.74	55.95	52.17	7.69	7.87	11.27	113.57
9	VRT-51	90.29	11.53	17.99	13.37	50.33	7.03	7.27	6.60	59.74	54.30	6.37	7.23	10.10	101.63
10	TOLcV-28	91.23	11.34	17.97	13.99	53.33	7.07	7.27	6.52	58.88	53.33	6.27	7.33	10.17	99.29
11	VRT-50	91.98	11.65	17.87	12.89	53.33	6.87	7.20	6.60	58.44	53.40	6.26	7.47	9.48	99.43
12	TOLcV-32	90.36	11.62	18.38	13.07	54.23	6.83	7.30	6.70	59.44	55.47	6.57	6.87	9.88	98.40
	<b>Tester</b>														
1	KASHI CHAYA	89.48	11.28	18.93	13.91	45.47	7.07	7.20	6.37	54.95	52.63	7.37	7.27	10.51	106.10
2	VAIBHAV	90.24	11.47	18.44	13.82	47.63	7.17	7.23	6.40	58.73	54.30	6.83	7.17	10.21	100.55
3	KASHI AMAN	90.26	11.31	18.30	13.93	47.73	6.93	7.23	6.34	59.67	54.00	6.77	7.33	9.93	98.05
4	VRT-50	89.30	11.54	17.81	13.67	48.60	6.90	7.33	6.32	54.92	53.77	6.37	6.77	9.94	97.10
5	VRT-30	88.71	11.42	18.67	14.06	49.10	6.93	7.40	6.34	64.57	54.77	6.43	6.20	9.36	94.85
	Mean	<b>92.17</b>	<b>11.91</b>	<b>17.90</b>	<b>13.79</b>	<b>49.26</b>	<b>7.09</b>	<b>7.31</b>	<b>6.52</b>	<b>59.06</b>	<b>53.69</b>	<b>6.76</b>	<b>7.36</b>	<b>10.30</b>	<b>102.61</b>
	Min	88.71	11.28	16.32	12.89	44.37	6.83	7.10	6.29	54.92	51.13	6.26	6.20	9.36	94.85
	Max	97.34	13.18	18.93	15.16	54.23	7.37	7.77	6.75	64.57	55.47	7.69	8.27	11.31	113.57

**Table 2: Mean performance of F<sub>1</sub>-hybrids and their crosses.**

Sr. No	Parents	Plant height (cm)	Number of fruit per cluster	Number of branches per plant	Number of fruits per plant	50% flowering	Number of cluster per plant	Number of flower per cluster	Number of fruit per cluster	Days to first fruit harvest	Fruiting duration	Fruit length	Fruit width	Lycopene content	Yield per plant (g)
1	H-88-78-5 × KASHI CHAYA	93.34	12.41	14.51	14.02	45.53	7.20	7.17	6.67	64.26	56.57	6.70	6.93	11.24	117.02
2	H-88-78-5 × VAIBHAV	90.48	11.39	19.83	15.13	54.00	6.73	7.37	6.67	65.12	87.67	6.50	7.30	11.19	117.55
3	H-88-78-5 × KASHI AMAN	92.66	12.09	18.41	15.23	50.63	6.63	7.53	6.63	56.40	88.29	6.70	7.27	11.20	118.86
4	H-88-78-5 × VRT-50	92.65	12.09	17.68	14.67	50.93	6.50	6.33	6.50	69.23	85.60	6.53	7.30	11.19	113.17
5	H-88-78-5 × VRT-30	92.59	12.49	19.10	14.63	56.27	6.20	6.63	6.50	64.69	88.00	6.57	6.77	11.19	109.36
6	H-88-78-4 × KASHI CHAYA	92.52	12.22	15.38	13.87	45.27	7.27	7.40	6.80	63.26	56.17	6.87	6.83	10.49	118.34
7	H-88-78-4 × VAIBHAV	90.04	12.08	19.99	15.22	52.07	6.77	7.53	6.73	64.75	84.83	6.67	7.37	10.53	111.61
8	H-88-78-4 × KASHI AMAN	92.37	12.15	18.42	15.23	47.32	6.47	7.40	6.77	56.28	90.20	6.77	7.37	10.54	112.37
9	H-88-78-4 × VRT-50	92.77	12.71	17.51	14.80	51.93	6.57	6.50	6.50	68.55	85.30	6.67	7.33	10.54	109.12
10	H-88-78-4 × VRT-30	93.11	13.25	19.02	14.23	55.07	6.17	7.60	6.63	61.89	87.57	6.73	7.27	10.53	109.06
11	VRT-67 × KASHI CHAYA	97.88	12.21	16.84	14.07	44.20	7.37	7.70	7.10	54.42	54.10	7.10	8.00	11.33	124.86
12	VRT-67 × VAIBHAV	89.66	12.12	20.28	14.86	51.93	7.03	6.80	6.77	63.66	85.97	6.77	6.70	11.28	116.09
13	VRT-6 × KASHI	92.92	12.18	18.45	14.97	51.30	6.50	7.47	6.70	62.70	88.73	6.73	6.70	11.28	115.59

	AMAN														
14	VRT-67 × VRT-50	93.50	12.39	17.93	15.47	52.93	6.53	6.73	6.47	64.75	86.63	6.63	6.60	11.34	111.21
15	VRT-67 × VRT-30	93.37	12.21	18.73	14.30	55.20	6.20	6.73	6.40	63.63	85.67	6.53	7.30	11.26	110.70
16	TOLeV-16 × KASHI CHAYA	93.81	13.15	15.84	14.33	46.17	7.17	7.50	6.67	56.81	57.60	6.80	7.27	10.30	115.09
17	TOLeV-16 X×VAIBHAV	92.00	12.38	21.31	14.39	52.93	7.07	6.77	6.83	60.59	85.50	6.80	7.20	10.30	114.03
18	TOLeV-16 × KASHI AMAN	91.63	12.38	18.60	15.57	51.93	6.43	7.53	6.70	63.77	87.67	6.83	7.20	10.28	110.41
19	TOLeV-16 × VRT-50	89.78	10.22	18.59	12.97	55.87	5.70	7.27	5.73	70.50	91.17	8.13	7.97	12.22	98.51
20	TOLeV-16 × VRT-30	93.95	12.10	17.81	14.30	55.19	6.53	6.63	6.37	62.93	86.60	6.43	7.37	10.27	108.63
21	VRT-16-1 × KASHI CHAYA	92.60	13.02	16.93	14.20	48.23	7.17	7.60	6.87	55.86	87.00	6.97	7.43	10.28	113.85
22	VRT-16-1 × VAIBHAV	89.74	11.49	19.73	14.56	53.00	6.83	6.63	6.83	59.89	84.60	6.80	7.33	10.21	113.17
23	VRT-16-1 × KASHI AMAN	91.80	12.09	19.38	15.37	51.10	6.50	7.43	7.03	64.64	86.43	7.13	7.27	10.23	112.23
24	VRT-16-1 × VRT-50	90.00	10.38	18.67	13.37	56.87	5.67	7.50	5.67	71.21	91.07	8.30	7.80	12.41	94.93
25	VRT-16-1 × VRT-30	92.63	12.34	17.95	14.63	56.09	6.47	6.50	6.40	65.69	85.03	6.50	6.73	10.23	109.25
26	VRT-06 ×KASHI CHAYA	94.18	12.27	17.81	14.18	46.17	7.00	7.43	7.07	58.35	88.59	6.73	7.60	10.36	109.92
27	VRT-06 × VAIBHAV	90.66	11.39	19.95	15.03	54.03	6.83	6.67	6.77	68.16	87.27	6.80	7.43	10.33	108.49
28	VRT-06 × KASHI AMAN	92.74	12.06	18.34	15.07	51.97	6.50	6.67	6.30	65.59	87.00	6.33	7.37	10.31	107.99
29	VRT-06 × VRT-50	89.59	10.50	18.59	12.93	57.23	5.63	7.53	5.67	70.54	91.23	8.30	7.87	12.30	91.60
30	VRT-06	93.80	12.29	18.09	14.57	57.41	6.67	6.41	6.40	64.61	84.20	6.50	7.27	10.31	108.40

	×VRT-30														
31	VRT-19 × KASHI CHAYA	93.98	12.09	18.46	14.06	44.98	7.03	7.50	6.83	57.87	87.83	6.77	6.70	10.22	107.84
32	VRT-19 × VAIBHAV	92.19	11.27	18.55	14.83	54.43	6.77	7.07	6.50	68.25	87.03	6.47	6.77	10.18	106.94
33	VRT-19 × KASHI AMAN	92.64	12.77	18.77	14.73	53.00	6.53	6.77	6.30	66.75	85.40	6.33	6.73	10.19	106.64
34	VRT-19 × VRT-50	90.03	10.33	18.75	13.00	59.30	5.73	7.60	5.67	72.59	92.47	8.37	7.47	12.41	94.38
35	VRT-19 ×VRT-30	94.12	11.88	18.11	14.57	56.17	6.67	6.38	6.57	62.17	87.20	6.63	7.27	10.17	106.17
36	H-88-78-1 × KASHI CHAYA	98.11	11.92	22.64	15.50	44.47	7.27	7.83	6.93	56.12	88.60	6.87	7.23	11.28	123.16
37	H-88-78-1 X×VAIBHAV	92.63	11.26	18.81	14.53	53.07	6.77	7.23	6.47	69.70	86.57	6.50	7.27	11.23	114.73
38	H-88-78-1 × KASHI AMAN	92.81	12.59	19.31	14.23	52.93	6.47	6.50	6.20	66.70	84.53	6.37	7.20	11.26	114.95
39	H-88-78-1 × VRT-50	89.81	11.16	18.48	12.93	58.33	5.67	7.80	5.63	72.55	93.83	8.40	7.50	12.53	98.04
40	H-88-78-1 × VRT-30	94.39	11.78	19.32	14.56	55.23	6.57	6.30	6.27	61.29	86.33	6.37	7.37	11.24	110.63
41	VRT-51 × KASHI CHAYA	92.91	11.75	18.66	15.10	55.20	7.13	7.40	6.77	58.44	88.60	6.70	7.30	10.13	112.27
42	VRT-51 × VAIBHAV	93.70	11.26	18.95	14.63	53.03	6.70	7.30	6.57	68.74	88.06	6.50	7.33	10.13	113.12
43	VRT-51 × KASHI AMAN	93.32	12.37	19.09	14.07	51.40	6.50	6.53	6.27	65.70	87.96	6.27	7.27	10.14	111.72
44	VRT-51 × VRT-50	89.22	10.24	17.88	12.97	57.77	5.67	7.63	5.50	72.82	90.50	8.37	7.50	12.56	96.82
45	VRT-51 × VRT-30	94.52	11.78	19.65	14.22	57.52	6.60	6.37	6.27	59.86	85.77	6.33	6.77	10.07	108.22
46	TOLeV-28 × KASHI CHAYA	91.66	11.67	18.62	15.16	53.00	7.03	7.30	6.57	59.41	86.60	6.43	7.40	10.18	110.08

47	TOLeV-28 × VAIBHAV	92.04	12.28	19.22	14.90	54.00	6.70	7.23	6.57	70.00	85.60	6.50	7.43	10.18	110.64
48	TOLeV-28 × KASHI AMAN	93.18	12.46	18.42	14.87	49.37	6.50	6.37	6.27	69.28	87.70	6.43	7.40	10.14	108.97
49	TOLeV-28 × VRT-50	92.99	11.14	18.54	14.17	50.73	6.00	6.83	6.30	69.89	88.60	6.40	7.33	10.14	112.97
50	TOLeV-28 × VRT-30	94.54	12.04	19.31	14.47	55.23	6.53	6.47	6.30	60.47	88.89	6.40	7.27	10.14	111.27
51	VRT-50 ×KASHI CHAYA	92.11	11.48	19.51	15.00	54.07	7.00	7.23	6.83	60.11	86.80	6.73	6.77	9.50	109.68
52	VRT-01 ×VAIBHAV	94.11	12.40	18.63	14.67	53.10	6.70	7.23	6.43	68.85	86.35	6.47	6.73	9.50	109.63
53	VRT-01 X×KASHI AMAN	93.04	11.89	18.80	14.57	50.33	6.37	6.37	6.73	69.53	87.23	6.83	6.70	9.50	107.61
54	VRT-01 × VRT-50	92.56	12.18	18.49	14.50	53.47	6.37	7.20	6.23	62.95	86.60	6.43	7.50	9.47	108.73
55	VART-01 × VRT-30	92.59	12.10	19.46	14.37	57.23	6.53	6.53	6.47	64.57	85.57	6.57	7.33	9.47	107.51
56	TOLeV-32 × KASHI CHAYA	92.28	11.50	20.95	15.17	53.03	6.87	7.37	6.60	61.18	87.30	6.53	7.23	9.70	110.90
57	TOLeV-32 × VAIBHAV	93.92	12.17	18.56	15.50	50.27	6.70	7.47	6.37	64.32	87.02	6.47	7.23	9.73	110.57
58	TOLeV-32 × KASHI AMAN	92.78	12.11	18.49	14.60	50.03	6.43	6.33	6.53	69.29	88.78	6.60	7.27	9.74	109.54
59	TOLeV-32 × VRT-50	93.88	12.38	18.92	14.80	54.80	6.23	7.30	6.27	63.61	86.80	6.30	7.37	9.77	108.53
60	TOLeV-32 × VRT-30	94.29	12.13	19.93	14.43	56.19	6.17	6.70	6.17	64.30	85.67	6.30	7.40	9.66	105.64
	Mean F <sub>1</sub>	<b>92.65</b>	<b>11.94</b>	<b>18.68</b>	<b>14.52</b>	<b>52.67</b>	<b>6.58</b>	<b>7.05</b>	<b>6.48</b>	<b>64.33</b>	<b>85.31</b>	<b>6.77</b>	<b>7.24</b>	<b>10.60</b>	<b>109.99</b>
	Min	89.22	10.22	14.51	12.93	44.20	5.63	6.30	5.50	54.42	54.10	6.27	6.60	9.47	91.60
	Max	98.11	13.25	22.64	15.57	59.30	7.37	7.83	7.10	72.82	93.83	8.40	8.00	12.56	124.86
	Mean All	<b>92.55</b>	<b>11.93</b>	<b>18.51</b>	<b>14.36</b>	<b>51.92</b>	<b>6.69</b>	<b>7.11</b>	<b>6.48</b>	<b>63.17</b>	<b>78.33</b>	<b>6.77</b>	<b>7.26</b>	<b>10.53</b>	<b>108.36</b>
	Min	88.71	10.22	14.51	12.89	44.20	5.63	6.30	5.50	54.42	51.13	6.26	6.20	9.36	91.60
	Max	98.11	13.25	22.64	15.57	59.30	7.37	7.83	7.10	72.82	93.83	8.40	8.27	12.56	124.86
	SE(d) ±	0.52	0.09	0.19	0.36	0.59	0.11	0.13	0.15	0.82	0.68	0.13	0.08	0.10	1.11
	C.D. at 5%	1.03	0.18	0.37	0.70	1.17	0.22	0.25	0.29	1.62	1.35	0.26	0.16	0.19	2.20
	C.V. (%)	0.69	0.92	1.25	3.04	1.40	2.04	2.19	2.79	1.59	1.07	2.35	1.35	1.14	1.26

## DISCUSSION

### Mean performance of parents

From the present findings the mean performance of parents with respect to plant height in F<sub>1</sub> hybrids Line (H-88-78-5), tester (Kashi Aman) and (H-88-78-1x Kashi Chaya). It may be noted that plant height was superior due to interaction components which was highly significant in terms of plant height. The present results are in line with [7,8,9]. The number of fruit per cluster as in F<sub>1</sub> hybrids Line (ToLcv-15), tester (Kashi Aman) and (H-88-78-1x Kashi Chaya). It may be well noted that number of fruit per cluster was superior due to interaction components which was highly significant in terms of number of fruit per cluster. The number of branches per plant as in F<sub>1</sub> hybrids Line (H-88-78-1), tester (Kashi Chaya) with (H-88-78-1x Kashi Chaya). The present study revealed that maximum branches were significant and superior as most significant contribution of line and tester observed for number of branches. The findings were similar to the studies with [10,11,12] in tomato. The number of fruits per plant as in F<sub>1</sub> hybrids Line (H-88-78-5), tester (VRT-30) with (H-88-78-1x Kashi Chaya). The present study revealed that maximum number of fruits per plant were significant and superior as most significant contribution of line and tester observed for number of fruits per plant. The data on days to 50% flowering was as in F<sub>1</sub> hybrids Line (TOLcv-32), tester (VRT-30) and (H-88-78-1x Kashi Chaya). The present study revealed that maximum number of days to 50% flowering were non-significant and superior in comparison to minimum number of days to 50% flowering as most significant contribution of line and tester observed for days to 50% flowering. The present findings were similar to the studies [13,15,14] in tomato. For number of cluster per plant was as in F<sub>1</sub> hybrids Line (TOLcv-32), tester (VRT-30) with (H-88-78-1x Kashi Chaya). The present study revealed that maximum number of cluster per plant were significant and superior in comparison to minimum number of cluster per plant as most significant contribution of line and tester observed for days to 50% flowering. The number of flower per plant was as in F<sub>1</sub> hybrids Line (H-88-78-1), tester (VRT-30) with (H-88-78-1x Kashi Chaya). It may be well referred from the present study that number of flower per plant were significantly better under the hybrid lines and tester. The test of significance of variance due to line x tester exhibited a significant variation in terms of number of flower per plant. The number of fruit per cluster was as in F<sub>1</sub> hybrids Line (H-88-78-5), tester (Vaibhav) with (H-88-78-1x Kashi Chaya). The mean performance of line, tester and their crosses clearly indicated that there is no single genotype involved or revealed the superiority in all the traits. As a general trend it may be well established from the present result that better performance of the lines, tester and their hybrids clearly indicated significant variation in terms of number of fruit per cluster. The present result is in agreement with that of [16,18,17] in tomato. For days to first fruit harvest was as in F<sub>1</sub> hybrids Line (H-88-78-1), tester (VRT-50) with (H-88-78-1x Kashi Chaya). It may be well noted that days to first fruit harvest was superior due to interaction components which was highly significant in terms of days to first fruit harvest. The present results are in line with [19,18,17] in tomato. Fruiting duration was as in F<sub>1</sub> hybrids Line days (H-88-78-5), tester (Kashi Chaya) with (H-88-78-1x Kashi Chaya). It may be inferred from the current study that fruiting duration was superior due to interaction components which was

highly significant in terms of fruiting duration. The present results are in line with [20,21,22] in tomato. For fruit length was as in F<sub>1</sub> hybrids Line (H-88-78-5), tester (Kashi Chaya) with (H-88-78-1x Kashi Chaya). The mean performance of line, tester and their crosses clearly indicated that there is no single genotype involved or revealed the superiority in all the traits for fruit length and its architecture. As a general trend it may be well established from the present result that better performance of the lines, tester and their hybrids clearly indicated significant variation in terms of fruit length. The present result is in agreement with that of [23,24,25,26,27] in tomato. Fruit width was as in F<sub>1</sub> hybrids Line (H-88-78-5), tester (Kashi Aman) with (H-88-78-1x Kashi Chaya). As a general trend it may be well established from the current result that better performance of the lines, tester and their hybrids clearly indicated significant variation in terms of fruit width. The present result are in accordance with that of [28,29,30] in tomato. The Lycopene content was as in F<sub>1</sub> hybrids Line (H-88-78-1), tester (Kashi Chaya) with (H-88-78-1x Kashi Chaya). It may be noted from the present results that lycopene content was higher due to interaction components and the general combiner for most of the traits which was highly significant in terms of lycopene content. The present study are in agreements with that of [31]. Yield per plant (g) was as in F<sub>1</sub> hybrids Line (H-88-78-1), tester (Kashi Chaya) with (H-88-78-1x Kashi Chaya). It may be noted that yield per plant was superior due to interaction components of the best hybrids and their combination. The best hybrid combination reflected the high significant values in terms of plant yield.

## **CONCLUSION**

These pertaining to GCA and SCA effects of different parameters related yield and quality are very important tools for identifying suitable parents to be used in any breeding program. Among all the different line employed in the present research. The (H-88-78-5), tester (Kashi Aman) with (H-88-78-1x Kashi Chaya) was found to be a good general combiner for maximum characters. Path coefficient analysis also revealed that highest number of fruit/cluster exerted the highest direct effect on lycopene content. For Phenotypic path with yield and yield components Path coefficient analysis also revealed the highest pattern for many characters viz., number of fruit duration which exerted the highest direct effect on lycopene content. Thus, obtained results can help in deciding the suitable parents as well as crosses for particular trait that can be better and further utilized in breeding programmes in tomato.

## **FUTURE SCOPE**

Thus, the most suitable inbred lines to be employed in future breeding programmes can be identified thanks to this study, which uses a variety of tomato genotypes. The consequences of crossing showed that some parent were a good general match for many character, indicating that some parents will need to be chosen for genetic enhancement depending on specific qualities taken into account. Because tomatoes are a highly consumed produce, experts are focusing on developing superior hybrids with desired parent combinations through crop improvement programmes. Additionally, hybrids that produce larger yields assist farmers in meeting the ongoing need of the market.

## REFERENCE

1. Rick CM. Tomato. Hybridization of crop plants. 1980 Jan 1:669-80.
2. Esquinas-Alcazar, J. T. "Genetic resources of tomatoes and wild relatives." (1982): 65-pp.
3. Peixoto JV, Neto CM, Campos LF, Dourado WD, Nogueira AP, Nascimento AD. Industrial tomato lines: morphological properties and productivity. Genet. Mol. Res. 2017 Apr 13;16(2):1-5.
4. Kumar BS, Mali SC, Patel AI, Prajapati MR. Appraisal of combining ability and gene action for yield, its governing characters and quality traits in tomato (*Solanum lycopersicum* L.). Electronic Journal of Plant Breeding. 2023;14(2):752-7.
5. Kempthorne, Oscar. "An introduction to genetic statistics.1957; xvii-545.
6. FAOSTAT F. Food and agricultural organization. Roma, Italy. 2014.
7. Amin A, Wani KP, Dar ZA, Jabeen N, Mushtaq F. Heterosis studies in tomato (*Solanum lycopersicum* L.). Journal of Pharmacognosy and Phytochemistry. 2017;6(6):2487-90.
8. Jadav NK, Patel SY, Malviya AV, Patel UV, Vasava HV. Combining ability analysis and gene action for yield, quality and its component traits of tomato (*Solanum lycopersicum* L.). Trends in Biosciences. 2017;10(13):2434-42.
9. Bhalala KC, Acharya RR. Assessment of combining ability using Line× tester analysis over environments in tomato (*Solanum lycopersicum* L.). Journal of Pharmacognosy and Phytochemistry. 2019;8(3):4478-85.
10. Gowthami KJ, Raut N, Jawadagi RS, Chittapur R, Haveri N. Assessment of combining ability in tomato genotypes (*Solanum lycopersicum* L.) for quality traits. Emergent Life Sciences Research. 2022 Jun;8:63-9.
11. Mondal C, Sarkar S, Hazra P. Line× Tester analysis of combining ability in tomato (*Lycopersicon esculentum* Mill.). Journal of Crop and Weed. 2009 Sep 30;5(1):53-7.
12. Kumar K, Sharma D, Singh J, Sharma TK, Kurrey VK, Minz RR. Combining ability analysis for yield and quality traits in tomato (*Solanum lycopersicum* L.). Journal of Pharmacognosy and Phytochemistry. 2018;7(6):1002-5.
13. Kumar K, Sharma D, Singh J, Chandel N. Heterosis for yield and yield attributing traits in tomato (*Solanum lycopersicum* L.). Vegetable Science. 2021;48(1):41-8.
14. Singh PK. Studies on heterosis and combining ability for yield and yield attributes in Tomato (*Lycopersicon esculentum* Mill.). Journal of Pharmacognosy and Phytochemistry. 2017;6(6):2140-3.
15. Raj T, Bhardwaj ML, Pal S. Estimation of heterosis for quality traits in tomato (*Solanum lycopersicum* L.). Journal of Pharmacognosy and Phytochemistry. 2018;7(1):2401-4.
16. Singh S, Singh AK, Singh BK, Singh V, Shikha K. Line× tester analysis for yield and component traits in tomato (*Solanum lycopersicum* L.). Journal of Pharmacognosy and Phytochemistry. 2021;10(1):2044-9.

17. Singh N, Ram CN, Singh V, Kumar D. Line× Tester Analysis and Estimating Combining Abilities for Yield and Some Quality Attributes in Tomato (*Solanum lycopersicum* L.). International Journal of Plant & Soil Science. 2022 Jul 12;34(21):176-94.
18. Kumar R, Srivastava K, Singh NP, Vasistha NK, Singh RK, Singh MK. Combining ability analysis for yield and quality traits in tomato (*Solanum lycopersicum* L.). Journal of Agricultural Science. 2013 Feb 1;5(2):213.
19. El-Mansy AB, Metwally EI, ElKasas AI, El-Tantawy AM, Mahmoud MI. Performance, Heritability and Correlation Coefficients for Some Important Traits in Tomato Under North Sinai Condition. Sinai Journal of Applied Sciences. 2015 Aug 1;4(2):79-94.
20. Reddy BR, Singh AK, Pal AK, Reddy YS, Eswara Reddy G. Combining ability and heterosis studies in tomato (*Solanum lycopersicum* L.) for quality traits and yield. Int. J. Chem. Stud. 2020;8(2):2788-92.
21. Usama M, Hussain A, Rafique A, Naheed K, Kareem M, Hussain Z. Identification of better parents for quality and yield attributing traits in tomato using line× tester analysis. Journal of Physical, Biomedical and Biological Sciences. 2022 Jul 10;2022(1):2-.
22. Kumari S, Sharma MK. Line x tester analysis to study combining ability effects in tomato (*Solanum lycopersicum* L.). Vegetable Science. 2012;39(1):65-9.
23. Narasimhamurthy YK, Gowda PH. Line× tester analysis in tomato (*Solanum lycopersicum* L.): identification of superior parents for fruit quality and yield-attributing traits. International Journal of Plant Breeding. 2013;7(1):50-4.
24. Rehana S, Ullah MZ, Zeba N, Narzis N, Husna A, Siddique AB. Estimation of heterosis for yield and yield attributing traits in tomato crossed with line and tester method. Prog. Agric. 2019;30:179-85.
25. Joshi A, Kohli UK. Combining ability and gene action studies for processing quality attributes in tomato (*Lycopersicon esculentum* Mill.). Indian Journal of Horticulture. 2006;63(3):289-93.
26. Damor HI, Acharya RR, Patel AA. Genetic Analysis for Fruit Yield and its Component Traits in Tomato (*Solanum lycopersicum* L.) Population. Journal of Plant Development Sciences. 2021;13(1):1-9.
27. Pandey SK, Dixit J, Pathak VN, Singh PK. Line x Tester analysis for yield and quality characters in tomato (*Solanum lycopersicon* (Mill.) WETTSD). Veg. Sci. 2006;33(1):13-7.
28. Tekkam S, Pidigam S, Adapa KK, Natarajan S, Amarapalli G. Evaluation of parents and hybrids for yield and yield contributing traits in tomato (*Solanum lycopersicum* L.) for per se performance. Journal of Pharmacognosy and Phytochemistry. 2021;10(1):1770-6.
29. Umesh BC, Patil RV. Combining ability analysis for yield and quality traits in double cross F1s of tomato (*Solanum lycopersicum*L.). Journal of Pharmacognosy and Phytochemistry. 2021;10(4):296-9.

30. Metwally EI, El-Kassas AI, El-Tantawy AM, Mahmoud MI, El-Mansy AB. Heterosis and combining ability in tomato by line x tester. *Journal of Plant Production*. 2015 Feb 1;6(2):159-73.
31. Panthee DR, Perkins-Veazie P, Anderson C, Ibrahem R. Diallel analysis for lycopene content in the hybrids derived from different colored parents in tomato. *American Journal of Plant Sciences*. 2015 Jun 5;6(9):1483-92.

UNDER PEER REVIEW