

Review Article

Grain Legumes in Agricultural Systems: Assessing Their Role in Enhancing Soil Health and Promoting Sustainable Agricultural Practices-A Systematic Review

Suggested title: The Role of Grain Legumes in Enhancing Soil Health and Promoting Sustainable Agricultural Practices: A Systematic Review

ABSTRACT

The imperative for sustainable agricultural practices is widely recognized, with the diversification of cropping systems playing a key role in achieving this goal. Grain legumes are vital to farming systems, offering benefits in terms of food and nutrition security as well as income generation. Utilizing legume-based cropping systems can address multiple agricultural challenges including pest and disease management, low soil fertility, biodiversity conservation, and food insecurity. This chapter discusses the role of legumes in sustainable agricultural systems, specifically focusing on the advantages of integrating legumes within different agroecosystems, [cropping system](#) and farming practices. Legumes enhance crop productivity potential, providing a strategic solution to low crop yields and supporting the shift towards more sustainable agricultural practices. Additionally, due to their health and environmental benefits, it is crucial to focus on breeding grain legumes that optimize their natural ability for biological nitrogen fixation and thrive despite abiotic and biotic stresses([Sepcifically mention the stresses](#)).

INTRODUCTION

Legumes, comprising over 20,000 species, represent the third-largest plant family and are a cornerstone of global food security, ranking just behind cereals in terms of consumption (Shafique et al., 2014). These plants play a crucial role in diets worldwide by providing a significant source of nutritional protein when paired with cereals (Mashungwa et al., 2019). One of the remarkable features of legumes is their ability to convert atmospheric nitrogen into ammonia through symbiotic relationships with rhizobia, a specific type of soil-borne bacteria, thereby enhancing soil fertility (Gonzalez-Rizzo et al., 2009). This symbiosis enables legumes to adapt to adverse environmental conditions and nourish neighboring crops, contributing to sustainable agricultural practices (Popoola et al., 2014).

Legumes are cultivated in various agricultural setups, including intercropping, monoculture, and rotational systems. Intercropping, long practiced by smallholder farmers in regions like Asia, Africa, and Latin America, is noted for maximizing yields from limited resources and space (Yu *et al.*, 2015). In arid areas, crops like pigeon peas([scientific name??](#))

Formatted: Font: Italic

are commonly interplanted with cereals due to their drought resilience (Semahegn, 2022). This method not only conserves space but also enhances cereal growth and productivity through nitrogen fixation and the enrichment of soil organic content (Yuvaraj et al., 2020). When planning intercropping, farmers must consider various factors including species compatibility, planting timing, and the physical characteristics of the plants involved (Semahegn, 2022).

Conversely, monocropping legumes is generally discouraged due to its association with increased pest pressure ([Specific the name of pests](#)) and nutrient depletion ([specify the name of the nutrient](#)) in the soil (Semahegn, 2022). Crop rotation, another sustainable practice, involves alternating different crops like maize and soybean or alfalfa on the same plot to optimize nutrient availability and water usage, which leads to maintained or increased yields (Zhen et al., 2016). Mixed cropping systems, incorporating legumes with cereals or tuber crops, leverage complementary resource use, reducing pest incidents and soil erosion while improving biomass production and yield stability.

Recent years have seen a heightened focus on the roles of legumes in enhancing both human and animal nutrition and in addressing environmental degradation on smallholder farms (ICRISAT, 2016). Acknowledging these benefits, the Food and Agriculture Organization (FAO) of the United Nations highlighted the importance of pulses, designating 2016 as the International Year of Pulses to promote their cultivation (FAO, 2016). This review aims to underline the significant contributions of legume production systems to soil health and the sustainability of agricultural practices.

MATERIALS AND METHODS

The data utilized in this study were predominantly obtained from secondary sources, including scientific publications and official websites of esteemed research institutions. The collection process involved systematic desk research, where relevant materials were identified, downloaded, read, and referenced according to established best practices. [Broadly categorize the sources of data/information](#)

RESULTS AND DISCUSSION

Global Status of Legumes

From 2016 to 2020, the worldwide annual production of grain legumes was approximately 115.28 million tons, as shown in Table 1. The top producers of these legumes are India, Myanmar, Canada, China, and Brazil, which collectively account for half of the global output. The primary legumes cultivated globally include common beans, chickpeas, cowpeas, lentils, pigeon peas, groundnuts, and soybeans, with their respective production details also summarized in Table 1.

According to Nigrum et al. (2021), the common bean leads in cultivation across 126 countries, making it the most extensively grown legume. Groundnuts are cultivated in 114 countries, while soybeans are grown in 89. Other widely grown legumes are chickpeas,

lentils, cowpeas, and pigeon peas. These staples are cultivated on nearly 212 million hectares worldwide, yielding approximately 421 million tonnes annually.

In terms of production efficiency, soybeans dominate with 56.5% of the productivity share, followed by common beans at 14.2%, and groundnuts at 12.8%. Other legumes such as chickpeas contribute 6.0%, cowpeas 5.8%, pigeon peas 2.5%, and lentils 2.2% of the total global production area (Nigrum et al., 2021). [Mention the major legumes production nations' name](#)

[Mention the grass pea also](#)

Table 1: Average Annual Global Production of Grain Legumes from 2016 to 2020

Legume	Production in Million metric tonnes
Field peas	14.9
Chickpeas	50.19
Cowpeas	9.80
Faba beans (Broad beans)	22.55
Lentils	13.34
Pigeon peas	4.50

Table 2: Comparative Chemical Composition of Selected Legumes (Percentage-Based)

Legumes	Botanical names	Proteins	Fat	Carbohydr rates	Fibre
Soybean	<i>Glycine max</i>	37-41	18-21	30-40	4-6
Cowpea	<i>Vigna unguiculata</i>	22-26	1-2	60-65	4-5
Groundnut	<i>Arachis hypogaea</i>	20-33	42-48	22-25	3-4
Hyacinth beans	<i>Lablab Purpureus</i>	24-28	1-2	65-70	7-9

Significance of Grain Legumes for Human and Livestock Nutrition

Grain legumes are pivotal in soil enhancement, and they significantly contribute to the food, feed, and fuel sectors due to their diverse nutritional and chemical properties, which vary by type, species, and soil health. This section explores how grain legumes benefit these key areas.

Nutritional Resource for Agricultural Communities

Grain legumes have been a fundamental component of human diets for millennia, tracing back to the early domestication of crops, and are crucial for achieving food and nutritional security in agricultural settings. The seeds of grain legumes are rich in proteins, starch, soluble vitamins, insoluble fiber, and various micro- and macro-nutrients, along with numerous bioactive phytochemicals (see Table 2) (Watson et al., 2017). Legumes contain

significantly more protein than cereals, typically ranging from 20% to 45% protein content, compared to 7%–17% found in cereals (Day, 2013; Zander et al., 2016). They are excellent sources of the essential amino acid lysine, vital for protein synthesis. However, they generally lack sulphur-containing amino acids like methionine and cysteine, which are crucial for cellular integrity and growth (Friedman, 1996). Most legumes, with the exceptions of soybeans and groundnuts, are low in fat and are also rich in minerals such as calcium and phosphorus. The health benefits of legumes are substantial and vary by species. Soybean-based products, for instance, are known to mitigate heart and vascular diseases by reducing cholesterol levels and managing hypertension (Harland and Haffner, 2008; Sirtori et al., 2009). Further, soybean and lupin have been shown to lower cholesterol in humans (Sirtori et al., 2012). The crucial and varied roles of food legumes within agricultural systems and the diets of economically disadvantaged populations make them key crops for meeting development goals aimed at reducing poverty and hunger, boosting human health and nutrition, and strengthening ecosystem resilience.

Role of Grain Legumes in Livestock Nutrition

Grain legumes serve as an excellent feed option for ruminants, either as part of concentrated compound feeds or whole plant feeds (Sherasia et al., 2018). Their utilization in ruminant diets depends on their chemical composition, how well they complement forage diets, and the rate and extent of their nutritional breakdown in the rumen (Watson et al., 2017). Legumes primarily augment the protein content in animal feeds. The protein degradability of grain legumes in the rumen, similar to most cereal grains, often exceeds 80% (Luke, 2016). Soybeans stand out as the most popular legume and the principal source of supplemental plant protein in animal feeds. Świątkiewicz (2021) estimates that soybeans account for 84% of the high-protein oilseed meals used in compound livestock rations globally, especially in the poultry and pig industries. The reliance on soybean meal is particularly strong in the pig sector due to its high crude protein content (44%) and a favorable amino acid profile, which includes amino acids that are often deficient in other grain legumes like methionine, cysteine, and tryptophan (Watson et al., 2017). However, current research is exploring other legume-based feeds to reduce dependency on soybeans. In the aquaculture sector, efforts are being made to evaluate the use of plant-based proteins, as detailed in a review by Ayadi et al. (2012). Studies have also shown that in the cattle and sheep industries, alternatives such as peas, lupins, rapeseed, or faba beans can replace soybean-based meals as the protein source during different lactation phases without adversely affecting milk production (Khorasani et al., 2001; Froidmont and Bartiaux-Thill, 2004; White et al., 2007; Van der Pol et al., 2008; Tufarelli et al., 2012).

Potential of Legumes in Nitrogen Fixation

Legumes enhance soil fertility through nitrogen fixation, benefiting biodiversity and soil quality (Yuvaraj et al., 2020; Okumu et al., 2017; Couto-Vazquez and González-Prieto, 2016). They form symbiotic relationships with nitrogen-fixing bacteria from various genera of Alphaproteobacteria, Azorhizobium, and Betaproteobacteria (Vasconcelos et al., 2020). These bacteria invade the roots and develop nodules, sites of biological nitrogen fixation.

Here, the bacteria convert atmospheric nitrogen (N₂) into ammonia, a readily available form of nitrogen for plants, facilitated by the nitrogenase enzyme (Howard and Rees, 1996; Ferguson et al., 2013; Ferguson et al., 2019). The efficiency of this biological process depends on the legume species, the bacterial species involved, and abiotic factors such as temperature, water availability, and soil mineral nitrogen content (Otieno et al., 2018; Swaroop and Lal, 2018; Kebede, 2021).

Legumes as Climate-Smart Crops

Besides improving soil health, legumes are recognized as climate-smart crops due to their role in environmental quality enhancement through carbon sequestration and mitigation of other pollutants. Kumar et al. (2018) conducted a meta-analysis indicating that legumes store 30% more soil organic carbon (SOC) than other plant species. The type of legume, growth habits, root morphology, leaf structure, climate, soil structure, cropping system, and stage of agronomic practices all influence their capacity for carbon sequestration and the amount of organic carbon they return to the soil (Kumar et al., 2018). Guan et al. (2016) reported that perennial legumes increase SOC, a process influenced by factors such as turnover, root cell epidermal sloughing, and root exudation of soluble carbon compounds. Legumes also contribute significantly to ecosystem services by reducing dependency on synthetic fertilizers, thereby lowering greenhouse gas emissions (Watson et al., 2017; Stagnari et al., 2017). For example, Gregorich et al. (2005) observed a linear increase in nitrous oxide emissions from soils treated with mineral fertilizers. Legumes have shown potential in reducing greenhouse gas (GHG) emissions, especially nitrous oxide, when used as green manures compared to nitrogen fertilizers, which tend to increase emissions. For instance, the application of alfalfa and other legume crops as green manures has been associated with lower annual nitrous oxide emissions, prompting recommendations for their consideration in national GHG records for agriculture (Abberton, 2010). Legumes are frequently employed as intercrops and cover crops, enhancing soil stability and boosting organic matter, which is crucial for soil formation, fertility, and yields (Martin Körschens, 2002; Howieson et al., 2008). Research by Otieno et al. (2020) demonstrated sustainable and high production levels of dry beans utilizing residual fertilizer from preceding maize crops. However, it's vital to recognize that the management of agroecosystems incorporating legumes also significantly affects their ability to reduce GHG emissions.

Legumes as Biofuels

Legume residues, which often contain higher protein concentrations than cereal crops—up to 10%—offer significant potential as a source of biofuel. Their biomass not only has high protein content but also allows for the extraction of protein as a by-product in biofuel production (Jensen et al., 2012). Since the early 20th century, various high oil and starch content plants like canola (*Brassica napus*), juncea (*B. juncea*), and soybean (*Glycine max*) have been used for biodiesel production (Biswas et al., 2011). Peanut oil, for instance, was first utilized in diesel engines in 1904. The oil from these legumes is extracted by crushing the seeds and squeezing out the oil, which is then transesterified to produce biodiesel. These oil crops can also be transformed into high-value biochemicals and

biomaterials, helping reduce reliance on fossil fuels. Currently, soybean serves as a major feedstock for biodiesel in the United States, while canola and rapeseed are more commonly used in Europe. These plants are highly suitable for biodiesel manufacture due to their ability to produce significant amounts of oil from their seeds annually and their high biomass yield per unit area. Biomass feedstocks for energy production can derive from plants grown specifically for energy, or from plant parts, leftovers, industrial wastes, and materials from animal and human activities (US Department of Energy, 2013).

Grain Legume Production Systems

Grain legumes are versatile crops produced in various forms—such as dry grain, green forage, arable silage, and green manure—depending on climatic and soil conditions and the intended use (Watson and Stoddard, 2017). Several species are cultivated, including pea (*Pisum sativum* L.), lupins (*Lupinus* spp.), faba bean (*Vicia faba* L.), chickpea (*Cicer arietinum* L.), lentil (*Lens culinaris* Medik.), common bean (*Phaseolus vulgaris* L.), and soybean (*Glycine max* (L.) Merr.) (Vanlauwe et al., 2019). Legumes may be grown as standalone crops or intercropped with cereals like maize.

Formatted: Font: Italic

Formatted: Font: Italic

Intercropping

Intercropping involves growing two or more crop species from different families simultaneously, with one being the primary crop and the others serving secondary roles such as nitrogen fixation and soil improvement. This practice is utilized to enhance species interaction, increase biodiversity, and help farmers cope with climate anomalies (Meena and Lal, 2018). The selection of legume intercrops depends on the primary crop, legume species available, maturity periods, growth stature, and farmer preference. Planting configurations and spacing must consider species characteristics, mechanization potential, growth status of the main crop and intercrop, and prevailing climatic conditions and soil fertility. Legume-based intercropping systems typically result in higher production from the same field and improve the efficiency of natural resource use compared to mono-cropping (Inal et al., 2007). Fustec et al. (2010) noted that leguminous crops enhance soil functions through biological nitrogen fixation, offering advantages over monoculture farming systems. Practicing legume-based intercropping promotes eco-friendly agriculture by minimizing the use of inorganic nitrogen fertilizers, thus supporting sustainable agricultural production systems.

Crop Rotation

Crop rotation is a farming system where different crops are grown in a specific recurrent sequence on the same land, as opposed to the continuous cultivation of a single crop (monoculture) (Sumner, 2018). This system enhances productivity and sustainability in crop production systems, especially when legumes are included in rotations with maize or sorghum (Keeler et al., 2009). Schwember (2020) highlighted that cultivating grain legumes as alternative crops in rotation allows for soil recovery and diversifies production. Through crop rotation, soil biodiversity is enhanced, soil fertility is improved, and incidences of pests and diseases are reduced (Espinoza et al., 2012). Legumes as part of crop rotations also improve soil structure, soil permeability, microbial activity, water holding capacity, and

organic matter content, reducing reliance on chemical fertilizers and increasing biological diversity, thus boosting crop yields and the sustainability of production systems. Knight (2012) indicated that nitrogen fixation and soil microbial functions are influenced by the frequency with which legumes are rotated. However, Brouwer (2006) noted a trend toward specialization in agricultural production systems, which has led to a gradual reduction in diversity and a degradation of biological functions, such as replacing ecological pest management with pesticides or biological nitrogen fixation with mineral fertilizers (Zander et al., 2021). Farmers are encouraged to rotate different species of crops with varying life cycles to maximize pest and disease management while optimizing nutrient availability (Cook, 2013; Garrison et al., 2014; Reckling et al., 2016).

Mixed Cropping

Mixed cropping involves planting a combination of legumes with cereals or tuber crops, a practice particularly common in marginal agroecological environments. This method utilizes growth factors complementarily, including soil nutrients, light, and water, optimizing the use of available resources (Nigli et al., 2008; Dore et al., 2011). It not only reduces the prevalence of pests, diseases, and soil erosion but also enhances biomass and overall yields (Staniak et al., 2014). Adjustments can be made to this cropping system to adapt to variables such as the onset of the rainy season or variations in soil fertility across fields (Weltzien et al., 2017). Importantly, legume-cereal mixtures require lower nitrogen fertilizer inputs than sole cereal crops, resulting in higher protein content in cereal seeds. Research by Zarea et al. (2008) found that mixed cropping of forage legumes enhances the bacterial community in the soil, increasing the presence of free-living N₂-fixing bacteria and *Azotobacter*. Additionally, experiments involving yellow lupine mixed with wheat and oats showed the competitive potential of legumes against cereals (Staniak et al., 2014). The allelopathic interactions within these mixtures can significantly affect plant growth and yield.

Monocropping

Monocropping, the practice of growing a single crop repeatedly on the same land, is often discouraged due to its economic and environmental drawbacks compared to more diverse farming systems involving legumes (Nigli et al., 2008; Dore et al., 2011; Kebede, 2020). This method may lead to issues such as nitrate leaching and increased N₂O emissions, as seen in studies where legume monocrops produced higher cumulative N₂O emissions than fertilized wheat (Senbayram et al., 2016). Due to these limitations, farmers are advised to employ legumes within intercropping, rotational, or mixed cropping systems to mitigate these issues.

Contribution of Grain Legumes to Sustainable Agricultural Production

As global populations rise, pressure on water and land resources intensifies, exacerbated by soil degradation due to global warming, pollution, and loss of soil fertility (Swaroop and Lal, 2018). Legumes offer a viable alternative to support soil stability and enhance soil health when included in crop rotation and intercropping systems. They are integral to integrated soil fertility management, capable of fixing atmospheric nitrogen in

symbiosis with rhizobia bacteria, thus providing significant organic matter that improves the soil's chemical, physical, and biological properties (Sa et al., 2017). Studies have shown that including legumes as intercrops, particularly in maize farms, not only improves productivity but also maintains nutrient availability (Choudhary and Choudhury, 2018). Chimonyo et al. (2019) reported increased maize productivity within a legume-maize cropping system, attributed to improved soil water holding capacity and fertility. This enhancement in soil quality due to legume integration into cropping systems underlines their role in improving soil resilience to erosion and other degradative processes (Kintl et al., 2015).

Table 3: Summary of Legume-Based Practices and Their Impact on Soil Health.

Legume-based system and use of residue	Soil health parameter evaluated	References
Maize-legume rotation	Total N, avail P, exchange K, Mg	Uzoh et al. (2019)
Legume residue	Soil organic carbon, total N, exchangeable Ca and Mg	Kolawole (2013)
Legume intercrop	Soil organic carbon, improving chemical, biological, and physical soil environment, reducing pest damage	Hu et al. (2021)
Legume cover crops	Conservation, SOC and nitrogen stocks, BNF, reduction in nitrous oxide emission	Nees et al. (2010)
Legume residue	Physical properties of soil, i.e., structure, texture, density, stability, porosity	Jena et al. (2022)

Enhancing Ecosystems through Legume Integration in Farming Systems

Incorporating legumes into farming systems significantly strengthens ecosystems by boosting beneficial soil microbes and enhancing soil biodiversity (Meena et al., 2014). These plants facilitate mineralization, offering protection against pests and diseases, and reducing soil erosion (Lal, 2013). Legumes possess extensive root systems and secrete root exudates that greatly improve soil nutrient dynamics, structure, and overall quality (Sugiyama and Yazaki, 2012). They play a crucial role in recycling essential nutrients like nitrogen, phosphorus, and carbon. Various legumes, such as lupin (*Lupinus angustifolius* L.), vetches (*Vicia sativa* L.), velvet bean (*Mucuna pruriens* Bak.), fenugreek (*Trigonella foenum-graecum* L.), and clovers (*Trifolium* sp.), along with *Crotalaria spectabilis* and *Sesbania rostrata*, are utilized as green manure. When used in this capacity, these plants enhance soil organic matter and nutrient availability, enriching the nutrient base for successive plantings and boosting nitrogen stocks within rotational systems (Hauggaard-Nielsen and Jensen, 2005).

Challenges of Sustainable Productivity and the Role of Legumes

Sustainable productivity poses significant challenges, particularly for developing countries where the misuse of agrochemicals has deteriorated soil health. Inappropriate and

excessive use of fungicides to combat soil-borne diseases negatively impacts microbial composition, soil fertility, and grain production (Shahid et al., 2020). Legume-based crop rotations offer a more environmentally friendly solution by significantly enhancing soil microbial health and supporting nutrient mineralization, thus improving soil functions and overall productivity.

Contribution of Legume Residues to Soil Sustainability

Legume residues are valuable sources of mineral nitrogen for subsequent crops due to their relatively high nitrogen content (Kebede, 2021). Decomposed legume plant parts significantly contribute to below-ground nutrient transfer (Louarn et al., 2015). Serving as both live and dead soil covers, legumes help reduce soil moisture loss, lower evapotranspiration rates, and enhance soil rootability (Moura et al., 2015). Using legume crop residues as an alternative method to improve soil fertility is noted by Sangare et al. (2016), although these residues do not immediately provide nitrogen but contribute to a long-term nitrogen pool in the soil (Formowitz et al., 2009). Studies have shown that incorporating organic residues into the soil leads to increased crop yields (Kouyate et al., 2000; Gachengo et al., 1999; Shafi et al., 2007; Bakht et al., 2009). Research by Moura et al. (2014) demonstrated that high-biomass legume trees significantly increased carbon stocks in the litter and total organic carbon. Moura (2009) emphasized the importance of regularly adding residues to maintain a balance between carbon inputs and decomposition rates. Combining mineral fertilizers with organic residues from legumes is promoted as a method to enhance their efficiency in soil amendment (Kouelo et al., 2013).

[Also add a subheading of constraints of legume production :](#)

[Viz. Panda et al\(2019\) noted the major constraints of lentil production were marketing facilities, resource support, inputs availability and its cost.](#)

[Panda, C. K., Kumar, R., & Meena, L. K. \(2019\). Lentil production economics and constraints: An empirical study in Mokama taal of Bihar. *Lentil Production Economics and Constraints: An Empirical Study in Mokama Taal of Bihar*. 14\(3\), 567-571](#)

Conclusion

The diverse and pivotal role of food legumes in farming systems and diets, particularly in impoverished communities, makes them crucial for achieving developmental goals such as reducing poverty and hunger, improving human health and nutrition, and enhancing ecosystem resilience. It is essential to incorporate legumes into agricultural systems to ensure resilient and sustainable agricultural livelihoods. Understanding the unique characteristics of each legume type and how they function within an agricultural context is key to addressing issues of low crop yields and promoting sustainable production.[Also mention constraints in legume production.](#)

REFERENCES

Abberton, M. (2010). Enhancing the role of legumes: potential and obstacles. *Integrated Crop Management*, 11, 177-187.

Ashoka, P., Meena, R. S., Kumar, S., Yadav G. S., & Layek, J. (2017) Green nanotechnology is a key for eco-friendly agriculture. *Journal of Cleaner Production* 142, 4440-4

Ayadi, F. Y., Rosentrater, K. A., & Muthukumarappan, K. (2012). Alternative protein sources for aquaculture feeds. *Journal of Aquaculture Feed Science and Nutrition*, 4(1), 1-26.

Bakht, J., Shafi, M., Jan, M. T., & Shah Z. (2009). Influence of crop residues management, cropping system and N fertilizer on soil N and C dynamics and sustainable wheat production. *Soil and Tillage Research*, 104, 233-240.

Biswas, B., Scott, P. T., & Gresshoff, P. M. (2011). Tree legumes as feedstock for sustainable biofuel production: Opportunities and challenges. *Journal of Plant Physiology*, 168(16), 1877-1884.

Chimonyo, V. G. P., Snapp, S. S., & Chikowo, R. (2019). Grain legumes increase yield stability in maize based cropping systems. *Crop Science*, 59(3), 1222-1235.

Choudhary, V. K., & Choudhury, B. U. (2018). A staggered maize-legume intercrop arrangement influences yield, weed smothering and nutrient balance in the eastern Himalayan region of India. *Experimental Agriculture*, 54(2), 181-200.

Cook, D., Grum, D. S., Gardner, D. R., Welch, K. D., & Pfister, J. A. (2013). Influence of endophyte genotype on swainsonine concentrations in *Oxytropis sericea*. *Toxicology*, 61, 105-111.

Day, L. (2013). Proteins from land plants-potential resources for human nutrition and food security. *Trends in Food Science and Technology*, 32(1), 25-42.

Dhakal, Y., Meena, R. S., & Kumar, S. (2016). Effect of INM on nodulation, yield, quality and available nutrient status in soil after harvest of green gram. *Legume Research*, 39(4), 590-594.

Doré, T., Makowski, D., Malézieux, E., Munier-Jolain, N., Tchamitchian M., &

Tittonell, P. (2011). Facing up to the paradigm of ecological intensification in agronomy: Revisiting methods, concepts and knowledge. *European Journal of Agronomy*, 34(4), 197-210.

Energy efficiency and renewable energy. 2013. Energy 101: feedstocks for biofuels and more. <https://www.energy.gov/eere/videos/energy-101-feedstocksbiofuels-and-more>

Espinoza, S., Ovalle, C., Zagal, E., Matus, I., Tay, J., Peoples, M. B., and del Pozo, A. (2012). Contribution of legumes to wheat productivity in Mediterranean environments of Central Chile. *Field Crops Research*, 133, 150-159.

- Ferguson, B. J., Mens, C., Hastwell, A. H., Zhang, M., Su, H., Jones, C. H., Gresshoff, P. M. (2019). Legume nodulation: the host controls the party. *Plant, cell & Environment*, 42(1), 41-51.
- Ferguson, B., Lin, M. H., Gresshoff, P. M. (2013). Regulation of legume nodulation by acidic growth conditions. *Plant Signaling & Behavior*, 8(3), e23426.
- Formowitz, B., Joergensen, R. G., & Buerkert, A. (2009). Impact of legume versus cereal root residues on biological properties of West African soils. *Plant and Soil*, 325(1), 145-156.
- Friedman M. (1996). Nutritional value of proteins from different food sources: a review. *Journal of Agricultural and Food Chemistry*, 44, 6-21.
- Froidmont, E., & Bartiaux-Thill, N. (2004). Suitability of lupin and pea seeds as a substitute for soybean meal in high-producing dairy cow feed. *Animal Research*, 53(6), 475-487.
- Fustec, J., Lesuffleur, F., Mahieu, S., & Cliquet, J. B. (2010). Nitrogen rhizodeposition of legumes: a review. *Agronomy and Sustainable Development* 30, 57–66
- Gachengo, C. N., Palm, C. A., Jama, B., & Otieno, C. (1999). Tithonia and senna green manures and inorganic fertilizers as phosphorus sources for maize in Western Kenya. *Agroforestry Systems.*, 44, 21-36.
- Garrison, A. J., Miller, A. D., Ryan, M. R., Roxburgh, S. H., & Shea, K. (2014). Stacked crop rotations exploit weed–weed competition for sustainable weed management. *Weed Science*, 62, 166–176.
- Gomes, A. M., & Vasconcelos, M. W. (2014). “The legume grains: when tradition goes hand in hand with nutrition,” in ISEKI Food Series volume 10 Traditional Foods; General and Consumer Aspects, eds K. Kristbergsson and J. Oliveira (Boston, MA: Springer).
- Gregorich, E. G., Rochette, P., VandenBygaart, A. J., & Angers, D. (2005). Greenhouse gas contributions of agricultural soils and potential mitigation practices in Eastern Canada. *Soil Tillage Research*, 81, 53–72
- Guan, X. K., Turner, N. C., Song, L., Gu, Y. J., Wang, T. C., & Li, F. M. (2016). Soil carbon sequestration by three perennial legume pastures is greater in deeper soil layers than in the surface soil. *Biogeosciences*, 13(2), 527-534.
- Harland, J. I., & Haffner, T. A. (2008). Systematic review, meta-analysis and regression of randomised controlled trials reporting an association between an intake of circa 25 g soya protein per day and blood cholesterol. *Atherosclerosis*, 200(1), 13-27.
- Hauggaard-Nielsen H, Jensen E. S. (2005). Facilitative root interactions in intercrops. *Plant Soil*, 274, 237–250
- Howard, J. B., & Rees, D. C. (1996). Structural basis of biological nitrogen fixation. *Chemical reviews*, 96(7), 2965-2982.

- Howieson, J. G., Yates, R. J., Foster, K. J., Real, D., Besier, R. B. (2008). Prospects for the future use of legumes. In *Nitrogen-fixing leguminous symbioses* (pp. 363-394). Springer, Dordrecht.
- Hu, L., Huang, R., Deng, H., Li, K., Peng, J., Zhou, L., & Ou, H. (2021). Effects of different intercropping methods on soil organic carbon and aggregate stability in sugarcane field. *Polish Journal of Environmental Studies*, 31(4), 3587-3596.
- Inal A, Gunes A, Zhang F, Cacak I. (2007). Peanut/maize inter-cropping induced changes in rhizosphere and nutrient concentrations in shoots. *Plant Physiology and Biochemistry*, 45.
- Jena, J., Maitra, S., Hossain, A., Pramanick, B., Gitari, H. I., Praharaj, S., Jatav, H. S. (2022). Role of legumes in cropping system for soil ecosystem improvement. *Ecosystem services: types, management and benefits*. Nova Science Publishers, Inc, New York.
- Jensen, E. S., Peoples, M. B., Boddey, R. M., Gresshoff, P. M., Hauggaard-Nielsen, H., JR Alves, B., & Morrison, M. J. (2012). Legumes for mitigation of climate change and the provision of feedstock for biofuels and biorefineries. A review. *Agronomy for Sustainable Development*, 32(2), 329-364.
- Kebede, E. (2020). Grain legumes production and productivity in Ethiopian smallholder agricultural system, contribution to livelihoods and the way forward. *Cogent Food & Agriculture*, 6(1), 1722353.
- Khorasani, G. R., Okine, E. K., Corbett, R. R., & Kennelly, J. J. (2001). Nutritive value of peas for lactating dairy cattle. *Canadian Journal of Animal Science*, 81(4), 541-551.
- Kintl, A., Elbl, J., Záhora, J., Kynický, J., Brtnický, M., & Mikajlo, I. (2015). Evaluation of grain yield in mixed legume-cereal cropping systems. *Ad Alta: Journal of Interdisciplinary Research*, 5(1), 96-98
- Kolawole, G. O. (2013). Effects of leguminous plant residues and NPK fertilizer application on the performance of yam (*Dioscorea rotundata* 'cv'ewuru) in south-western Nigeria. *Archives of Agronomy and Soil Science*, 59(3), 423-434.
- Kouelo, F. A., Hounnandan, P., & Gerd, D. (2013). Contribution of seven legumes residues incorporated into soil and NP fertilizer to maize yield, nitrogen use efficiency and harvest index in degraded soil in the center of Benin. *International Journal of Biological and Chemical Sciences*, 7(6), 2468-2489.
- Kouyate, Z., Franzluebbers, K., Juo, A. S. R., & Hossner L. (2000). Tillage, crop residues, legume rotation, and green manure effects on sorghum and millet yields in the semiarid tropics of Mali. *Plant and Soil*, 225, 141-151.
- Książak, J., & Staniak, M. (2011). Effect of root excretions from spring cereal seedlings on legume seeds germination. *Journal of Food, Agriculture & Environment*, 9(3), 4.

- Kumar, S., Meena, R. S., Lal, R., Yadav, G. S., Mitran, T., Meena, B. L., EL-Sabagh, A. (2018). Role of legumes in soil carbon sequestration. In *Legumes for soil health and sustainable management* (pp. 109-138). Springer, Singapore.
- Lal, R. (2013) Intensive agriculture and the soil carbon pool. *Journal of Crop Improvement* 27:735–751.
- Layek, J., Das, A., Mitran, T., Nath, C., Meena, R. S., Yadav, G. S., & Lal, R. (2018). Cereal+ legume intercropping: An option for improving productivity and sustaining soil health. In *Legumes for soil health and sustainable management* (pp. 347-386). Springer, Singapore.
- Luke, (2016). Feed Tables and Nutrient Requirements [online]. Natural Resources Institute Finland, Jokioinen. <http://www.mtt.fi/feedtables>. Accessed on 3rd December 2022
- Mandal MK, Banerjee M, Banerjee H, Alipatra A, Malik G. C. (2014). Productivity of maize (*Zea mays*) based intercropping system during kharif season under red and lateritic tract of West Bengal. *Bioscan* 9(1):31–35
- Martin Körschens (2002) Importance of soil organic matter (SOM) for biomass production and environment (a review), *Archives of Agronomy and Soil Science*, 48:2, 89-94
- Mashungwa, G.N., Moroke, T.S., Kgosiesele, E. and Kashe, K. (2019). Grain legume production and their potential for sustainable agriculture in Botswana between 2008 and 2015 – A review. *Botswana Journal of Agriculture and Applied Sciences* 13 (Issue 1 – Special): 80–90.
- McIntyre, B., Gold, C., Kashaija, I., Ssali, H., Night, G., & Bwamiki, D. (2001). Effects of legume intercrops on soil-borne pests, biomass, nutrients and soil water in banana. *Biology and Fertility of Soils*, 34(5), 342-348.
- Meena, V. S, Maurya, B. R., Meena, R. S., Meena, S. K, Singh, N. P., & Malik, V. K. (2014). Microbial dynamics as influenced by concentrated manure and inorganic fertilizer in alluvium soil of Varanasi, India. *African Journal of Microbiology Research* 8(1), 257–263
- Moura, E. G., das, C. F., Aguiar, A., Piedade, A. R., & Rousseau, G. X. (2015). Contribution of legume tree residues and macrofauna to the improvement of abiotic soil properties in the eastern Amazon. *Applied Soil Ecology*, 86, 91-99.
- Nees, B., Anderberg, S., & Olsson, L. (2010). Structuring problems in sustainability science: the multi-level DPSIR framework. *Geoforum* 41(3), 479–488.
- Nigli U, Slab A, Schmid O, Halberg N, Schlüter M. 2008. Vision for an Organic Food and Farming Research Agenda to 2025. Report IFOAM EU Group and FiBL 2008.
- Okumu, O. O. (2018). Effect of lablab (*Lablab purpureus* L.) green manure on population of pathogenic and non-pathogenic soil microorganisms and bean (*Phaseolus vulgaris* L.) crop establishment (Doctoral dissertation, University of Nairobi).

- Okumu, O. O., Muthoni, J., Ojiem, J., Narla, R., & Nderitu, J. (2018). Effect of lablab green manure on population of soil microorganisms and establishment of common bean (*Phaseolus vulgaris* L.).
- Okumu, O., Muthoni, J., Narla, R., Nderitu, J., Lauren, J., & Ojiem, J. (2017, April). Evaluation of Common Bean Production Systems and Fertilizer use in Nandi South. In Innovation Research Symposium 2017. University of Nairobi
- Otieno, H. M. O., Chemining'wa, G. N., Zingore, S., Gachene, C. K. (2020). Tillage method and residual N, P, K, Zn, B, Mg, Ca, and S nutrients effect on growth and yield of dry bean grown after the harvest of maize. *Turkish Journal of Agriculture-Food Science and Technology*, 8(1), 18-26.
- Otieno, H. M., Chemining'wa, G. N., Zingore, S. (2018). Effect of Farmyard Manure, Lime and Inorganic Fertilizer Applications on Soil pH, Nutrients Uptake, Growth and Nodulation of Soybean in Acid Soils of Western Kenya. *Journal of Agricultural Science*, 10(4).
- Peterson, E., Issac, D. J., Luce, C. H., & Rieman, B. E. (2010). Effects of climate change and wildlife on-stream temperatures and Salmonin thermal habitat in a mountain river network. *Ecological Applications*, 20, Iss. 5, 1350-71
- Reckling, M., Hecker, J.-M., Bergkvist, G., Watson, C.A., Zander, P., Schläfke, N., Stoddard, F.L., Eory, V., Topp, C.F.E., Maire, J. and Bachinger, J. (2016a) A cropping assessment framework – evaluating effects of introducing legumes into crop rotations. *European Journal of Agronomy* 76, 186–197
- Semahegn, Z. (2022). Intercropping of cereal with legume Crops. *International Journal of Research in Agronomy*, 5(1), 26-31
- Senbayram, M., Wenthe, C., Lingner, A., Isselstein, J., Steinmann, H., Kaya, C., & Köbke, S. (2016). Legume-based mixed intercropping systems may lower agricultural born N₂O emissions. *Energy Sustain Society*, 6, 2.
- Shafi, M., Bakht, J., Jan, M. T., & Shah, Z. (2007). Soil C and N dynamics and maize (*Zea mays* L.) yield as affected by cropping systems and residue management in Northwestern Pakistan. *Soil and Tillage Research*, 94, 520-529.
- Shafique, A., Rehman, S., Khan, A., & Kazi, A. G. (2014). Improvement of legume crop production under environmental stresses through biotechnological intervention. In *Emerging technologies and management of crop stress tolerance* (pp. 1-22). Academic Press.
- Sherasia, P. L., Garg, M. R., & Bhandari, B. M. (2018). Pulses and their by-products as animal feed. United Nations.
- Sirtori, C. R., Galli, C., Anderson, J. W., & Arnoldi, A. (2009). Nutritional and nutraceutical approaches to dyslipidemia and atherosclerosis prevention: Focus on dietary proteins. *Atherosclerosis*, 203 (1), 8-17.

- Sirtori, C. R., Triolo, M., Bosisio, R., Bondioli, A., Calabresi, L., De Vergori, V., & Arnoldi, A. (2012). Hypocholesterolaemic effects of lupin protein and pea protein/fibre combinations in moderately hypercholesterolaemic individuals. *British Journal of Nutrition*, 107(8), 1176-1183.
- Stagnari, F., Maggio, A., Galieni, A., & Pisante, M. (2017). Multiple benefits of legumes for agriculture sustainability: an overview. *Chemical and Biological Technologies in Agriculture*, 4(1), 1-13.
- Staniak, M., Księżak, J., & Bojarszczuk, J. (2014). Mixtures of legumes with cereals as a source of feed for animals. *Organic agriculture towards sustainability*, 6, 123-145.
- Sugiyama, A., & Yazaki, K. (2012). Root exudates of legume plants and their involvement in interactions with soil microbes. In *Secretions and exudates in biological systems* (pp. 27-48). Springer, Berlin, Heidelberg.
- Sumner, D. R. (2018). Crop rotation and plant productivity. In *CRC handbook of agricultural productivity* (pp. 273-314). CRC Press.
- Swaroop, R. M., & Lal, R. (2018). Legumes and sustainable use of soils. In *Legumes for soil health and sustainable management* (pp. 1-31). Springer, Singapore.
- Świątkiewicz, M., Olszewska, A., Grela, E. R., Tyra, M. (2021). The effect of replacement of soybean meal with corn dried distillers grains with solubles (Cddgs) and differentiation of dietary fat sources on pig meat quality and fatty acid profile. *Animals*, 11(5), 1277.
- Tufarelli, V. R., Khan, U., & Laudadio, V., 2012. Evaluating the suitability of field beans as a substitute for soybean meal in early-lactating dairy cow: production and metabolic responses. *Animal Science Journal*, 83, 136–140.
- Uzoh, I. M., Igwe, C. A., Okebalama, C. B., & Babalola, O. O. (2019). Legume-maize rotation effect on maize productivity and soil fertility parameters under selected agronomic practices in a sandy loam soil. *Scientific Reports*, 9(1), 1-9.
- Van der Pol, M., Hristov, A.N., Zaman, S., & Delano, N., (2008). Peas can replace soybean meal and corn grain in dairy cow diets. *Journal of Dairy Science*, 91, 698–703.
- Vanlauwe, B., Hungria, M., Kanampiu, F., & Giller, K. E. (2019). The role of legumes in the sustainable intensification of African smallholder agriculture: Lessons learnt and challenges for the future. *Agriculture, Ecosystems and Environment*, 284, 106583.
- Vasconcelos, M. W., Grusak, M. A., Pinto, E., Gomes, A., Ferreira, H., Balázs, B., ... & Iannetta, P. (2020). The biology of legumes and their agronomic, economic, and social impact. In *The Plant Family Fabaceae* (pp. 3-25). Springer, Singapore.
- Watson, C. A., Reckling, M., Preissel, S., Bachinger, J., Bergkvist, G., Kuhlman, T., & Stoddard, F. L. (2017). Grain legume production and use in European agricultural systems. *Advances in Agronomy*, 144, 235-303.

Weltzien, E., & Christinck, A. (2017). Participatory breeding: developing improved and relevant crop varieties with farmers. In *Agricultural Systems* (pp. 259-301). Academic Press.

White, C. L., Staines, V. E., & Staines, M. V. H. (2007). A review of the nutritional value of lupins for dairy cows. *Australian Journal of Agricultural Research*, 58, 185–202.

Yu, Y., Stomph, T. J., Makowski, D., Van der Werf, W. 2015. Temporal niche differentiation increases the land equivalent ratio of annual intercrops: A meta-analysis. *Field Crops Research*, 184, 133-144.

Yuvaraj, M., Pandiyan, M., & Gayathri, P. (2020). Role of legumes in improving soil fertility status. *Legume Crops-Prospects, Production and Uses*.

Zander, P., Amjath-Babu, T. S., Preissel, S., Reckling, M., Bues, A., Schläfke, N., Watson, C. (2016). Grain legume decline and potential recovery in European agriculture: a review. *Agronomy for Sustainable Development*, 36(2), 1-20.

Zarea, M. J., Ghalavand, A., & Jamshidi, E. (2008). Role of forage legumes mixed cropping on biomass yield and bacterial community composition.

Zhao, J., Yang, Y., Zhang, K., Jeong, J., Zeng, Z., & Zang, H. (2020). Does crop rotation yield more in China? A meta-analysis. *Field Crops Research*, 245, 107659.

UNDER PEER REVIEW