

TENDON INJURIES OF THE HAND; A REVIEW OF CASES AT A SUBURBAN TERTIARY HOSPITAL, NIGERIA.

ABSTRACT

Background: Tendon injuries of the hand are one of the commonest presentations of hand injuries which itself bears a social and economic burden to the society as the most commonly affected are the active and working age groups. Most of these injuries will require surgical intervention to restore aesthetics, structure and function.

Objective: With few reports on tendon injuries to the hand, this study seeks to assess the pattern of these injuries, etiology and reappraise the preventive modalities with the aim of reducing incidence in our immediate environment.

Method: A retrospective review of patients who presented and were managed surgically for tendon injuries of the hand with or without injury to other structures of the hand between January 2017 and December 2021. The operation registers as well as case notes of patients were the sources of the information. Information obtained included patients' biodata, aetiology and pattern of injury, hand involved (whether right or left), collateral injuries as well as surgery offered. The results were analysed using SPSS version 20 and descriptive statistics used to represent frequency distribution.

Results: In the five years under review, a total of 27 patients were managed for hand injuries during the period of study, however, only 21 cases which involved tendon injuries were analyzed making it 77.7% occurrence in hand injuries. There were 17 males [84.2%] and 4 females [15.7%] giving a male to female ratio of 4.25:1. The mean age was 30.3 ± 17.10 years and the highest incidence of tendon injury; 9 (42.8%) occurred within the age range of 21-30 years of age. Industrial accident due to grinding machines, 8(38%) was the commonest cause of tendon associated hand injuries, followed by matchet cut injury to the hand following assault 6(28.6%). The overall most common tendon injury was flexor tendon injury (71.4%) with the right hand being most commonly affected; 16(76.2%). All patients had tendon repair done alongside repair of other associated injuries of the hand.

Conclusion: Tendon injuries of the hand is a common occurrence following injury to the hand and seen among the economically active groups with crippling implications. In addition to its proper management by a skilled surgeon, preventive efforts must be implemented to prevent its occurrence as majority of its etiology is preventable.

Key words: *Tendon, Injury, Hand, Suburban, Nigeria*

1. INTRODUCTION

The hand, as the human executing organ, is in the center of daily life activities in professions and sports. In this outstanding position, the hand is always exposed to injuries and overuse¹. Tendon injuries are the second most common injuries of the hand². Tendons are strong, dense, connective tissue structures that attach muscles to bones. Injury patterns are differentiated into open or closed, sharp or blunt, traumatic or degenerative lesions. Most injuries are open injuries to the flexor or extensor tendons, but less frequent injuries, e.g., damage to the functional system tendon sheath and pulley or dull avulsions². Proper

management of hand tendon injuries requires a good knowledge of the anatomy and physiology of the hand and forearm. Hand surgeons^{3,4,5} agree that the goal of treatment of hand injury is to restore an aesthetically pleasing, painless, tactile, mobile, stable finger that can sense pain, temperature, pressure, stereogenesis and fine touch.

One-third of all accidents, and about 20% of all lacerated wounds occur in the hand^{4,5}. Hand injuries are a leading cause of time lost from work and workers compensation claims has been documented by most workers. The United States Bureau of labor statistics reports that hand injuries are the second most common injury resulting in days away from work^{5,6}. In a study conducted at the Lagos university teaching hospital Lagos, Nigeria⁷, hand injuries are among the common emergency hand problems that present in the accident and emergency department which demand immediate attention and treatment. Management of hand injury is a teamwork, the team must also involve the nurses, physiotherapist, the patients, their relatives, the hospital administrators with the hand surgeons⁸. Although orthopedic surgeons are involved in the management of hand injuries, however much are managed by plastic surgeons especially with respect to tendon injuries.

In a study carried out in two tertiary hospitals in Delta State, Nigeria⁹ between January 2013 and December 2015. The commonest cause of hand injury was due to machete/ knife cuts and stab wounds, followed by road traffic accidents, burn injuries and machinery accidents respectively. On the types of hand injuries sustained, deep lacerations were the commonest and comprised; 15 flexor tendon lacerations, 14 extensor tendon lacerations, 4 combined tendon and nerve lacerations and 1 isolated nerve laceration. The right hand was the most frequently injured hand. Majority of the hand injured patients had tendon repair for both flexor and extensor tendons⁹. In another study carried out in a suburban hospital in Bayelsa,¹⁰ road traffic accidents (RTA) remains the major cause of hand injury affecting more males and manual workers in the productive age group.

Edo state is located in the south-south zone of Nigeria and in it are two (2) Teaching Hospitals, with Irrua Specialist Teaching Hospital located in the suburb of the state. The hospital (ISTH) has a well established Plastic and Reconstructive surgery unit. The unit manages cases of hand injuries that present to or referred to the hospital. Looking through the unit records, many patients have been managed for different tendon injuries of the hand. This study is conducted to assess the pattern of these injuries, etiology and reappraise the preventive modalities with the aim of reducing incidence of these injuries in our immediate environment.

2. MATERIALS AND METHODS

This study was conducted at Irrua Specialist Teaching Hospital (ISTH), Edo state. "Irrua Specialist Teaching Hospital (formerly Otibhor Okae Teaching Hospital) was established by decree 92 of 1993 to provide tertiary health services to people of Edo State and beyond¹¹. This hospital is a 350-bed tertiary facility servicing the central and northern part of Edo State, Nigeria.

This was a hospital-based five (5) year retrospective review of patients who presented and were managed surgically for tendon injuries to the hand with or without injury to other structures of the hand between January 2017 and December 2021. Patients that were managed conservatively and poorly documented for were excluded. The operation registers as well as case notes of patients were the sources of the information. Information obtained included patients' biodata, aetiology and pattern of injury, hand involved (whether right or left), collateral injuries as well as surgery offered. The results were analysed using SPSS version 20. Descriptive statistics were used to represent frequency distribution. Ethical approval for this

study was obtained from the Health Research Ethics Committee of Irrua Specialist Teaching Hospital, Irrua, Edo state.

3. RESULTS

In the five years under review, a total of 27 patients were managed for hand injuries during the period of study, however, only 21 cases which involved tendon injuries were analyzed making it 77.7% occurrence in hand injuries. There were 17 males [81%] and 4 females [19%] females giving a male to female ratio of 4.25:1. Their age ranged from 11 to 60 years with a mean age of 30.3 ± 17.10 years. Most of the patients were machinery operators 8 (38%), followed by trading 6 (28.6%) and commercial cyclists 3 (14.3%).

Table 1. Age and sex distribution of patients

Variable	Frequency (%)
Age (years)	N = 21
< 10	0 (0)
11 - 20	3 (14.3)
21 - 30	9 (42.8)
31 - 40	6 (28.6)
41 - 50	2 (9.5)
51 - 60	1 (4.8)
Mean \pm SD	= 30.3 \pm 17.10 years
Gender	
Male	17 (81)
Female	4 (19)

The highest incidence, 9 (42.8%) of hand injuries occurred within the age range of 21-30 years of age. Majority 9 (42.9%) of the patients were students, followed by traders 4(19%). Industrial accident due to grinding machines, 8 (38%) was the commonest cause of tendon associated hand injuries, followed by matchet cut injury to the hand 6 (28.6%) following assault. Gun-shot injury and home accidents were the least cause of injuries. Other associated hand injuries sustained by our patients, deep lacerations were the commonest (42.9%), followed by nerve injuries (23.8%) and crush injury to the hand (14.3%). Tendon injuries sustained includes 15 flexor tendon injury (71.4%), and 6 extensor tendon injury (28.6%), leaving the most commonly injured to be the flexor tendons. The right hand was the most commonly affected 16 (76.2%). All patients had tendon repair done alongside repair of other associated injuries of the hand.

Table 2. Occupation of patients

Variable	Frequency (%)
Occupation	N = 21
Machinery operator	8 (38)
Trader	6 (28.6)
Commercial cyclist	3 (14.3)

Student	2 (9.5)
Unemployed (housewife)	1 (4.8)
Others (Artisan)	1 (4.8)

Table 3. Mechanism of injury/Aetiology

Mechanism	Frequency (%) N = 21
Industrial accident (grinding machines)	8 (38)
Matchet cut (following assault)	6 (28.6)
Road traffic accident (RTA)	5 (23.8)
Gun-shot injury	1 (4.8)
Home accident	1 (4.8)

Table 4. Pattern of tendon injury

Variable	Frequency (%)
Injury	N = 21
Flexor tendon in forearm	10 (47.6)
Extensor tendon in the forearm	6 (28.6)
Zone I flexor tendon	3 (14.3)
Zone II flexor tendon	2 (9.5)

Table 5. Associated hand injuries

Injury	Frequency (%) N = 21
Laceration	9 (42.9)
Nerve injury	5 (23.8)
Crush injury	3 (14.3)
Amputation	2 (9.5)
Vascular injury	2 (9.5)




Figure 1; ZONES OF THE FLEXOR TENDON OF THE HAND
 (figure with permission of Bal et al¹²).

Zone I - Contains flexor digitorum profundus only

Zone II "No Man's Land" - Extends from insertion of flexor digitorum superficialis (FDS) to proximal edge of A1 pulley

Zone III - Extends from the proximal end of A1 to the distal edge of of carpal tunnel

Zone IV - Lies within the carpal tunnel

Zone V - Lies proximal to the carpal tunnel




Figure 2; PULLEY AND FLEXOR TENDON ANATOMY
 (figure with permission of Dr Jared Vagy DPT: Will Anglin).

4. DISCUSSION

Hand injuries though not a common cause of mortality can be a source of severe pain, cosmetic embarrassment, loss of productivity and even depression to the owner. Tendon injuries of the hand are serious injuries that should be treated by a skilled Plastic surgeon. The objective is to restore as much function as possible¹³. When treatment is delayed, or inadequate, hand injuries may result in crippling complications especially when they become infected^{4,5}.

The results from our study showed more males sustaining tendon related hand injuries with the peak incidence in the 3rd and 4th decades of life. This is the age of active physical activities and productivity which again portrays the negative effect it will have in the life of the individual, family and society at large. In a study conducted in the United states by Chah SS et al¹⁴ and Laren et al¹⁵ Netherlands and Denmark hand injuries were commoner in males than in females and are more frequent among individuals aged above 18 years. In another study conducted in Nigeria by Inyang U C et al¹⁶ hand injuries occurred more also in males than in females in the ratio of 8:1, and the peak age incidence of these injuries was between 20-40 years. This can be explained by most manual work being carried out by men in the society and also the predilection towards employment of males for high risk manual jobs.

The commonest cause of tendon related hand injuries recorded from our study was injuries occupational accidents from grinding machines followed by machete cuts following assault. This contrasts a similar study in Delta state, where the leading cause of hand injuries resulted from assaults and fights involving use of machetes and knives followed closely by hand injuries from road traffic accidents (RTA). This disparity could be explained by the prevalence of low scale industries such as cassava processors, bakeries and local grinders in our study area being a semi-urban center. Most patients were right handed which could explain the reason why the right hand was most commonly injured. This could mean loss of productivity especially the machinery operators, farmers and others who relied a lot on their hands to earn a living, and inevitably affect the economic potential of the individual, family and country at large. This can be seen in a study in Mulago, Uganda¹⁷ where half of the participants were unable to fully use the injured hand for work for at least a month following hand injury. In this study, deep lacerations were the commonest injuries sustained and so a lot of tendon injuries, both flexors and extensors, as well as nerve and bone injuries were sustained. Consequently, the commonest surgical procedures performed were tendon repairs (flexors more than extensors)¹⁶.

In line with our findings, most of the injuries are completely preventable (occupational accidents, machete cuts and RTA's rank high in the aetiology of tendon related hand injuries). It has been noted by David S. S. and Goel K., that "hand injuries are as preventable as the other bodily traumas"¹⁸. It was the belief of Makobore et al, that targeted campaigns to sensitize workplaces and reduce accidents on the road may contribute to prevention of hand injuries¹⁹ this also holds true from our findings in this study. Furthermore, gangsterism and violence in communities can be eradicated or brought to the barest minimum, at least, by appropriate legislation and also law enforcement by the relevant security agencies. There should also be measures applied to road users and industrial organizations and their workers, as well as education of the populace.

5. CONCLUSION

Tendon injuries of the hand are common following injury to the hand and seen among the economically active groups with economic, cosmetic, psychological and social implications. In addition to management

by a skilled surgeon, measures should be put in place to prevent its occurrence as most of its etiology is preventable.

REFERENCES

1. Voigt C. [Tendon injuries of the hand]. *Chirurg* 2002; 73: 744-64; quiz 765-7
2. Schöffl V, Heid A, Küpper T. Tendon injuries of the hand. *World J Orthop* 2012; 3(6): 62-69 Available from: URL: <http://www.wjgnet.com/2218-5836/full/v3/i6/62.htm> DOI: <http://dx.doi.org/10.5312/wjo.v3.i6.62>
3. Chan SS, Rochette LM, Smith GA. Epidemiology of pediatric hand injuries presenting to the United States emergency department 1990 to 2009. *J Trauma Acute Care Surg.* 2012;72(6):1688-1694.
4. Adeyemi-Doro. Operated hand injuries and hand infections in an urban emergency service. *Nigerian Medical Practitioner.* 1988;16(5-6):171-173.
5. Adeyemi-Doro HO. Management of hand infections. *Nigerian Medical Practitioner.* 1981;1(4):5-10.
6. Bureau of Labour statistics, U.S department of Labour: Non-fatal occupational injuries and illness requiring days away from work; 2011. Available:http://www.bls.gov/news_release/archives/osh211082.pdf (March 3 2014)
7. Adeyemi-Doro HO. Manual of emergency surgery, edited by Adeyemi-Doro H O. University of Lagos Press. 1991;224-251.
8. O.M. Oluwatosin et al, Pattern And Management Of Hand Injuries In Ibadan, Nigeria: A FiveYearReview; *The Tropical' Journal of Health Sciences* Vol. 13(2005).
9. Otene C. I., Ikubor J. E., Idiakhwa O. O., Otene C. O; The Burden of Hand Injuries in Delta State, Nigeria; *Int. J of Forensic Med Invest* 2016; 2(1)20-24
10. Benjamin and Ejike; Hand Injuries in a Suburban Hospital Bayelsa State Nigeria; *British Journal of Medicine & Medical Research* 20(8): 1-8, 2017; 20(8): 1-8, 2017; Article no.BJMMR.30731
11. Irrua Specialist Teaching Hospital (ISTH). Available:<https://www.isth.org.ng/about/>

12. Bal et al Anatomic and Functional Improvements Achieved by Rehabilitation in Zone II and Zone V Flexor Tendon Injuries; American journal of physical medicine & rehabilitation / Association of Academic Physiatrists
13. Robert E. Page: Hand Injuries. Surgery. 1986; 759-764.
14. Chan SS, Rochette LM, Smith GA. Epidemiology of pediatric hand injuries presenting to United States emergency department 1990 to 2009. J Trauma Acute Care Surg. 2012;72(6):1688-1694.
15. Larsen CF, Mulder SS, Johnson AM, Stam C. The epidemiology of hand injury in the Netherland and Denmark. European Journal of Epidemiology. 2004;19:323- 327.
16. Inyang UC, Dim EN, Ofoegbu CKP et al. pattern of hand injury in Ilorin North central Nigeria; A review of 42 cases. Pioneer Medical Journal. 2013;3(5):1-11
17. Makobore P., Galukande M., Kalanzi E., Kijjambu S. C. The Burden of Hand Injuries at a Tertiary Hospital in Sub-Saharan Africa. Emerg Med Int, 2015; ID 838572.
18. David S S, Goel K. Knowledge, attitude, and practice of sugarcane crushers towards hand injury prevention strategies in India. Injury Prevention 2001; 7: 329–330.
19. Makobore P., Galukande M., Kalanzi E., Kijjambu S. C. The Burden of Hand Injuries at a Tertiary Hospital in Sub-Saharan Africa. Emerg Med Int, 2015; ID 838572.

UNDER PEER REVIEW