

A Review on efficacy effect of newer insecticides and biopesticides against brinjal shoot and fruit borer [*Leucinodesorbonalis* (Guenee)]

Abstract:-

Eggplant (*Solanum melongena*) is one of the most important vegetable of hot and wet climatic zones. There are 26 insects-pests species and few non insect pests species infesting brinjal shoot and fruit borer (*Leucinodesorbonalis*) major pest considered the main constraint as it damages the crop throughout the year. The loss of yield due to the pest ranges from 70 to 92 percent. This situation has posed a hindrance to the farmers in commercial cultivation of brinjal . After reviewing the research papers, it was found that newer insecticide and biopesticides were significantly superior, such as spinosad 45SC, followed by Indoxacarb 14.5SC and Emmaectin benzoate 0.5SG.

Keyword:

INTRODUCTION

Brinjal cultivation in India faces significant challenges from a variety of pests at different stages of growth—severely by number of pests, shoot and fruit borer (*Leucinodesorbonalis*), epilachna beetle (*Epilachnavigintioctopunctata*Fab.), jassid (*Amrascabiguttulabiguttula*Shida), aphid, thrips and white flies (*Bemiciatabaci*Gennadius). Among these pests, the shoot and fruit borer, (*Leucinodesorbonalis*G.), is widely regarded as the most destructive pest affecting brinjal crops in India. The infestation on brinjal can be as high as 75 to 92% Brinjal crop is attacked regularly or Brinjal in India is sporadically attacked by at least 50 insect pests, with Aphids, Jassids, Whiteflies, and Shoot and Fruit Borers being categorized as major pests that regularly afflict the crop.(PoojaandKumar,2022) Among the insects pest the most important *Leucinodesorbonalis*Guenee(Lepidoptera: Pyralidae). It stands out as the most destructive and consistently recurring pest, inflicting significant crop losses.(Umaphy *et al.*, 1991) and furthermore liable for weakening of natural product this characteristic ultimately impacts the market value of organic produce (Mathiranjani *et al.*, 2000)and (Das *et al.*,2008). The infestation intensity by the pest can reach levels as high as over 90% (Bhutani *et al.*, 1976; Kallou *et al.*, 1988). Yield losses have been estimated to range from 86% (Naresh *et al.*, 1986) in Haryana to approximately 83 to 91% in Uttar Pradesh (Patel *et al.*, 2021).

DAMAGING PATTERN AND BIOLOGY OF *Leucinodesorbonalis*

At beginning phase of the crop, the female moth lays eggs generally on the lower side of the leaves close to midrib, at the highest point of shoot or some of the time even on

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: Font: Italic

Commented [N1]: Reference

Formatted: Font: Bold

delicate shoots itself (Basu *et al.*, 1968; Singh and Singh, 2003). Drooping, wilting or withering of shoots are typical symptoms of shoot damage during the beginning phase of growth (Srinivasan *et al.*, 1998; Rakibuzzaman *et al.*, 2019). After fruit formation larvae generally enter from the underside of the calyx or bud or fruit (Briere *et al.*, 1999). The holes observed on the fruits are indeed the exit points of the larvae. (Singh and Singh, 2003). Significant harm is noticed each year, influencing antagonistically, the quality and yield of the harvest (Colinet *et al.*, 2015).

It alone results in damage as high as 85-90%, with instances of up to 100% damage also documented. The larvae bore into tender shoots, causing wilting and dead heart. In later stages, they bore into the tender fruits, rendering them unsuitable for human consumption. So far, (*Leucinodesorbonalis*) is considered as a major pest of brinjal as shoot and fruit borer in established crop in main field (Halder *et al.*, 2015)

The flat-oval eggs were laid most frequently at night. The average number of eggs laid per female was 121.5 ± 0.449 and of these 79.24% were viable. There were 6 larval instars. The egg, larval and pupal periods were 6.0 ± 0.058 , 15.38 ± 0.214 and 11.5 ± 1.11 days, resp. The average longevities of males and females were 4.0 ± 0.707 and 7.5 ± 0.267 days, respectively. It's also reported that there will be a reduction in vitamin C levels content to an extent of 68 per cent in the infested fruits. It was reported that the shoot and fruit borer (on shoot) were more prevalent during the vegetative phase of crop. (Yadav *et al.*, 2015).

Damage is caused by caterpillars that are creamy white when young but become light pink as they mature when full grown having length 18-23 mm. Generally larva undergoes 5-6 instars, (Raina and Yadav 2017)

It seems to match the developmental stages of *Leucinodesorbonalis*, the eggplant fruit and shoot borer, quite closely. Let's break down the stages:

Newly hatched larva: Tiny, creamy or dirty white in color with a dark brown or light black head. It has three pairs of thoracic legs and five pairs of pro legs. This is typical of many moth larvae, including *Leucinodesorbonalis*, upon hatching.

2nd instar larva: Resembles the first instar larva but is larger in size and may appear somewhat darker in color. This is also common in many moth larvae as they molt and grow between instars.

3rd instar larva: Longer than the 2nd instar larva and darker in color. The prothoracic legs, the legs near the head, are dark brown in color. This stage represents further growth and development.

4th instar larva: Takes on a more pinkish coloration. This could indicate changes in the larva's diet or metabolism as it progresses through its development.

Commented [N2]: reference

Commented [N3]: reference

5th instar larva: Has a pinkish-brown color with three distinct segments of the thorax and five pairs of well-developed prolegs. This is likely the final instar before pupation. These descriptions align well with the developmental stages typically observed in *Leucinodes orbonalis* larvae. Monitoring these stages can be crucial for implementing control measures effectively and managing infestations in eggplant crops. (Shaukat *et al.*, 2018)

The pupal period of BSFB varies: in summer, it lasts for 7-10 days, while in winter, it extends to 13-15 days pupation (Shaukat *et al.*, 2018). Pupation of this pest takes place generally in soil and also in dried shoots, leaves or plant refuse or debris fallen on the ground. (Butani and Verma 1976) The BSFB moth is white in colour with a blackish-brown spot on the dorsum of the thorax and abdomen. Whitish wings and pinkish brown tinge and are tinged ringed with small hair along the apical and anal margin (Raina and Yadav 2018).

Table 1. Newer insecticides and their mode of actions on insect pests

S.No.	Newer insecticides	Mode of action
1	Chlorantraniliprole	Coragen binds to a specific receptor in muscles known as the ryanodine receptor. When chlorantraniliprole attaches to this receptor, it prompts muscle cells to release calcium, resulting in abnormal muscle function. This results in paralysis and eventual demise of the insect. The ryanodine receptor differs between insects and mammals, with coragen binding much more strongly to the insect receptor (Cordova <i>et al.</i> , 2006 & Sattelle <i>et al.</i> , 2008). It primarily affects insects when ingested (FHHERA, 2019). Additionally, coragen is toxic to insect eggs, larvae, and pupae upon contact (Brugger <i>et al.</i> , 2018 & Krishan <i>et al.</i> , 2021).
2	Flubendiamide	Masaki <i>et al.</i> (2006) showed that FBD stimulates Ca ²⁺ pump activity is hindered by reducing the coupling between RyRs and the pump, leading to a decrease in internal calcium concentration. This particular mode of action of FBD causes various disruptions in muscle function in the target insect, resulting in symptoms of poisoning such as rapid cessation of feeding and contractile paralysis and regurgitation leading to the death of insect.
3	Thiacloprid	Thiacloprid is an insecticide of the neonicotinoid class. Its mechanism of action is akin to other neonicotinoids and entails disrupting the insect's nervous system through stimulating nicotinic acetylcholine receptors (USEPA, 2012 & Schuld and Schmuck, 2000).
4	Carbofuran	It is a systemic insecticide, Carbofuran also has contact activity against pests. It is one of the most toxic pesticides still in use (APDM 2020).
5	Indoxacarb	Insecticidal activity occurs via blockage of the sodium channels in the insect nervous system and the mode of entry is via the stomach and contact routes (USEPA 2000).
6	Emamectin benzoate	When sprayed to foliage, emamectin benzoate penetrates the leaf tissue and forms reservoir within treated leaves, which provides residual activity against pests that feed on foliage and ingest the substance when feeding. The proposed formulation is intended to translocate within the tree's vascular system upon injection (USEPA 2009).

Commented [N4]: ingredients/elements

Commented [N5]: ingredients/elements

Commented [N6]: ingredients/elements

Commented [N7]: ingredients/elements

Commented [N8]: ingredients/elements

Commented [N9]: ingredients/elements

7	Spinosad	The specific mode of action of spinosad is to alter the function of nicotinic and GABA-gated ion channels, causing rapid excitation of the insect's nervous system, resulting in involuntary muscle contractions, tremors, paralysis, and death (Fulton <i>et al.</i> , 2013)
---	----------	---

Commented [N10]: ingredients/elements

Table 2. Bio pesticide and their mode of action on insect pest

S.No.	Bio pesticide	Mode of action
1	Bt	The most approved and widely accepted primary mode of action of the Bt toxin is the lysis of epithelial cells in the midgut of the insect. The toxin acts from the exterior of the cells and enters the plasma membrane, but it does not enter the cytoplasm. (Liu <i>et al.</i> 2018)
2	<i>Beauveria bassiana</i>	Insect death is caused by starvation as the fungus takes over the internal structures of the insect, ultimately resulting in outward penetration of the cuticle and sporulation on the mummified body of the host (Urquiza <i>et al.</i> ,2015).
3	<i>Metarhizium anisopliae</i>	The fungus <i>Metarhizium anisopliae</i> strain F52 infects insects that come in contact with it. Once the fungus spores attach to the outer surface of the insect, they germinate and begin to grow. Upon penetrating the external exoskeleton of the insect, they proliferate rapidly inside, including the insect to die (USEPA 2003).
4	Neem oil	Most active component for repelling and killing pests (USEPA.,2012)

Commented [N11]: full meaning

Efficacy of various insecticides and biopesticides

Studies on chemical control of *L. orbonalis* revealed that, treatments of Emamectin benzoate 5% SG and Coragen 18.5% SC proved to be the most effective. The insecticides viz., Azadirachtin 5% EC and *Bacillus thuringiensis* 5% WP were found least effective against shoot and fruit borer. The remaining treatments viz., Spinosad 45% SC, Lambda cyhalothrin 5% EC, Pyriproxyfen 5% EC + Fenprothrin 15% EC were found moderately effective (Vinayaka *et al.*,2019) Naik and his co-workers study resulted that the results of chemical control trial profenofos @ 0.1% and spinosad @ 0.015% were most effective in reduction of shoot infestation of *L. orbonalis* besides recording higher brinjal fruit yield. Among the 15 treatments tested, profenofos proved to be the most effective, followed by spinosad individually. Their combinations with novaluron and Azadirachtin were highly effective in reducing the population as well as in giving higher yields (Naik *et al.*, 2008).

Results of another trail stated that Emamectin benzoate 5SG @ 50gm/lit was found most effective and showed (8.71%) shoot infestation and per cent fruit infestation (7.22%) followed by Spinosad 45 SC @ 0.02ml/lit (10.13) and (7.69), Cypermethrin 25 EC @ 2ml/lit (11.51)(Shyamrao *et al.*,2018). The treatments, namely chlorantraniliprole 18.5 SC (0.4ml/l), spinosad 45 SC (0.5ml/l), chlorpyrifos 10 SC (2ml/l), indoxacarb 14.5 SC (1ml/l), *Bacillus thuringiensis* (Bt) (2g/l), Azadirachtin 0.03EC (5ml/l), *Metarhizium anisoplae* (2.5g/l), *Beauveria bassiana* (2.5g/l) and chlorpyrifos 20EC (2.5 ml/l) were applied thrice at fifteen days interval starting from initiation of BSFB infestation. Mean shoot infestation was lowest in coragen plots (6.32%) followed by spinosad, chlorpyrifos indoxacarb. Among bio-pesticides, *Beauveria* and *Bt* were identified as effective treatments in reducing shoot infestation. Coragen achieved the lowest fruit infestation (8.25%) and highest marketable fruit yield (250.30q/ha) followed by spinosad and chlorpyrifos.(Tripura *et al.*, 2017). the different insecticidal treatments, by application of Emamectin benzoate 25 WG@0.4gm/lit recorded lowest fruit damage of 6.95% with highest yield of 351.46 qt/ha. However, it was at par with Spinosad 45 SC@0.5 ml/lit with fruit damage of 8.06% and yield of 341.75 q/ha. (Mane and Kumar2020). The bio-insecticides and botanicals tested included: Emamectin benzoate 5 SG @ 75 g a.i/ha, Spinosad 45 SL @ 18 g a.i/ha, NSKE 5%, Karanj seed extract 5%, Onion extract 5%, Garlic extract 5%, Tobacco extract 5%, Cannabis (bhang) leaf extract 5%, Wood ash 10 g/ plant and compared with control. Results showed that Emamectin benzoate 5 SG (75 g a.i/ha) treated plots exhibited the lowest infestation and resulted in higher fruit yield (313.85 q/ha) followed by Spinosad 45 SL (18 g a.i/ha) and NSKE (5%) which gave 300.58 and 284.33 q/ha fruit yield respectively. The least effective treatment was Wood ash (10 g/plant), yielded only 225.14 q/ha healthy fruits. The highest cost-benefit ratio was obtained from Emamectin benzoate 5 SG 75 g a.i/ha (1:21.23) treated plots. Tobacco leaf extract 5% treated plots though ecofriendly but gave least cost-benefit ratio (1:1.27). This study suggests the utilization of bio-insecticides and botanicals as they demonstrate significant efficacy in managing the brinjal shoot and fruit borer infestation(Verma *et al.*, 2021).

(Jat and Shrivastva 2023)revealed that all the treatments were found significant superior over control. The economics pattern of newer insecticides viz., Spinosad 45 SC, followed by Indoxacarb 14.5 SC and Emamectin benzoate 5 SG. It can be concluded that the most effective control of brinjal shoot and fruit borer was achieved with the insecticide Spinosad 45 SC after two sprays at the recommended intervals and doses.

Conclusion

References

Vinayaka KS, Balbir Singh, SS Yadav and Sudhanshu Bala Nayak. Efficacy of Different Insecticide against brinjal fruit borer (*Leucinodesorbonalis*Guenne) and their impact on fruit yield. *Journal of Entomology and Zoology Studies* 2019; 7(3): 63-66

Formatted: Justified

Commented [N12]: follow journal rules of referencing. used recent and update reference. give journal link with DOI number

Formatted: Font: Italic

Ajit Tripura, ML Chatterjee, Rachna Pande and Sandipp Patra. Biorational management of brinjal shoot and fruit borer (*Leucinodesorbonalis* Guenee) in mid hills of Meghalaya. *Journal of Entomology and Zoology Studies* 2017; 5(4): 41-45

Formatted: Font: Italic

Ankur Prakash Verma, Umesh Chandra, Anuj Shakya, Reetesh Pratap Singh, Bhupendra Singh and Khursheed Alam. Efficacy of bio-insecticide and Botanicals against brinjal shoot and fruit borer (*Leucinodesorbonalis* Guenee). *The Pharma Innovation Journal* 2021; 10(12): 1071-1077

Formatted: Font: Italic

PD Mane and Randhir Kumar. Bio-efficacy of new chemicals against shoot and fruit borer (*Leucinodesorbonalis*) of brinjal. *Journal of Entomology and Zoology Studies* 2020; 8(6): 1692-1693

Formatted: Font: Italic

Nawale Jayant Shyamrao, Ashwani Kumar, Patil AA and Narode MK. Efficacy of Certain insecticide and bio-pesticide against brinjal shoot and fruit borer (*Leucinodesorbonalis* Guenee). *Journal of Entomology and Zoology Studies* 2018; 6(5): 292-295

Formatted: Font: Italic

Naik CV, Rao PA, Krishnayya PV, Rao VS. Seasonal Incidence and Management of *Leucinodesorbonalis* Guenee on Brinjal. *Annals of Plant Protection Sciences* 2008;16(2):329-332

Formatted: Font: Italic

Raina J, Yadav GS. Influence of abiotic factors on population dynamics of *Leucinodesorbonalis* Guenee on brinjal in Hissar agro-climatic conditions. *Annals of Plant Protection Sciences*. 2017;25(2):277-280

Formatted: Font: Italic

Jyoti Raina, GS Yadav. BSFB: Bio-ecology and management. *Journal of pharmacognosy and phytochemistry*, 2018;7(4):444-449.

Formatted: Font: Italic

Formatted: Font: Italic

R Srinivasan. Integrated pest management for BSFB in south and Southeast Asia: past, present and future. *Journal of Biopesticides*, 2008, 105-112.

Formatted: Font: Italic

Cordova, D.; Benner, E. A.; Sacher, M. D.; Rauh, J. J.; Sopa, J. S.; Lahm, G. P.; Selby, T. P.; Severson, T. M.; Flexner, L.; Gutteridge, S.; Rhoades, D. F.; Wu, L.; Smith, R. M.; Tao, Y. Anthranilic Diamides: A New Class of Insecticides with a Novel Mode of Action, Ryanodine Receptor Activation. *Pestic Biochem Phys* 2006, 84, 196-214.

Sattelle, D. B.; Cordova, D.; Cheek, T. R. Insect Ryanodine Receptors: Molecular Targets for Novel Pest Control Chemicals. *Invert Neurosci* 2008, 8, 107-119.

Final Human Health and Ecological Risk Assessment for Chlorantraniliprole Rangeland Grasshopper and Mormon Cricket Suppression Applications; US Department of Agriculture, Marketing and Regulatory Programs, Animal and Plant Health Inspection Service, 2019.

Commented [N13]: rewrite it

Brugger, K. E.; Kole, P. G.; Newman, I. C.; Parker, N.; Scholz, B.; Suvagia, P.; Walker, G.; Hammond, T. G. Selectivity of Chlorantraniliprole to Parasitoid Wasps. *Pest Manag Sci* 2010, 66, 1075-1081.

Krishnan, N.; Zhang, Y.; Aust, M. E.; Hellmich, R. L.; Coats, J. R.; Bradbury, S. P. Monarch Butterfly (*Danaus Plexippus*) Life-Stage Risks from Foliar and Seed-Treatment Insecticides. *Envir Tox Chem* 2021, 40 (6), 1761-1777.

Masaki, T., Yasokawa, N., Tohnishi, M., Nishimatsu, T., Tsubata, K., Inoue, K., Hirooka, T., 2006. Flubendiamide, a novel Ca²⁺ channel modulator, reveals evidence for functional cooperation between Ca²⁺ pumps and Ca²⁺ release. *Mol. Pharmacol.* 69, 1733–1739.

Formatted: Font: Italic

Thiacloprid Pesticide Fact Sheet, United States Environmental Protection Agency Archived 2012-06-27.

Schuld M, Schmuck R (2000). "Effects of Thiacloprid, a New Chloronicotinyl Insecticide, On the Egg Parasitoid *Trichogrammacacaeciae*". *Ecotoxicology*. 9 (3): 197–205.

Ravichandra, N.G. (2018). *Agrochemicals in Plant Disease Management*. Scientific Publishers. p. 110.

M.H. Fulton, P.B. Key, M.E. DeLorenzo (2013) Insecticide Toxicity in Fish Physiology Volume 33, 2013, Pages 309-368. [journal name?](#)

Liu L, Chen Z, Yang Y, Xiao Y, Liu C, Ma Y, Soberón M, Bravo A, Yang Y, Liu K (2018) A single amino acid polymorphism in ABCC2 loop 1 is responsible for differential toxicity of *Bacillus thuringiensis* Cry1Ac toxin in different Spodoptera (Noctuidae) species. *Insect Biochem Mol Biol* 100:59–65.

Formatted: Font: Italic

Ortiz-Urquiza, Z. Luo, and N. O. Keyhani, "Improving mycoinsecticides for insect biological control," *Applied Microbiology and Biotechnology*, vol. 99, no. 3, pp. 1057–1068, Feb. 2015.

Hansa Kumari Jat and Dr. VK Shrivastva, "Cost benefit analysis of newer insecticide used in control of brinjal shoot and fruit borer (*Leucinodesorbonalis* Guenee) at Gwalior region of Madhya Pradesh", *The Pharma Innovation Journal* 12(3): 3091-3093, 2023.

Formatted: Font: Italic

Pesticide Product Information System; U.S. Environmental Protection Agency, U.S. Government Printing Office: Washington, DC, 2012

Basu AC, Pramanik LM. Acaricidal tests of nine pesticides against the two-spotted spider mite, a serious pest of brinjal (eggplant) in west Bengal. *Journal of Economic Entomology* 1968;61(3):768-770.

Formatted: Font: Italic

Butani DK, Verma S. Pests of vegetables and their control Brinjal. *Pesticides* 1976;17(9):6-13

Briere JF, Pracros P, Le Roux AY, Pierre JS. A novel rate model of temperature-dependent development for arthropods. *Environmental Entomology* 1999;28(1):22-9

Formatted: Font: Italic

Colinet H, Sinclair BJ, Vernon P, Renault D. Insects in fluctuating thermal environments. *Annual Review of Entomology* 2015;7(60):123-40.

Formatted: Font: Italic

Das BC, Sarker PK, Rahman MM. Aphidicidal activity of some indigenous plant extracts against bean aphid *Aphis craccivora* Koch (Homoptera: Aphididae). *Journal of Pest Science* 2008;81(3):153-9.

Formatted: Font: Italic

Kaloo G. Solanaceous crops. *Vegetable breeding* 1988;2:520-70.

Mathirajan VG, Natarajan K, Kuttalam S, Chandrasekaran S, Regupathy A. Efficacy of lambda cyhalothrin (karate 5 EC against brinjal shoot and fruit borer (*Leucinodesorbonalis* Guen.). *Pesticide Research Journal* 2000;12(1):117-119.

Formatted: Font: Italic

Rakibuzzaman M, Mahato AK, Husna MA, Maliha M, Jamal Uddin AF. Influence of natura one and neem oil on growth and yield of brinjal (*Solanum melongena*). *Journal of bioscience and Agriculture Research* 2019;20(02):1694-9.

Formatted: Font: Italic

Singh YP, Singh PP. Bioefficacy of insecticides in combination with carbofuran against brinjal shoot and fruit borer (*Leucinodesorbonalis* Guen.) at medium high altitude hills of Meghalaya. *Indian Journal of Plant Protection* 2003;31(2):38-41.

Formatted: Font: Italic

Srinivasan G, Babu PCS, Reddy PP, Kumar NKK, Verghese A. Advances in IPM for horticultural crops. In: Proceedings of the first national symposium on pest management in horticultural crops: environmental implications and thrusts 1998, 87-93. [journal name??](#)

Umapathy G, Baskaran P. Bioefficacy of certain synthetic pyrethroids against major pests of brinjal. *Madras Agric. J* 1991;78:8-10.

Formatted: Font: Italic

Akhilesh Kumar Patel, Pradeep Kumar, Lokesh Kumar Harinkhede, Amit Kumar Patel and Brajendra Ahirwar. Comparative efficacy of different botanicals and biopesticides for management of brinjal shoot and fruit borer (*Leucinodesorbonalis*Guenee) under Bundelkhand region. *The Pharma Innovation Journal* 2021; SP-10(9): 470-474.

Formatted: Font: Italic

Pooja,G.S.andKumar,A.(2022).Fieldefficacyandeconomicsofdifferentinsecticidesagainstbrinjal shoot and fruit borer (*Leucinodesorbonalis*Guenee). *The Pharma InnovationJournal*,(7):4545-4548.

Jaydeep Halder, Totan Adakand Sujana Majumder. Bioefficacy, phytotoxicity and insecticide residue dynamics of chlorantraniliprole in brinjal (*Solanum melongena*) under field condition. *Indian Journal of Agricultural Sciences* 92 (10): 1219–1224, October 2022.

Naresh JS, Malik VS, Balan JS. Estimation of fruit damage and larval population of brinjal fruit borer, *Leucinodesorbonalis*Guenee and its parasitism by *Trathala* sp. on brinjal. *Bulletin of Entomology New Delhi* 1986;27(1):44-7.

Formatted: Font: Italic

Halder J, Kushwaha D and Rai A B. 2020. Biology and feeding potential of *Eocantheconafurcellata*(Wolff) on its lesser known prey, *Spilosomaobliqua*(Walker). *Journal of Biological Control* 34(2): 109–12.

Halder J, Kushwaha D, Rai A B, Singh A and Singh B. 2017. Potential of entomopathogens and neem oil against two emerging insect pests of vegetables. *Indian Journal of Agricultural Sciences* 87(2): 220–24.

Sahu M, Adak T, Patil N B, Pandi G G P, Gowda G B, Yadav M K, Annamalai M, Golive P and Jena M. 2019. Dissipation of chlorantraniliprole in contrasting soils and its effect on soil microbes and enzymes. *Ecotoxicology and Environmental Safety* 180: 288–94.

Formatted: Font: Italic

Vijayasree V, Bai H, Beevi S N, Mathew T B, Kumar V, George T and Xavier G. 2013. Persistence and effects of processing on reduction of chlorantraniliprole residues on cowpea fruits. *Bulletin of Environmental Contamination and Toxicology* 90(4): 494–98.

Singh M, Sachan SK. Comparative efficacy of some biopesticides against shoot and fruit borer, *Leucinodesorbonalis*Guenee in Brinjal. *Plant Archives* 2015;15(2):805-8.

Formatted: Font: Italic

Srinivasan R. Integrated Pest Management for eggplant fruit and shoot borer (*Leucinodesorbonalis*) in south and southeast Asia: Past, Present and Future. *Journal of Biopesticides* 2008;1(2):105-12.

Formatted: Font: Italic

Jat KL, Pareek BL. Field evaluation of ecofriendly insecticides against brinjal shoot and fruit borer, *Leucinodesorbonalis* Guen. *Indian Journal of Plant Protection* 2001;29(1):53-6.

Formatted: Font: Italic

Jyothi DP. Biorational approaches for the management of brinjal shoot and fruit borer (Doctoral dissertation UAS, Dharwad) 2006.

Choudhary B. Vegetables. National book trust, New Delhi 1970, 25-50.

Anand GS, Sharma RK, Shankarganesh K. Evaluation of bio-efficacy and compatibility of emamectin benzoate with neem based biopesticide against fruit borers of brinjal and okra. *Indian Journal of Agricultural Sciences* 2014;84(6):746-53.

Formatted: Font: Italic

Dhankar BS. Progress in resistance studies in the eggplant (*Solanum melongena* L.) against shoot and fruit borer (*Leucinodesorbonalis* Guen.) infestation. *Trop Pest Manag* 1988;34:343-345.

Formatted: Font: Italic

Dhaliwal NK, Aggarwal N. Development and survival of brinjal shoot and fruit borer *Leucinodesorbonalis* Guenee (Crambidae: Lepidoptera) at constant and alternating temperatures. *International Journal of Tropical Insect Science* 2021;41(2):1717-28.

Formatted: Font: Italic

UNDER PEER REVIEW