

Original Research Article

**Ethno-Botanical Uses and Socio-Cultural Importance of White Yam
(*Dioscorea rotundata* Poir.) in South-Eastern, Nigeria**

ABSTRACT

Indigenous knowledge and uses of white yam (*Dioscorea rotundata* Poir.) in South-Eastern, Nigeria was investigated. White yam is the most important species of yam in West Africa cultivated for the consumption of its starchy tuber. Yam plays a major role in the socio-cultural significance of the people of South-Eastern, Nigeria. A well-structured, pre-tested questionnaire on the uses and socio-cultural importance of yam as well as economic status/income generation of farmers was designed and administered to obtain crucial information from the respondents within the study area. Five South-Eastern states of Nigeria namely; Abia, Anambra, Ebonyi, Enugu and Imo states were randomly surveyed. The results showed that 99.16% of the respondents in Anambra state use yam for religious worship, bride price/traditional marriages, funerals, cultural and masquerade festivals. Respondents utilized yam in diverse food forms such as fries/chips, cooked/boiled, roasted, pounded and porridge. It was also observed that respondents utilized yam as a means of livelihood to generate income and support their families. The levels of income generation from yam farming varied. Overall, respondents within the study area who were aware of the indigenous knowledge of yam were significantly ($p < 0.05$) higher than those who were unaware. This ethno-botanical study revealed the traditional concepts and practices relating to socio-cultural importance of yam.

Keywords: Ethno-Botanical Uses, Socio-Cultural Importance, South-Eastern States of Nigeria, White Yam.

1. Introduction

The term “Ethno-botany” is a multidisciplinary science that deals with the study of how the people of an exacting culture and region make use of plants. [1] stated that “Ethno-botany” is an interdisciplinary field of research with specific focus on the empirical knowledge of indigenous people with respect to natural plant substances that influence health and wellbeing and their associated risk. Different *Dioscorea* species plays a remarkable position in the traditional medicines for the treatment of various diseases [2]. There are numerous reports available on ethno-medicinal uses of different *Dioscorea* species worldwide [3]-[5]

Comment [u1]: See journal's rules

White yam (*Dioscorea rotundata* Poir.) according to [6], [7] is the most important specie of yam in West Africa. It is a monocotyledonous, perennial herb cultivated for the consumption of its starchy tuber [8]. [9] reported that the principal food nutrient in yam is mostly carbohydrate and low content of protein as well as ascorbic acid. [10]-[11] reported that consumption of yam may be by preparation of varieties of palatable dishes from yam tubers. [7] pointed out that the yam tuber is the only economical part of the crop, consumed roasted, fried, boiled, pounded or used as flour for baking and steaming for swallowing with soup. [12] reported that yams contribute significantly to food security and its availability in the market for a considerable part of the year helps prevent food shortages because it stores relatively longer than other food crops.

[13]-[14] reported that yam is considered a man's crop and has ritual and socio-cultural significance. In addition to its nutritional value, [15]-[16] reported that yam also plays a major role in the socio-cultural significance of the people mostly the South-Eastern Nigeria and also in the middle belt of Nigeria among the Tiv tribe [17]. [18] reported that yam is a highly revered cultural crop and key festivities like marriage, chieftaincy ceremonies, conflict resolution, peace accords and sacrifices to the gods are tied to it. [19] emphasized that our dietary footprints and food habits are to a large extent, ethnically, regionally and culturally interconnected. This study aims to emphasize the uses and socio-cultural importance of yam by indigenous people of the South-Eastern, Nigeria.

2. Materials and Methods


2.1. Study Area

Geographically, South-Eastern Nigeria extends from latitudes $4^{\circ} 40^{\prime}$ to $7^{\circ} 20^{\prime}$ North and $6^{\circ} 00^{\prime}$ to $8^{\circ} 20^{\prime}$ East longitude. The map of Nigeria showing the boundary area of the South-

Comment [u2]: Authors must state this in the method:
- when did the data collection take place?
- Where was it conducted? in the participants' homes?
- How was the information recorded? recorded or written down?
- What were the inclusion criteria for the participants? How did the researchers reach the participants?
- As the research involved human beings, the authors must present ethics committee approval;
- How the data collected was processed and analyzed

Translated with DeepL.com (free version)

Eastern States of Nigeria is shown below (Figure 1). According to reports, 98% of the indigenous ethnic groups in South-East are Igbos by tribe while 2% of the population is the Igala people who lived in the North-Western part. The natural vegetation in many parts of South-Eastern states is mainly grassland and woodland as well as tropical rainforest. Also, the annual rainfall is between 1,400mm in the north to 2,500mm in the south with soil pH in some parts ranging from 3.5 to 6.4 [20].



Comment [u3]: Present a figure with better resolution

Figure 1: Map showing the Five South-Eastern States of Nigeria [21].

2.2. Ethno-Botanical Study of White Yam (*Dioscorea rotundata* Poir.).

The method of [22], [18], [23], [24] was adopted in this present study. This involved the use of a well-structured, pre-tested questionnaire to obtain vital information from the respondents in the field. The fieldwork process used interviews, public meetings, local informants, review of secondary and grey literatures and keen observations [18]. The survey on Ethno-botanical

study was conducted between September, 2022 to December, 2022 and September, 2023 to December, 2023 in five states of South-Eastern, Nigeria. The states comprises of Anambra, Abia, Enugu, Ebonyi and Imo.

The interviews were segmented into two demographic categories (60 years and above and below 60 years) to understand generational differences and associated perceptions related to the topic of study [18]. A public meeting was organised at the point of yam loading and offloading. [18]reported that these informal yam stations signify a wider representation of yam farmers and distribution of stakeholders from all demographic categories. These stakeholders were interviewed to discern their level of perception of yam farming and livelihood practices. A single interview lasted on average between 45 minutes to one hour. Four local informants were used to facilitate community entry, interactions with interviewees, clarification of issues and interpretations of local dialects.

Field notes were recorded and later categorized and thematized. The questionnaires were sorted accordingly and data collected were used for analysis. The plant materials collected during the field study were pressed, preserved and dried following the standard method of preparation of herbarium techniques [25]. These herbarium specimens were deposited with the taxonomist and curator, Mr. FinianIroka of Botany Department, NnamdiAzikiwe University Awka, Anambra State, Nigeria with their respective assigned voucher numbers, after proper identification and authentication.

2.3. Sampling Frame

The respondents to the questionnaire were the Igbo speaking ethnic group of South-Eastern States of Nigeria. The choice of selecting these areas was because they were predominantly

found in Igbo land and also to compare the previous study by [18], [24] who concentrated their study in one out of the five South-Eastern states, namely; Akwa-Ibom, Anambra and Cross River states. In addition, the South-Eastern states have similar cultural and traditional beliefs as well as reverence for yam. All the people speak the same language, Igbo with slight variations in their pronunciations and intonations.

2.4. Sample Size/Sampling Procedure: Administration of questionnaire

A well-structured, pre-tested questionnaire was designed and administered to assist in obtaining crucial information from the respondents in the study area. One hundred and twenty (120) questionnaires were randomly distributed to each of the five South-Eastern States making a ground total of six hundred (600) sampled questionnaires used in this survey.

The questionnaire was constructed to get vital information from respondents as follows;

- i. Personal data/details such as age, sex, educational qualification, occupation, etc.
- ii. Indigenous knowledge of yam cultivation.
- iii. Labour practices and gender roles in yam cultivation.
- iv. Methods of yam cultivation adopted in their locality.
- v. Yam storage methods and distribution across livelihoods.
- vi. Socio-cultural perspectives on yam distribution practices.
- vii. Uses of yam in worship, marriages, funerals and festivals.
- viii. Disease awareness on yam.
- ix. Causes of post harvest disease of yam.
- x. Methods of post harvest disease control of yam.

3. Results

The socio-cultural use of yam by the respondents in their various communities was surveyed in this study. Respondents gave various uses of yams in their community and their interactions within the community with yams for different cultural activities. These respondents interviewed had indigenous knowledge of yams in their community. Some of the uses of yams in their locality include; religious worship/religious rites, bride price/traditional marriages, funerals, cultural festivals such as chieftaincy title taking “ichiozo” and masquerade festival. In Anambra state, 99.16% of the respondents reported that they use yam for religious worship/religious rites, bride price/traditional marriages, funerals, cultural festivals and masquerade festivals. This is followed by Abia state and Imo state where 97.41% and 96.37% of the respondents reported the same respectively (Table 1). The least was observed in Enugu state where 87.83% of the respondents reported that they use yams for all the socio-cultural practices. In Ebonyi state, 6.09% of the respondents reported that they use yam for religious worship/religious rites. In Enugu state, 7.63% of the respondents reported that they use yam for bride price in their community. There was a significant ($p < 0.05$) difference between the respondents who use yam for all the socio-cultural practices in their community to those who use yam for socio-cultural practices separately (Table 1).

Yam is an important staple food source and has its usefulness in several other social and cultural functions. The diverse food forms and preparations from yam consumed by the respondents were investigated. The various food forms from yam which respondents consumed includes; fries/chips, cooked/boiled, roasted, pounded and porridge. In Anambra state, 50.15% of the respondents reported that they consume yam in all the diverse food forms enumerated such as fries/chips, cooked/boiled, roasted, pounded and porridge. This is

closely followed by Ebonyi state where 49.95% of the respondents affirmed to consuming yam in all the aforementioned diverse food forms. The least was recorded in Imo state where 47.37% of the respondents affirmed that they consume yam in the aforementioned diverse food forms (Table 2). There was a significant ($p < 0.05$) difference between the respondents who consume yam in all the aforementioned diverse food forms to those who consume yam separately in their diverse food forms. In Abia state, 18.10% of the respondents interviewed reported that they prefer consuming yam as fries/chips. In Anambra state, 8.40% of the respondents preferred consuming yam when cooked or boiled. Respondents who had preference to roasted yam were recorded in Enugu state with 12.71% respondents. The highest value for preference for pounded yam was recorded in Anambra state where 10.40% of the respondents registered their preference. In Abia state, 26.73% of the respondents registered their preference for the consumption of yam in porridge form (Table 2). Plate 1 below shows the diverse food forms and preparations from yams consumed by respondents in their communities.

The local names of white yam (*Dioscorea rotundata* Poir.) varieties cultivated and consumed by the respondents in their community were evaluated. Some of the prevalent yam varieties cultivated and consumed by the respondents in their various communities were identified and recorded with their local names (Table 3). The local names of these white yam varieties cultivated and consumed by the respondents in their locality varied in their pronunciations and spellings based on the various dialects found within the study area.

The survey on the sustainability of yam farmer's income revealed that 99.14% of the respondents in Abia state reported that their income was not sustainable. This was followed by Anambra state where 97.48% of the respondents reported that their income was not

sustainable. The least value ($p < 0.05$) was obtained in Imo state with 91.05% of the respondents stating that their income was not sustainable (Table 4). Some of the yam farmers agreed that their income was sustainable. The highest percentage of yam farmers who affirmed that their income was sustainable was observed in Imo state with 8.95% of the respondents while the least was observed in Abia state with 0.86% respondents. Overall, respondents who reported that their income was not sustainable was significantly ($p < 0.05$) higher than those who said that their income was sustainable (Table 4).

Respondents were interviewed to evaluate if yam farmers could support his family and make a decent livelihood out of yam farming. The highest value ($p < 0.05$) was obtained in Abia state where 99.14% of the respondents affirmed that they cannot support their family and make a decent livelihood from yam farming. This was followed by Anambra state with 97.48% of the respondents reporting same. The least was observed in Imo state with 90.70% of the respondents reporting same. Respondents who cannot support their family and make a decent livelihood from yam farming were significantly ($p < 0.05$) higher than those who could support their family and make decent livelihood from yam farming (Table 4).

The level of income generation from yam farming by yam farmers was evaluated. Depending on their profit margins annually, their level of income generation from yam farming was determined. In Anambra state, 70.59% of the respondents revealed that they make an average income between ₦40,000 to ₦59,000 from yam farming annually. This was followed by reports from Imo state and Ebonyi state where 69.65% and 66.04% of the respondents reported same respectively. The least value was recorded in Enugu state where 63.90% of the respondents reported same (Table 5). There was a significant ($p < 0.05$) difference between yam farmers who earned an income between ₦40,000 to ₦59,000 to those who earned higher

Comment [u4]: Provide information in US dollars or Euros so that readers in other regions can have a reference parameter.

Comment [u5]: Idem

income from yam farming. Yam farmers that earned an income between ₦60,000 to ₦79,000 was significantly ($p < 0.05$) higher than those who earned income between ₦80,000 to ₦99,000. The highest percentage was recorded in Anambra state where 27.73% of the respondents earned income between ₦60,000 to ₦79,000. This was followed by the values obtained in Abia state and Ebonyi state where 25.17% and 22.61% of the respondents reported that they earned the same income respectively (Table 5). Just few yam farmers reported that they earned an income of ₦100,000 and above. The highest value was recorded in Ebonyi state where 3.96% of the respondents reported that they earned an income of ₦100,000 and above. On the other hand, 19.30% of the respondents in Imo state reported that they earned an income below ₦40,000. The least value was obtained in Ebonyi state and Anambra state where 0.00% and 0.84% of the respondents respectively reported that they earn an income below ₦40,000 (Table 5).

Comment [u6]: Idem

Comment [u7]: Idem

Comment [u8]: Idem

Comment [u9]: Idem

Comment [u10]: Idem

Comment [u11]: Idem

Comment [u12]: Idem

Respondents provided additional information about their yam farming activities in their locality. Most of the respondents interviewed reported that they lack financial aid/support from the government and need such urgently. In Enugu state, 59.32% of the respondents reported that they require financial support from the government to boost their yam farming business. This is followed by reports from Ebonyi state and Anambra state where 52.17% and 49.58% of the respondents reported that they urgently need financial assistance in the form of loans and grants from the government. Similarly, in Imo state, 63.16% of the respondents reported that they lacked sufficient support either as a subsidy or improved seedlings for use. In Anambra state, 35.29% of the respondents reported that flooding of yam stead affected storage periods during the peak of the raining season. About 15.13% of the respondents pointed out that income from yam production was not sustainable. In Ebonyi state, 30.44% of

the respondents reported that yam is a male crop and requires tedious labour. About 17.39% of the respondents stated that yam is a cultural crop and has its norms and cultures (Table 6).

UNDER PEER REVIEW

Table 1: Socio-Cultural Uses of White Yam (*Dioscorea rotundata* Poir.)

Socio-Cultural Uses of Yam (%)						
States	Religious Worship	Bride Price	Funerals	Cultural Festivals	Masquerade Festival	All of the Above
Abia	0.00	2.59	0.00	0.00	0.00	97.41
Anambra	0.00	0.00	0.00	0.84	0.00	99.16
Ebonyi	6.09	3.48	0.87	1.74	0.00	87.83
Enugu	0.00	7.63	0.00	2.54	0.00	89.83
Imo	0.00	3.63	0.00	0.00	0.00	96.37

Comment [u13]: All figure and table titles must answer what? where? and when?
See all cases

UNDL

Table 2: Diverse Food Forms and Preparations from Yams Consumed by Respondents

DiverseFoodForms (%)						
States	Fries/Chips	Cooked/Boiled	Roasted	Pounded	Porridge	All of the Above
Abia	18.10	4.31	0.86	1.72	26.73	48.28
Anambra	14.29	8.40	6.72	10.40	10.04	50.15
Ebonyi	13.04	6.09	8.76	6.68	15.48	49.95
Enugu	11.69	6.78	12.71	3.39	16.96	48.47
Imo	15.48	8.21	7.96	9.86	11.12	47.37

Table 3: Local Names of White Yam Varieties in the South-Eastern Nigeria

States	Local Names of Cultivated and Consumed White Yam Varieties
Abia	Abiaotulugo, Ji-Mgbada, Ji-Agba-aka, Epe, Ji-Igwe, Ji-Arueri, Nkoroto
Anambra	Epe, Mumuye, Adaka, Ji-Iyoo
Ebonyi	Ji-Ekpuruma, Ji-Mbila, Ji-Iyoo, Oke-Ji, Ji-Oku, Ji-Mmanu (Nwuny ji)
Enugu	Igume, Unegbe, Ji-aga, Nwanyikochoo, Aga-Nvuma, Ono
Imo	Ede-Ji, Ji-Awom, Nkukpo-Oba, Ji-Oke, Ngaraga, Nkukpoagwa, Ji-Oma, Nwanyi-Aga




Plate 1: Diverse Food Forms and Preparations from Yams Consumed by Respondents

(a) Fries/Chips (b) Cooked/Boiled (c) Roasted yam (d) Roasted yam paired with palm oil sauce (e) Pounded yam paired with “ofensala” (f) Porridge

Table 4: Sustainability of Yam Farmer’s Income for Family Support and Livelihood from Yam Farming

States	Sustainability of Yam Farmer’s Income from Yam Farming		Family Support and Livelihood	
	Yes (%)	No (%)	Yes (%)	No (%)
Abia	0.86	99.14	0.86	99.14
Anambra	2.52	97.48	2.52	97.48
Ebonyi	7.83	92.17	9.13	90.87
Enugu	6.61	93.39	4.24	95.76
Imo	8.95	91.05	9.30	90.70

Table 5: Level of Income Generation from Yam Farming (Naira)

States	Level of Income Generation from Yam Farming (%)				
	₦40,000- ₦59,000	₦60,000- ₦79,000	₦80,000- ₦99,000	₦100,000 and above	Below ₦100,000
Abia	64.83	25.17	0.00	0.00	10.00
Anambra	70.59	27.73	0.84	0.00	0.84
Ebonyi	66.04	22.61	7.39	3.96	0.00
Enugu	63.90	17.63	5.93	3.22	9.32
Imo	69.65	3.51	7.02	0.53	19.30

Comment [u14]: Use an international currency as a referenc

Table 6: Additional Information about Yam Farming

States	Additional Information from Respondents	Frequency	Percentage(%)
Abia	Lack of financial support from the government	49	42.24
	None	67	57.76
Anambra	Financial aid/loans highly solicited for	59	49.58
	Flooding of yam stead affects storage period	42	35.29
	Income from yam production is not sustainable	18	15.13
Ebonyi	Grants and financial aid from the government	60	52.17
	Yam is a cultural crop and has its traditions	20	17.39
	Yam is a male crop that requires tedious labour	35	30.44
Enugu	Financial aid from the government is needed	70	59.32
	None	48	40.68
Imo	Financial aid from the government	42	36.84
	Lack of sufficient support for farmers either as subsidy or improved seedling for use	72	63.16

4. Discussion

Socio-cultural uses of yam by the respondents in their various communities across the South-Eastern states of Nigeria were surveyed. In this study, it was observed that respondents had indigenous knowledge of yam in their communities for various cultural and religious purposes such as religious worship/religious rites, bride price/traditional, funerals, cultural festivals such as chieftaincy title taking “ichiozo” and masquerade festivals (Table 1). A great majority of the respondents noted that they utilize yam for all the aforementioned cultural and religious purposes. This is in tandem with [26], [27], [18] who reported that among Nigeria’s Igbo ethnic group, yam is the most favourite food and has a purpose in social functions such as marriages, burials and other traditional ceremonies and rituals. [18] also added that all ethnic groups in South-Eastern Nigeria share a common perception of yam as a religious, social and cultural crop. This perception largely explains why the annual yam festival is a shared cultural heritage across the region.

It was observed that the respondents consumed yam in diverse food forms. Yam is an important staple food and has its usefulness in several other social and cultural functions [28]. The diverse food forms from yam consumed by the respondents includes; fries/chips, cooked/boiled, roasted, pounded and porridge. Most of the respondents affirmed that they consume yam in the aforementioned diverse food forms. This observation resonates with [29], [30], [18] who reported that beyond its cultural and religious values, yams are nutritious and beneficial to human health. They affirmed that yam can be eaten boiled, roasted, in a pounded form or processed into yam chips, snacks and flakes.

Local names of white yam varieties cultivated and consumed by the respondents in their community were identified and recorded (Table 3). The local names of these white yam

varieties cultivated and consumed by the respondents in their locality varied in their pronunciations and spellings based on the various dialects found within the study area. This is in tandem with the study by [24] who observed that respondents from Obudu local government area of Cross River state, Nigeria has a special breed of local white yam called “Atam yam” in their dialect.

Majority of yam farmers interviewed during this survey reported that their income was not sustainable and could not support their family and make a decent livelihood out of yam farming (Table 4). This observation is in tandem with the study of [14], [31], [24] who reported that yam farmers complement their income through extra paid labour services and through the sale of other farm produce they realised from their farms.

A great number of yam farmers affirmed that they make an income of ~~₦40,000-₦59,000~~ annually as their level of income generation from yam farming while about 7.71% of the respondents make an income of ~~₦100,000~~ and above (Table 5). This is in contrast with [31] who reported that ginger farmers make an annual income of ~~₦72,487.5~~ on average. There were yam farmers who affirmed that they make annual income of below ~~₦100,000~~ as well.

Most of the respondents interviewed reported that they require financial aid/support from the government urgently, grants or support either as subsidy or improved seedlings for use as additional information. Others reported that their income from yam cultivation was not sustainable. A 54 years old female respondent who retails yam tubers at the Otuocha market in Anambra state said that flooding of yam stead affects storage period at the peak of the raining season. A 39 years old male respondent in Ebonyi state reported that yam is a cultural and has its norms and culture. He also said that yam is a male crop requiring arduous labour.

This view is in sync with observation by [14] who reported that men and women hold different rights of production, labour, ownership, access and marketing of specific crop types. [18] pointed out that designating specific crops for specific gender category has the capacity to produce differential effects in productivity, vulnerability to shock and income generation prospects.

There were limitations encountered during the survey of ethno-botanical study. The first is the sample size. It is important to state that one hundred and twenty (120) questionnaires are not enough to obtain all the vital information from each state of South-Eastern, Nigeria. Secondly, relevant stakeholders such as traditional rulers, religious priests, public officials and urban dwellers were not interviewed. This inability to interview these individuals means that the findings in this study do not represent every segment of what is applicable about yam in South-Eastern, Nigeria. Furthermore, insecurity in the South-Eastern states of Nigeria restricted interviews in major yam cultivating zones of the region during this survey.

5. Conclusion

Respondents from the five South-Eastern states of Nigeria had indigenous knowledge of yam. This study has demonstrated the importance and functions of yam in structuring the social, economic, religious and cultural fabric of a society. Findings from this survey will be utilized as a basis for further investigations on the emerging indigenous practices and sustainability impact.

Disclaimer (Artificial Intelligence)

Authors hereby declare that NO generative AI technologies such as large language models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscript.

References

Comment [u15]: References must be up-to-date. Only 3 references from the last five years.

- [1] Gurib-Fakim, A. (2006). Medicinal Plants: Traditions of Yesterday and Drugs of Tomorrow. *Molecular Aspiration for Medicine*,**27**:1–93.
- [2] Kumar, S., Das, G., Shin, H. S., and Patra J. K. (2017). *Dioscorea* spp. (A Wild Edible Tuber): A Study on Its Ethnopharmacological Potential and Traditional Use by the Local People of Similipal Biosphere Reserve, India. *Frontiers Pharmacology*,**8**: 52.
- [3] Mishra, S., Swain, S., Chaudhury, S. S., and Ray, S. (2008). Wild Edible Tubers (*Dioscorea* spp.) and their Contribution to the Food Security of Tribes of Jeypore Tract, Orissa, India. *PGR Newsletter*,**56**: 63–67.
- [4] Sharma, L. N., and Bastakoti, R. (2009). Ethnobotany of *Dioscorea* L. with Emphasis on Food Value in Chepang Communities in Dhading District, Central Nepal. *Botanica Orientalis Journal of Plant Science*,**6**: 12–17.
- [5] Sheikh, N., Kumar, Y., Misra, A. K., and Pfoze, L. (2013). Phytochemical Screening to Validate the Ethnobotanical Importance of Root Tubers of *Dioscorea* species of Meghalaya, North East India. *Journal of Medicinal Plants Studies*,**1**: 62–69.
- [6] Onwueme, I. C. (1978). *The Tropical Tuber Crops, Yam, Cassava, Sweet Potato and Cocoyams*. John Willey and Sons, New York. 234 Pp.
- [7] Okigbo, R. N., Opara, P. U., and Anuagasi, C. L. (2015a). Efficacy of Extracts of Water Yam (*Dioscorea alata*) and aerial yam (*Dioscorea bulbifera*) Peels in the Control of White Yam (*Dioscorea rotundata*) Rot. *Journal of Agricultural Technology*, **11**(8): 1823- 1842.
- [8] Ezeibekwe, I. O., Opara, M. I., and Mbagwu, F. N. (2009). Antifungal Effects of *Aloe vera* Gel on Fungal Organisms Associated with Yam (*Dioscorea rotundata* Poir) Rot. *Journal of Molecular Genetics*, **1**(1): 11-17.
- [9] Igyor, M. A., Ikyo, S. M. and Gernah, D.I. (2004). The Food Potential of Potato Yam (*Dioscorea bulbifera*). *Nigerian Food Journal*,**22**(1): 209-215.
- [10] Fasasi, A. R. and Fasina, O. O. (2005). *Resources, Uses and Efficiency in Yam Production in Ondo State, Nigeria: Agricultural Rebirth for Improved Production in Nigeria*. In: Proceedings of the 39th Annual Conference of Agricultural Society of Nigeria (ASN), held at the University of Benin, Benin City, Nigeria. Pp 184-186.
- [11] Oladebo, J. O. and Okanlawon, O. M. (2010). *Profitability Level of Yam (Dioscorea spp.) Production in Oyo State of Nigeria*. In: Proceedings of the 44th Annual Conference of Agricultural Society of Nigeria, held at Ladoké Akintola University of Technology, Ogbomosho. Pp 57-59.
- [12] Opara, E. U., and Nwokocha, N. J. (2015). Antimicrobial Activities of Some Local Medicinal Plants against Post Harvest Yam Rot Pathogens in Humid South Eastern Nigeria. *Journal of Microbiology Research and Reviews*, **3**(1): 1-9.
- [13] Hahn, S. K., Osuiru, D. S. O., Akoroda, M. O., and Otoo, J. O. (1987). Yam Production and its Future Prospects. *Outlook on Agriculture*, **16**: 105-110.
- [14] Kleih, U., Philips, D., Mignouna, D., Ogbonna, M., and Siwoku, B. (2012). Nigeria Scoping Yam Value Chain Analysis. In: *Yam Improvement for Income and Food Security in West Africa*. Gatesopenresearch, London UK. 142 Pp.

- [15] Sangoyomi, T. E. (2004). Post-Harvest Fungal Deterioration of Yam (*Dioscorea* spp.) and its Control. Ph.D Dissertation (Unpublished), University of Ibadan, Nigeria. 179 Pp.
- [16] Okigbo, R. N., Anuagasi, C. L., and Nwanna, C. R. (2015b). Control of Yam Rot Fungi with Crude Extracts of *Chromolaenaodorata* and *Cympobogoncitratrus*. *Nigerian Journal of Mycology*, **7**: 44-54.
- [17] Orkwor, G. C. (1992). *Yams Production: Cropping systems, Ecology, Development and Research in Nigeria*. Paper Presented at the National Seed and Plant Quarantine Project, Nigeria.
- [18] Obidiegwu, J. E., and Akpabio, E. M. (2017). The Geography of Yam Cultivation in Southern Nigeria: Exploring its Social Meanings and Cultural Functions. *Journal of Ethnic Foods*, **4**: 28-35.
- [19] Kittler, P. G., Sucher, K. P., and Nelms, M. N. (2012). *Food and Culture*. Sixth Edition. Belmont, Wadsworth, CA, USA. 344 Pp.
- [20] Igwilo, I.O., Afonne, O.J., Maduabuchi, U.J. and Orisakwe, O.E. (2006). Toxicology Study of the Anam River in Otuocha, Anambra State, Nigeria. *Archaeology, Environmental and Occupational Health*, **61(5)**: 205 – 208.
- [21] Okeke, F. O. (2023). Map of South-East.Avaialabe online at https://www.researchgate.net/figure/map-of-South-East-Geopolitical-zone-of-Nigeria_fig1_350950645. Cited 8th September, 2023.
- [22] Okigbo, R. N. and Nwatu, C. M. (2015). Ethnostudy and Usage of Edible and Medicinal Mushrooms in Some Parts of Anambra State, Nigeria. *Natural Resources*, **6**: 79-89.
- [23] Nilofer, S., Yogendra K., Licha, J., and Nazir, A. B. (2017). Ethnobotanical Uses and Survey of *Dioscorea* Species of North East India: Its Conservation and Sustainable Utilization. *International Journal of Current Research in Biosciences and Plant Biology*, **4(12)**: 117-124.
- [24] Ema, E-O. S., Obidiegwu, J. E., Chilaka, C. A., and Akpabio, E. M. (2023). Indigenous Food Yam Cultivation and Livelihood Practices in Cross River State, Nigeria. *World*, **4**: 314-332.
- [25] Jain, S. K., and Rao, R. R. (1997). *A Hand Book of Field and Herbarium Technique*. Today and Tomorrow Publication, New Delhi. 434 Pp.
- [26] Basden, G. T. (1966). *Niger Ibos: A Descriptive of the Primitive Life, Customs and Animalistic Beliefs of the Ibo People of Nigeria by One Who, for Thirty-Five Years, Enjoyed the Privilege of their Intimate Confidence and Friendship*. Barnes and Noble, New York. 224 Pp.
- [27] Achebe, C. (1975). *Things Fall Apart*. Heinemann Educational Books, Ibadan, Nigeria. Pp 16-17.
- [28] Coursey, D. G. (1976). The Origins and Domestication of Yams in Africa. In: Harlan, J. R., De Wet, J. M. J., and Stemler, A. B. L. (Eds). *Origin of African Plants Domestication*. Mouton, The Hague. Pp 386-403.
- [29] Howard, B. J., and Warren, D. H. (1988). Chemistry of Tropical Root Crops. Significance for Nutrition and Agriculture in the Pacific. *ACIAR Monogram Series*, **6**: 27-37.
- [30] Degras, L. (1993). *The Yam: A Tropical Root Crop*. Macmillian Press Ltd., London, United Kingdom. 464 Pp.
- [31] Mazza, M., Ewuziem, J. E., and Uwandu, Q. C. (2019). Factors Affecting Farmers Income Generation from Ginger Production in Abia and Imo States, Nigeria. *Canadian Journal of Agriculture and Crops*, **4(2)**: 77-83.