

Effect of combined application of treated sewage sludge and inorganic fertilizers on yield and yield attributes of maize crop (*Zea mays* L.)

ABSTRACT

In present day scenario the production wastes have increased tremendously to point where its disposal is a concern. The sewage sludge generated from sewage treatment plants (STP) contains nutrients which are beneficial to plants. Due to these beneficial qualities, it can be used for agricultural purpose. The purpose of the conducted study was to know the effect of combined application of sewage sludge along with inorganic fertilizers of Maize crop. Randomized block design involving 10 treatments and 3 replications was adopted for the study. The experiment was carried out from October 2023 to February 2024 at college farm, College of Agriculture, PJTSAU, Rajendranagar, Hyderabad. The T10 treatment showed significant results when compared with T2 treatment. The results obtained revealed that application of sewage sludge along with inorganic fertilizers (T10) showed significant results when compared to application of inorganic fertilizers alone. The significantly highest grain yield, grain weight cob^{-1} , no. of grains cob^{-1} and no. of seed rows cob^{-1} were observed in T10 treatment followed by T9, T8 and T2. According to the results obtained, sewage sludge can be used as source of nutrients for agricultural application by reducing the quantities of inorganic fertilizers and also reduces resource consumption for disposal of sewage sludge.

Keywords: sewage sludge, inorganic fertilizers, disposal, nutrients, resource.

1. INTRODUCTION

Sewage sludge is an end production of sewage treatment plant (STP). The sewage sludge is the waste product after the aerobic or anaerobic digestion process. The disposal of sewage sludge has always been a concern. In general disposal of sewage sludge includes land filling, incineration etc, which consumes land, time and labor resources. Now a days due to

the increased inflow of people to urban areas led to increased quantities of sewage sludge produced.

Sewage sludge is also known for its nutritional value. The nutrient content of sewage sludge varies depending upon the source from which it is being collected and treated. Sewage sludge consists of organic matter, macro and micro nutrients, trace elements, microorganisms. Due to the presence of essential macro and micro nutrients in sewage sludge, it can be used as fertilizer material and the organic matter present in sewage sludge improves soil properties (Heckman *et al.*, 1986; Roberts *et al.*, 1988).

Use of sewage sludge in agriculture is a way of recycling the wastes produced from STP. Agricultural application of sewage sludge reduces the exploitation of land resources for its disposal (Ciešlik *et al.*, 2015). Application of sewage sludge in agriculture is a cost-effective and environmentally friendly method of disposal (Evans *et al.*, 2004). One of the demerits of application of sewage sludge was presence of heavy metals, so it is necessary to analyze the heavy metal content in sewage sludge (Sharma *et al.*, 2017). The composition of sewage sludge is based on the source of waste produced and treatment process used to treat the generated wastes (Rigby *et al.*, 2016). Agricultural use of sewage sludge after proper treatment improves the soil properties by providing the nutrients back to soil which removed before (Seleiman *et al.*, 2013; Bai *et al.*, 2017).

Various reports from studies conducted previously states that application of sewage sludge improves productivity of crops. The Application of 2% sewage sludge improved the yield of broad bean in calcareous loam soil (Dahdohand Hassan. 1997). Application of sewage sludge increased the productivity of bread wheat (Koutroubaset *et al.*, 2014), Maize (Moreira *et al.*, 2020). Application of 62 Mg ha⁻¹ was reported to increase the grain yield in maize (Binder *et al.*, 2002). Application of sewage was also reported to increase the yield of Barley (Antolín *et al.*, 2005), Sunflower (Koutroubaset *et al.*, 2020).

The present study conducted shows the effect of sewage sludge application in combination with inorganic fertilizers on yield and yield attributes of maize.

2. MATERIAL AND METHODS

Table. 1 characteristics of sewage sludge.

pH	Ec (dS m ⁻¹)	OC (%)	Total N (%)	Total P (%)	Total K (%)
5.9	4.38	12	1.09	0.16	0.019

2.1 Collection of Sewage Sludge

Sewage sludge was collected from STP (50 MLD) Attapur, Hyderabad. The sewage sludge was aerobically digested, treated and chlorinated before its disposal.

2.2 Location of Experimental Plot

The research work was carried out in College farm, College of Agriculture, Rajendranagar, PJTSAU, Hyderabad. Geographically the site is located 17°19'21.0" N latitude and 78°24'38.9"E longitude.

2.3 Treatments and Design of Experiment

Randomized block design was adopted for conducting experiment. The total no. of treatments includes 10 i.e., T1=control(0:0:0), T2 = 100% recommended dosage of fertilizers (RDF), T3= 2.5 t ha⁻¹, T4 = 5 t ha⁻¹, T5 = 50% RDF + 2.5 t ha⁻¹, T6= 50% RDF + 5 t ha⁻¹, T7=75 % + 2.5 t ha⁻¹, T8=75 % + 5 t ha⁻¹, T9=100%RDF+ 2.5 t ha⁻¹, T10=100%RDF+5 t ha⁻¹ were employed in 30 plots. Sewage sludge was incorporated in field 10 days before sowing.

2.4 Crop Management

The crop selected for carrying out the experiment was Maize (*Zeamays* .L). The seed rate of crop is 20 kg ha⁻¹, the recommended dosage of fertilizer is 240:80:80 kg NPK ha⁻¹. The hybrid variety selected was DHM-117. The land was ploughed with disc plough followed by cultivator and harrowing. Furrow and ridges were made with ridger. The spacing adopted for planting was 60×20 cm and the plot was 6×4.2 meters approx.

The crop was sown in *rabi* October 2023. The crop was irrigated regularly. One manual weeding after seed germination was done and pre-emergent herbicide spraying with Atrazine (1 kg 200 lit⁻¹ acre⁻¹) after 2 days of sowing and post emergent herbicide spray with Calaris (400 ml 200 lit⁻¹ acre⁻¹) after 20 days after sowing. The crop was infested with stem borer and fall army worm after one week of germination which were controlled with the foliar application of Chlorantraniliprole @ 4ml lit⁻¹ acre⁻¹.

The weather conditions during the crop period from October 2023 to February 2024. The weekly mean maximum temperature was between 27.7°C to 33.8°C with an average of 30.5°C mean weekly minimum temperature was between 15.3°C and 21.0°C with an average temperature of 17.5°C. The weekly morning relative humidity was 81% to 91% with an average of 85.7% and the weekly afternoon relative humidity was between 31.7% to 62.4% with an average of 41.8%. The total rain fall recorded was 14.4 mm. The weekly mean sunshine hours ranged between 3.5 to 9.3 with an average of 6.5 hours day⁻¹.

2.6 Data Analysis

2.6.1 Plant Sampling and data collection

Five plants from each plot were tagged and the recorded data was collected from the tagged plants. The data collected includes grain yield, grain weight cob⁻¹, no. of grains cob⁻¹, no. of seed rows cob⁻¹ and plant height (cm). The grain weight cob⁻¹, no. of grains cob⁻¹ and no. of seed rows cob⁻¹ were recorded from tagged plants and then averaged. The plant height was measured using meter scale from base to top of the plant for tagged plants in plot and then averaged. For grain yield, the cobs were harvested, dried and then threshed till the moisture content reaches 10% and then presented in t ha⁻¹.

2.6.2 Statistical analysis

The data collected was subjected to analysis of variance for simple randomized block design (Rao, 1983).

3. RESULTS AND DISCUSSION

Table. 2 Effect of different doses of combined application of sewage sludge and inorganic fertilizers.

Treatments	Grain yield (t/ha)	Grain weight ^{cob-1} (g)	No.of grains ^{cob-1}	No. of seed rows ^{cob-1}	Plant height (cm)
T1	2.10	25.17	205.20	12.60	145.07
T2	7.39	88.53	360.27	14.93	173.85
T3	2.18	26.10	224.83	13.40	154.90
T4	2.71	32.43	235.60	13.87	157.33
T5	5.43	65.09	323.13	13.60	168.33
T6	6.06	72.57	347.40	14.20	171.20
T7	7.48	89.66	358.60	14.93	179.97
T8	7.76	92.99	370.03	15.07	180.80
T9	8.11	97.13	375.33	15.13	194.83
T10	8.38	100.37	395.07	16.53	199.50
SEM	0.22	2.61	11.57	0.50	7.08
CD	0.65	7.77	34.37	1.50	21.03
CV	6.56	6.56	6.27	6.06	7.10

$P < 0.01^{**}$

3.1 Grain yield: The grain yield ranged between 2.10 t ha⁻¹ in control to 8.38 t ha⁻¹ in T10. The significant results were obtained with the application of T10 treatment (table 2). The highest yield was obtained with T10 followed by T9 T8, T7, T2, T6, T5, T4, T3 and T1. Significant difference was recorded between the yields obtained from T10 (100% RDF + % t ha⁻¹) and T2 (100% RDF). The yield obtained in T3 and T4 was slightly higher than T1, but significant difference was not observed. Likewise the results recorded with T10 showed significant difference with T2 but on par with T9, T8. This might be attributed to the presence of macro nutrients present in the sewage sludge similar results were reported by [Eghball and Power. \(1999\)](#). The decomposition of organic matter releases organic acids which solubilizes nutrients present in the soil and increases their availability ([Dahdoh and El-Hassanin. 1993](#)).

3.2 Grain weight cob⁻¹: Among the treatments employed the grain weight cob⁻¹ ranged from 25.17g to 100.37 g (table 2). The highest results were recorded with T10 followed by T9, T8, T7, T2, T6, T5, T4, T3 and T1. The weights recorded with T10 was on par with weights obtained with the T7 and T7 was on par with T2. The highest weights in T10 might be attributed to application of sewage sludge along with RDF. The lesser weights obtained in T1 might be due no application of fertilizers. The weights obtained were in similar trend as that of grain yield. Similar results were reported by Hussain. (1999) as he reported that difference in the weights might be attributed to different levels of fertilizers involved in treatments.

3.3 No. of grains cob⁻¹: The no. of grains cob⁻¹ ranged between 205 g to 395 g. The trend was similar to that obtained in grain yield. The highest results were recorded with T10 due to highest doses of fertilizers and sewage sludge applied and the lowest were recorded with T1 plot due to no application of fertilizers and manures (table 2). The results recorded with T10 showed significant difference with T2 and on par with T9 and T8. The results recorded with T2 were on par with T8. This might be due to the application of difference in levels of fertilizers and sewage sludge. The results were in accordance with the results reported by Sharar *et al.*, (2003) and Oktemet *et al.*, (2005) as they reported that no. of grains cob⁻¹ were affected with different doses of fertilizers applied.

3.4 No. of grain rows cob⁻¹: The recorded results ranged between 12.60 to 16.53(table 2). the highest results were recorded with T10 and the lowest results were recorded with T1. The T10 was significantly superior than T2 and was on par with T9 and T8. The trend obtained is similar to that obtained with grain yield. The highest results recorded were attributed to application of highest dosage of fertilizers and sewage sludge. The lesser results recorded were attributed to 0:0:0 application of fertilizers. Similar results were recorded by Ali *et al.*, (2002) and Younas. (2000).

3.5 Plant height:The plant height recorded ranged between 145.07 cm to 199.50 cm (table 2). the highest results were recoded with T10 and the lowest was recorded with T1. The plant height recorded with T10 was on par with T9, T8 and T7 but significant difference was reported with T2. The significantly higher value might be attributed to application of fertilizers and additional availability of nutrients through sewage sludge. The results obtained were in agreement with results reported by Sharar *et al.*, (2003) and Ayub *et al.*, (2002) who reported that availability of different levels of nutrients to plants effects plant height.

4. CONCLUSIONS: Application of sewage sludge in combination with sewage sludge improved Maize yield. T10 treatment showed significant results when compared to T2 treatment. Due to the beneficial properties of sewage sludge, it improves soil properties. Sewage sludge can also replace the inorganic fertilizers to some extent for crop application. Use of sewage sludge in agriculture helps in its disposal. So, it is concluded that application of sewage sludge in agricultural land is way of recycling generated organic wastes and reduces land resource consumption.

5. REFERENCES:

1. Ali J, Bakht J, Shafi M, Khan S and Shah WA. Uptake of nitrogen as affected by various combinations of nitrogen and phosphorus. *Asian Journal of Plant Sciences*. 2002;1: 367-369.
2. Antolín MC, Pascual I, García C, Polo A and Sánchez-Díaz M. Growth, yield and solute content of barley in soils treated with sewage sludge under semiarid Mediterranean conditions. *Field Crops Research*.2005;94:224–237.
3. Ayub M, Nadeem MA, Sharar MS and Mahmood N. Response of maize (*Zea mays* L.) fodder to different levels of nitrogen and phosphorus. *Asian Journal of Plant Sciences*.2002;9:352-354.
4. Bai S, Lyu T, Ding Y, Li Z, Wang D, You S and Xie Q. Campus Sewage treatment in multilayer horizontal subsurface flow constructed wetlands: Nitrogen removal and microbial community distribution. *Clean-Soil, Air, Water*. 2017.45(11):170054.
5. Binder DL, Dobermann A, Sander DH and Cassman KG. Biosolids as nitrogen source for irrigated maize and rainfed sorghum. *Soil Science Society of America Journal*. 2002;66:531–542.
6. Cieslik BM, Namiesnik J and Konieczka P. Review of sewage sludge management: standards, regulations and analytical methods. *Journal of Cleaner Production*.2015;90:1-15.
7. Dahdoh MSA and El-Hassanin AS.1993. Combined effects of organic source, irrigation water salinity and moisture level on the growth and mineral composition of barley grown on calcareous soil. *Desert Institute Bulletin*.1993;43:12
8. Dahdoh MSA, Hassan FA. Combined effect of sewage sludge and saline water irrigation on growth and elemental composition of broad bean. *Egyptian journal of Soil Sciences*. 1997;37(2):189-204.

9. Eghball B and Power JF. Phosphorus and nitrogen-based manure and compost applications: corn production and soil phosphorus. *Soil Science Society of America Journal*. 1999;63:895–901
10. Evans T, Lowe N and Matthews P. Sustainable biosolids – Welcomed practice through community partnership and the consequential economic benefits. *Water Science and Technology*. 2004;49:241–249.
11. Heckman JR, Angle JS and Chaney RL. Soybean nodulation and nitrogen fixation on soil previously amended with sewage sludge. *Biology and Fertility of Soils*. 1986;2:181-185.
12. Hussain I, Mahmood T, Ullah A and Ali A. Effect of nitrogen and sulphur on growth, yield and quality of hybrid maize. (*Zea mays* L.) *Pakistan Journal of Biological Sciences*. 1999;2(3):637-638.
13. Koutroubas SD, Antoniadis V, Damalas CA and Fotiadis S. Sunflower growth and yield response to sewage sludge application under contrasting water availability conditions. *Industrial Crops and Products*. 2020;154:112670.
14. Koutroubas SD, Antoniadis V, Fotiadis S and Damalas CA. Growth, grain yield and nitrogen use efficiency of Mediterranean wheat in soils amended with municipal sewage sludge. *Nutrient Cycling in Agroecosystems*. 2014;100:227–243.
15. Moreira RS, Lense GHE, Fávero LF, Oliveira Junior BMD and Mincato RL. Nutritional status and physiological parameters of maize cultivated with sewage sludge. *CienciaAgrotecnologia*. 2020;44:029919.
16. Oktem AG and Oktem A. Effect of nitrogen and intra spaces on sweet corn characteristics. *Asian Journal of Plant Sciences*. 2005;4:361-364.
17. Rao, G.N. 1983. *Statistics for Agricultural Sciences*. BS Publication, Hyderabad. 250-295.
18. Rigby H, Clarke BO, Pritchard DL, Meehan B, Beshah F, Smith SR and Porter NA. A critical review of nitrogen mineralization in biosolids-amended soil, the associated fertilizer value for crop production and potential for emissions to the environment. *Total Science Environment*. 2016; 541:1310–1338.
19. Roberts JA, Daniels WL, Bell JC and Martens DC. Tall rescue production and nutrient state on southwest Virginia mine soils. *Journal of Environmental Quality*. 1999;17:55-62.
20. Seleiman MF, Santanen A, Kleemola J, Fredrick LS and Pirjo SAM. Improved sustainability of feedstock production with sewage sludge and interacting mycorrhiza. *Chemosphere*. 2013;91(9):1236-1242.
21. Sharar MS, Ayub M, Nadeem N and Ahmad N. Effect of different rates of nitrogen and phosphorus on growth and grain yield of maize. *Asian journal of Plant Sciences*. 2003;2(3): 347-349.

22. Sharma B, Sarkar A, Rajee PS and Singh P. Agricultural utilization of biosolids: a review on potential effects on soil and plant grown. *Waste Management*. 2017; 64:117-132.
23. Younas MH, Rehman and Hayder G. Magnitude of variability for yield and associated traits in maize hybrids. *Asian Journal of Plant Sciences*. 2002;1(6): 694-696.

UNDER PEER REVIEW