


Optimization of Siderophore Production by *Bacillus subtilis*: Antifungal Efficacy Against *Alternaria triticina* and *Bipolaris sorokiniana* and Seed Germination Enhancement Potential

Abstract

Bacillus subtilis (UP11) is a gram-positive, plant growth-promoting rhizosphere bacterium (PGPR) isolated from the rhizosphere soil of wheat. It produces an active compound i.e., siderophore. Siderophores are low-molecular-weight and high-affinity molecules produced by bacteria under iron-limited conditions. In present study, we optimized the culture condition for siderophore production and also investigated the antagonistic activity against the foliar blight pathogens. Therefore, the optimum pH 7 was found maximum siderophore unit (62.95%), mannitol as carbon source (59.14%), ammonium nitrate as nitrogen source (63.04%), heavy metal $Pb(NO_3)_2$ shows (61.36%) highest yield in the succinate broth. The optimal iron concentration 50 μM for siderophore production. All these parameters and their effects were investigated using the one-factor-at-a-time (OFAT) approach. The resulted optimized siderophore is partially purified as in ethyl acetate extract which shows (59.64%, 50.88%) mycelial fungal inhibition against *Alternaria triticina* and *Bipolaris sorokiniana*. After that, this filtrate tested for germination potential, resulted shows that it potential act both biocontrol and plant growth promoting agent. It enhances the % germination, root and shoot length. Our findings indicate that *Bacillus subtilis* can produce substantial quantities of siderophores, which exhibit potent antagonistic activity and promoting growth of plant. This finding holds promise for advancing the development of novel biological control agents.



Graphical abstract – Siderophore producing bacteria (*Bacillus subtilis*) shows both antifungal and plant growth promoting activity

Keywords: Siderophore unit, *Bacillus subtilis*, antagonistic, antifungal

1. Introduction

Microorganisms produce various secondary metabolites for growth and survival. During iron stress, they produce secondary metabolites termed siderophores. Rhizosphere microbiota benefit plants through direct and indirect mechanisms. Direct benefits include the production of phytohormones, siderophores, antioxidants, cell wall-degrading enzymes, and volatile organic compounds (VOCs). Additionally, they enhance plant health by solubilizing minerals such as phosphorus, potassium, and zinc, and by fixing biological nitrogen. Indirect benefits are conferred through antibiosis, induced systemic resistance, and the formation of biofilms (Singh et al., 2022).

Iron is crucial for the growth and survival of most microorganisms. However, in natural environments, iron predominantly exists as insoluble Fe^{3+} (10-18 M at pH 7.0), making it challenging for microorganisms to access. Siderophores are small, low molecular weight secondary metabolites (400–2,000 Da) with a strong affinity for Fe^{3+} . They are produced and secreted by microorganisms to sequester iron from their surroundings as needed (Albelda-Berenguer et al., 2019). In aerobic environments, where Fe^{2+} is scarce, bacteria must actively uptake Fe^{3+} to maintain cellular iron homeostasis. Based on their iron-binding moieties, siderophores can be classified into three basic types: catecholate, hydroxamate, and carboxylate and mixed types (Khan et al., 2020). In iron-deficient environments, microorganisms produce siderophores to acquire iron for growth. Some plant growth-promoting rhizobacteria (PGPR) synthesize siderophores, enabling them to compete with pathogens for iron and inhibit pathogen growth (Leventhal et al., 2019). We previously reported that *Bacillus subtilis* used as biocontrol bacterium against foliar blight pathogens (Jaggi et al., 2020). The primary objective of this study was to identify and maximize siderophores produced by *Bacillus subtilis*.

Various factors, including carbon source, nitrogen source, pH, temperature, and metal ions, influence siderophore synthesis (Srivastava et al., 2022). Additionally, carbon and nitrogen sources play vital roles; for instance, maltose as a carbon source and NH_4NO_3 as a nitrogen source significantly enhance siderophore production in endophytic fungi from *Cymbidium aloifolium* (Chowdappa et al., 2020). Optimal conditions for maximum siderophore production vary between strains. For example, Bendale found that *Streptomyces fulvissimus* achieves peak siderophore production with sucrose as the carbon source (Bendale et al., 2009), whereas Santos identified glycerol as optimal for *Bacillus megatherium* growth (Santos et al., 2014). *Bacillus* sp. PZ-1 showed highest siderophore production with a glucose concentration of 21.84 g/L, *B. brevis* sucrose as carbon source and asparagine as nitrogen (Yu et al., 2017; Sheng et al., 2020). pH is critical; plays sensitive role, neutral pH generally supports maximum siderophore production in fluorescent pseudomonads (Sayyed et al., 2005; Tailor and Joshi, 2012). Iron availability crucially impacts organism growth by influencing siderophore production. Furthermore, the presence of metal ions in the medium, such as cadmium, influences siderophore production in *Streptomyces tendae*, MnCl_2 in *P. monteilii* B8 (Sayyed and Chincholkar, 2010; Srivastava et al., 2022).

Siderophore-producing bacteria are crucial for enhancing crop nutrition and suppressing phytopathogens (Yasmin et al., 2012; Sayyed et al., 2013). Ghazy and El-Nahrawy (2021) studied *Bacillus subtilis* MF497446 and *Pseudomonas koreensis* MG209738, showing their efficacy against *Cephalosporium maydis* in maize. Additionally, Wang et al. (2022) demonstrated that siderophore-producing *Pseudomonas putida* controls *Fusarium* wilt in cucumber, radish, and flax, highlighting their role as effective biocontrol agents (PGPB). Siderophore-producing bacteria offer a sustainable alternative to chemical pesticides in agriculture. They help immobilize toxic metals in soil (Trindade et al., 2021; Hardcore et al., 2022) and assist in bioremediation efforts (Rani et al., 2022). Commercializing these bacteria as bio-inoculants can effectively control fungal diseases by limiting iron availability to pathogens (Basu et al., 2021; Malgioglio et al., 2022), promoting sustainable agriculture. Our study reported that *Bacillus subtilis* produce hydroxamate-catechol mixed type siderophores and has a strong antagonistic effect on wheat foliar blight fungal pathogens. However, in this study also included the factor that influence the production of siderophore production and their effect on antifungal and seed germination were also evaluated.

2. Materials and Methods

2.1 Antagonistic Strains and Medium

Antagonistic wheat rhizobacteria strains (UP11) *Bacillus subtilis* accession no MN099431.1 previously isolated and stored at the Rhizosphere Biology Laboratory, Department of Microbiology, Govind Ballabh Pant University of Agriculture and Technology, Pantnagar, Uttarakhand were cultured and maintained on Nutrient Agar (NA).

2.2 Chrome Azurol S (CAS) medium assay

The CAS agar plate method (Schwyn and Neilands, 1987) was used to qualitatively assay siderophore production. Strains were grown in Nutrient Broth (NB) for 48 hours at 30°C and 120 rpm, then 0.05 ml of a suspension (2×10^8 cells/ml) was spotted onto CAS-agar plates and incubated for 5 days at 30°C. Clear halo around colonies indicated siderophore production.

Quantitative assay

To quantify siderophore activity in culture supernatant extracts, the CAS solution method (Schwyn and Neilands, 1987) was used. Tested strains were grown in Nutrient Broth for 48 hours on a rotary shaker at 30°C and 120 rpm, then centrifuged for 15 minutes at 3000 rpm. A mixture of 0.5 ml of supernatant (2×10^8 cells/ml), 0.5 ml CAS solution, and 10 μ l shuttle solution (sulfosalicylic acid) was incubated at room temperature for 2 hours. Absorbance was measured at 630 nm using a UV/Vis spectrophotometer. Reference samples (blanks) were prepared with Nutrient Broth medium, CAS solution, and shuttle solution. Siderophore units (SU) were calculated using the formula $SU = [(Ar - As) / Ar] \times 100\%$, where Ar represents the reference absorbance and As represents the sample absorbance.

2.3 Characterization of siderophores

Hydroxamate

Tetrazolium salt test: A small amount of tetrazolium salt and 1-2 drops of 2N NaOH were added to 0.1 mL of the test culture supernatant. The immediate appearance of a red to deep-red color indicated the presence of hydroxamate siderophores (Rashmi et al., 2013).

Catecholate

Arnow's test: 1 mL of the culture supernatant, 0.1 mL of 0.5 M HCl is added, followed by 1 mL of nitrate molybdate reagent and 1 mL of NaOH. The presence of catecholates in the supernatant is indicated by a color change from yellow to bright orange-red after 5 minutes of incubation (Arnow 1937).

Carboxylate

Vogel's chemical test: Prepare a test reagent by adding 3 drops of 2N NaOH to 1 drop of phenolphthalein. Add distilled water until the solution turns pink. Add the test sample to this solution. If the pink color disappears, it indicates the presence of carboxylate siderophores.

2.4 Effect of different medium on siderophore production

Bacillus subtilis was separately inoculated into nutrient broth, Luria-Bertani (LB) medium, *Bacillus* medium broth and succinate medium. Each medium was inoculated with a loopful of bacteria and incubated at 37°C for 48–72 hours in a rotary shaker set at 120 rpm. After incubation, the cultures were centrifuged to obtain the culture filtrate. 1 ml of cell filtrate was mixed with 1 ml of CAS (Chrome Azurol S) solution. The mixture was thoroughly mixed, and the absorbance was measured at 630 nm using a spectrophotometer. The percentage of siderophore units was then calculated based on the absorbance readings.

2.5 Optimization of the Cultural Conditions for Siderophore Production

One factor at a time (OFAT) was conducted to investigate the impact of various physicochemical parameters—including pH, carbon source, nitrogen source, and heavy metal, iron concentration on siderophore production of the UP11 strain.

Effect of carbon and nitrogen sources

The effects of different carbon sources including sucrose, glucose, maltose, sucrose, fructose, lactose, and mannitol as well as different nitrogen sources such as sodium nitrate, ammonium sulfate, soybean meal powder, yeast extract, ammonium nitrate, peptone, and potassium nitrate were investigated in succinate broth (100ml) by supplemented with 1 g/l of these different carbon and nitrogen sources separately. Then flasks were incubated at 37°C for 48–72 hours at 120 rpm. After incubation, 1 ml of the culture filtrate was mixed with 1 ml of CAS solution, and the absorbance was measured at 630 nm to calculate the percentage of siderophore units (Waday et al., 2022; Cornu et al., 2022).

Effect of pH and heavy metals

The pH of 100 ml of succinate broth was adjusted to 3.0, 5.0, 7.0, 9.0, and 11.0. To investigate the effect of metal ions, the succinate broth was supplemented with 10 μM of HgCl₂, ZnSO₄, MnCl₂, CdCl₂, MnSO₄, and NiCl₂ incubated at 37°C for 48–72 hours. The percentage of siderophore units was measured at 630 nm (Wang et al., 2021).

Effect of iron concentration

Log-phase bacterial cultures were separately inoculated into succinate broth containing varying concentrations of FeCl₃·6H₂O (0, 25, 50, 100, and 150 μM) and incubated at 37°C for 48–72 hours at 120 rpm to identify threshold level that inhibits siderophore production. Siderophore production was calculated as a percentage of siderophore units (Patel, 2022).

2.6 Preliminary Purification of Siderophore

After a 48-hour incubation at 30°C and 200 rpm, the liquid culture was centrifuged at 1,000 ×g for 10 minutes to collect the cell-free supernatant. It was tested for siderophores using the CAS assay. This supernatant was further purified by extraction with ethyl acetate. The pH of the supernatant was adjusted to 2 - 2.5 using strong acids before being quickly subjected to ethyl acetate extraction. The organic fractions were collected and dried by evaporation. The resulting dried siderophore crystals were suspended in sterile double-distilled water. The purified siderophore solutions were subsequently used in antifungal experiments.

2.7 Antifungal activity of purified siderophore

The purified siderophore was tested for antifungal activity against foliar blight fungal plant pathogens, *Alternaria triticina* and *Bipolaris sorokiniana* on potato dextrose agar. The fungal mycelium (5mm) were placed onto center then 5 mm wells were prepared. These wells were then filled with the purified siderophore and incubated at 28°C ± 2°C for 6 – 7 days. After the incubation, fungal growth inhibition was observed.

2.8 Seed germination

To evaluate the effect of bacterial siderophores on seed germination, wheat seeds were collected from a Norman E. Borlaug Crop Research Centre U.S. Nagar, Uttarkhand. Healthy seeds of similar sizes and shapes were selected for the experiments. The seeds were surface-sterilized by soaking in 75% ethanol, repeatedly washing with sterile deionized water to remove any chemicals, and drying with tissue paper. Fifteen seeds were soaked in the cell-free supernatant of the siderophore production medium in a Petri dish and incubated at 25°C for 1-2 hours. Seedlings were monitored and considered germinated when the root protruded at least 2 mm from the seed coat. Seedlings per plate were randomly selected to measure root and shoot lengths. The germination percentage was calculated using Equation (Kumar et al., 2018).

Germination percentage (% GP) = (Seeds germinated/Total seeds) × 100

2.9 Statistical analysis

All experiments were conducted in triplicates. Statistical analysis of the data was done using Statistical Package for Social Studies (SPSS), version 21.0. software. All data are expressed as means ± SE.

3. Results

3.1 Qualitative and Quantitative estimation of siderophore

Clear halo zone was observed on chrome-azurool S-agar medium (CAS) agar medium shows the strain UP11 was siderophore positive. CAS-Fe⁺³-HDTMA complex was broken by siderophores and formed Fe⁺³-siderophore complex. As dye released, resulted in clear zone formation (Murakami et al., 2021). The quantitative estimations were found to produce siderophores in the range of 53.26 ± 0.72 siderophore units (Fig. 1).

a





Figure1: Qualitative and Quantitative estimation of siderophore of *Bacillus subtilis* (a) Chrome-azurool S-agar medium plate assay for siderophore production (b) % Siderophore unit of UP11

3.2 Identification of type of siderophore production

The immediate appearance of a red to deep-red color confirmed the presence of hydroxamate siderophores. *B. subtilis* had a strong reaction to tetrazolium tests and Arnow's, which indicate the presence of a hydroxamate- catechol type siderophore in the supernatant of the culture. this indicating the occurrence of a mixed-type siderophore (both catechol and hydroxamate). The no disappearance of the pink color upon adding the test sample to the prepared reagent confirmed the absence of carboxylate siderophores (Table 1, Fig. 2).

Table1: Characterization of siderophore type in endophytic strain UP11 (*Bacillus subtilis*)

Strain	Peak at 425 nm (Hydroxamate)	Peak at 495 nm (catechol)	Hydroxamate (Tetrazolium test)	Catechol (Arnow's test)	Carboxylate (Vogel's test)
UP11	+	+	++	+	-

Catechol, carboxylate and hydroxamate-type siderophores were identified using Arnow', Vogel's and Tetrazolium test tests, respectively. ++ strong positive result; + positive result; – negative result




Figure 2: Positive results of different types of siderophore produced by *Bacillus subtilis*.

3.3 Influence of different media on siderophore production

Bacillus subtilis shows the maximum % siderophore unit in succinate medium (67.20%) followed upto on *Bacillus* broth (55.06%), LB (50.37%), NB (47.82%). Before optimizing physicochemical parameters, it is crucial to optimize the nutrient media to ensure maximum growth of the test culture and, consequently, maximum siderophore production. Among the four different nutrient media tested, succinate medium yielded the highest siderophore production at 630 nm (Fig. 3). Similarly, a study reported that succinate B medium to be ideal for siderophore production by *P. aeruginosa* KSB, *P. aeruginosa* PAUT14, *P. aeruginosa* PB19, *B. amyloliquefaciens*, *Bacillus subtilis* MF497446, and *Pseudomonas koreensis* (Bindu and Nagendra 2016; Srivastava et al., 2022; Ghazy and El-Nahrawy 2021). So, succinate media is further used for optimization culture condition for siderophore production.




Figure 3: Effect of different media on % Siderophore units of *Bacillus subtilis* strain. Values are presented as mean \pm SE (n = 3).

3.4 Optimization of culture condition for siderophore production

The literature indicates that several factors, such as initial pH, carbon source, nitrogen source, iron concentration and metal ions, influence siderophore synthesis and secretion (Srivastava et al., 2022). Additionally, these conditions can vary significantly depending on the bacterial species and strain used to

optimize siderophore production (Sheng et al., 2020). Therefore, the optimum pH 7 was found for maximum siderophore unit (62.95%), mannitol as carbon source (59.14%), ammonium nitrate as nitrogen source (63.04%), heavy metal $Pb(NO_3)_2$ shows (61.36%) highest yield in the succinate broth. The optimal iron concentration $50 \mu M$ for siderophore production. All these parameters and their effects were investigated using the one-factor-at-a-time (OFAT) approach (Fig. 4).




Figure 4. Effect of physicochemical parameters (a) carbon (b) nitrogen sources, (c) initial pH (d) heavy metals (e) iron concentration on siderophore production of *Bacillus subtilis*

3.5 Antifungal Activity of Crude Siderophore filtrate

Siderophore produced by UP11 has a stronger antifungal effect shows fungal mycelial inhibition (59.64 % ,50.88%) against *Alternaria triticina* and *Bipolarissorokiniana* (Fig. 5). These results indicated that siderophore is a crucial component produced by UP11 and plays a significant role in antifungal activity. This effect is due to siderophores sequestering iron, depriving fungi of this essential nutrient and inhibiting their growth. Additionally, siderophores may directly interfere with fungal iron acquisition systems and enhance plant defenses, making plants more resistant to fungal infections (Ghazy and El-Nahrawy 2021).




Figure 5: Antagonistic activity of purified siderophore extract of UP11 against *Alternaria triticina* (F1) and *Bipolarissorokiniana* (F2).

3.6 Seed Germination

This study investigated the effect of siderophores on the germination of wheat seeds. Seeds were incubated in siderophore-positive supernatant, with water-incubated seeds as controls. The siderophore presence was confirmed via the CAS assay, only siderophore-positive supernatant was used. Germination percentage was assessed after 36 hours, with sprouting monitored and root length measured every 12 hours. *Bacillus subtilis* enhanced seedling growth i.e., root, shoot length and germination percentage for each seed with the respective supernatant are shown in the figure 6. The siderophore positive *Bacillus subtilis* ethyl acetate filtrate shows the maximum germination root (3.60 cm) and shoot length (4.17 cm).

a






Figure 6: In vitro seed bioassay to check the effect of partial purified siderophore on seeds (a) control (water) (b) Ethyl acetate extract of siderophore (c) percent germination (d) Shoot and root length. Values are presented as mean \pm SE (n = 3).

Conclusions

Iron is an essential and abundant metal on Earth's surface, but it is unavailable to living organisms at physiological pH. Under iron-limiting conditions, siderophore-producing microorganisms can act as bio-fertilizers to enhance wheat growth. Iron deficiency hampers growth and impairs photosynthesis. *Bacillus* sp. WR12 has been shown to improve iron acquisition and mitigate these adverse effects (Yue et al., 2022). Our study found that *Bacillus subtilis* as an effective siderophore producing bacterium and it produces mixed siderophore types that enhance the antifungal as well as elevated the plant growth of wheat. So, it induces the disease resistance of foliar blight disease in wheat plant and enhancement of growth promotion and yield. Further research is needed to explore gene-level interactions related to siderophore production in plant growth-promoting rhizobacteria (PGPR), pathogenic fungi, and host plants.

Author Contributions

N.S: Conceptualization, methodology, preparation of experiments, analysis of data, preparation of final draft of manuscript, V.B and S.S: writing—review and editing M.S.: supervision reviewing, validation of data. All authors have read and approved the final version of the manuscript.

Funding

This research received no external funding.

Data Availability Statement

All the data are presented in the paper.

References:

Albelda-Berenguer, M., Monachon, M., & Joseph, E. (2019). Siderophores: From natural roles to potential applications. *Advances in applied microbiology*, 106, 193-225.

Arnou, B. Y. L. E. (1937). Hydrochloric Acid and Enough Distilled Water to Make a Volume of 1 Liter. Preserve under Toluene. 531. *Comp. A J. Comp. Educ.*, 531-537.

Basu, A., Prasad, P., Das, S. N., Kalam, S., Sayyed, R. Z., Reddy, M. S., & El Enshasy, H. (2021). Plant growth promoting rhizobacteria (PGPR) as green bioinoculants: recent developments, constraints, and prospects. *Sustainability*, 13(3), 1140.

Bendale, M. S., Chaudhari, B. L., & Chincholkar, S. B. (2009). Influence of environmental factors on siderophore production by *Streptomyces fulvissimus* ATCC 27431. *Current Trends in Biotechnology and Pharmacy*, 3(4), 362-371.

Bindu, P., & Nagendra, P. G. (2016). Siderophore production by *Pseudomonas aeruginosa* isolated from the paddy fields of Kuttanad, Kerala. *Int J Sci Res*, 5, 1577-1581.

Chowdappa, S., Jagannath, S., Konappa, N., Udayashankar, A. C., & Jogaiah, S. (2020). Detection and characterization of antibacterial siderophores secreted by endophytic fungi from *Cymbidium aloifolium*. *Biomolecules*, 10(10), 1412.

Cornu, J. Y., Gutierrez, M., Randriamamonjy, S., Gaudin, P., Ouedraogo, F., Sourzac, M., ... & Janot, N. (2022). Contrasting effects of siderophores pyoverdine and desferrioxamine B on the mobility of iron, aluminum, and copper in Cu-contaminated soils. *Geoderma*, 420, 115897.

Ghazy, N., & El-Nahrawy, S. (2021). Siderophore production by *Bacillus subtilis* MF497446 and *Pseudomonas koreensis* MG209738 and their efficacy in controlling *Cephalosporium maydis* in maize plant. *Archives of microbiology*, 203(3), 1195-1209.

Gregory, J. A., & David, M. F. (2017). Current understanding of iron homeostasis. *Am. J. Clin. Nutr*, 106, 1559S-1566S.

Handore, A. V., Khandelwal, S. R., Karmakar, R., & Handore, D. V. (2022). Exploration of bacterial siderophores for sustainable future. In *Climate Change and Microbial Diversity* (pp. 163-189). Apple Academic Press.

Jaggi, V., Joshi, S., Dasila, H., Pareek, N., & Sahgal, M. (2020). Functional and molecular characterization of wheat rhizosphere bacteria and their antagonistic activity against wheat foliar blight pathogens.

Khan, A., Gupta, A., Singh, P., Mishra, A. K., Ranjan, R. K., & Srivastava, A. (2020). Siderophore-assisted cadmium hyperaccumulation in *Bacillus subtilis*. *International microbiology*, 23, 277-286.

Kumar, P., Thakur, S., Dhingra, G. K., Singh, A., Pal, M. K., Harshvardhan, K., ... & Maheshwari, D. K. (2018). Inoculation of siderophore producing rhizobacteria and their consortium for growth enhancement of wheat plant. *Biocatalysis and agricultural biotechnology*, 15, 264-269.

Leventhal, G. E., Ackermann, M., & Schiessl, K. T. (2019). Why microbes secrete molecules to modify their environment: the case of iron-chelating siderophores. *Journal of the Royal Society Interface*, 16(150), 20180674.

Malgioglio, G., Rizzo, G. F., Nigro, S., Lefebvre du Prey, V., Herforth-Rahmé, J., Catara, V., & Branca, F. (2022). Plant-microbe interaction in sustainable agriculture: the factors that may influence the efficacy of PGPM application. *Sustainability*, 14(4), 2253.

Murakami, C., Tanaka, A. R., Sato, Y., Kimura, Y., & Morimoto, K. (2021). Easy detection of siderophore production in diluted growth media using an improved CAS reagent. *Journal of Microbiological Methods*, 189, 106310.

Patel, N. (2022). Siderophores. *Practical Handbook on Agricultural Microbiology*, 351-359.

Rani, R., Kumar, V., & Gupta, P. (2022). Plant growth-promoting rhizobacteria-assisted bioremediation of toxic contaminant: recent advancements and applications. In *Microbial Biodegradation and Bioremediation* (pp. 327-341). Elsevier.

- Rashmi, V., ShylajaNaciyar, M., Rajalakshmi, R., D'Souza, S. F., Prabakaran, D., & Uma, L. (2013). Siderophore mediated uranium sequestration by marine cyanobacterium *Synechococcus elongatus* BDU 130911. *Bioresource technology*, *130*, 204-210.
- Santos, S., Neto, I. F., Machado, M. D., Soares, H. M., & Soares, E. V. (2014). Siderophore production by *Bacillus megaterium*: effect of growth phase and cultural conditions. *Applied biochemistry and biotechnology*, *172*, 549-560.
- Sayyed, R. Z., Badgajar, M. D., Sonawane, H. M., Mhaske, M. M., & Chincholkar, S. B. (2005). Production of microbial iron chelators (siderophores) by fluorescent *Pseudomonads*.
- Sayyed, R. Z., Chincholkar, S. B., Reddy, M. S., Gangurde, N. S., & Patel, P. R. (2012). Siderophore producing PGPR for crop nutrition and phytopathogen suppression. In *Bacteria in agrobiolgy: disease management* (pp. 449-471). Berlin, Heidelberg: Springer Berlin Heidelberg.
- Schwyn, B., & Neilands, J. (1987). Universal chemical assay for the detection and determination of siderophores. *Analytical biochemistry*, *160*(1), 47-56.
- Sheng, M., Jia, H., Zhang, G., Zeng, L., Zhang, T., Long, Y., ... & Liu, H. (2020). Siderophore production by rhizosphere biological control bacteria *Brevibacillus brevis* GZDF3 of *Pinelliaternata* and its antifungal effects on *Candida albicans*. *Journal of Microbiology and Biotechnology*, *30*(5), 689.
- Sheng, M., Jia, H., Zhang, G., Zeng, L., Zhang, T., Long, Y., ... & Liu, H. (2020). Siderophore production by rhizosphere biological control bacteria *Brevibacillus brevis* GZDF3 of *Pinelliaternata* and its antifungal effects on *Candida albicans*. *Journal of Microbiology and Biotechnology*, *30*(5), 689.
- Singh, P., Chauhan, P. K., Upadhyay, S. K., Singh, R. K., Dwivedi, P., Wang, J., ... & Jiang, M. (2022). Mechanistic insights and potential use of siderophores producing microbes in rhizosphere for mitigation of stress in plants grown in degraded land. *Frontiers in Microbiology*, *13*, 898979.
- Srivastava, P., Sahgal, M., Sharma, K., Enshasy, H. A. E., Gafur, A., Alfarraj, S., ... & Sayyed, R. Z. (2022). Optimization and identification of siderophores produced by *Pseudomonas monteillii* strain MN759447 and its antagonism toward fungi associated with mortality in *Dalbergia sissoo* plantation forests. *Frontiers in Plant Science*, *13*, 984522.
- Tailor, A. J., & Joshi, B. H. (2012). Characterization and optimization of siderophore production from *Pseudomonas fluorescens* strain isolated from sugarcane rhizosphere. *Journal of Environmental Research and Development*, *6*(3A), 688-694.
- Trindade, I. B., Hernandez, G., Lebègue, E., Barrière, F., Cordeiro, T., Piccioli, M., & Louro, R. O. (2021). Conjuring up a ghost: structural and functional characterization of FhuF, a ferric siderophore reductase from *E. coli*. *JBIC Journal of Biological Inorganic Chemistry*, *26*, 313-326.
- Waday, Y. A., Girma Aklilu, E., Bultum, M. S., Ramayya Ancha, V., & Beyene, D. (2022). Isolation and Characterization of Plant Growth Promoting Rhizobacteria from Coffee Plantation Soils and Its Influence on Maize Growth. *Applied and Environmental Soil Science*, *2022*(1), 5115875.
- Wang, H., Liu, R., You, M. P., Barbetti, M. J., & Chen, Y. (2021). Pathogen biocontrol using plant growth-promoting bacteria (PGPR): Role of bacterial diversity. *Microorganisms*, *9*(9), 1988.

Wang, Y., Huang, W., Li, Y., Yu, F., & Penttinen, P. (2022). Isolation, characterization, and evaluation of a high-siderophore-yielding bacterium from heavy metal-contaminated soil. *Environmental Science and Pollution Research*, 29, 3888-3899.

Yasmin, S., Alcazar-Fuoli, L., Gründlinger, M., Puempel, T., Cairns, T., Blatzer, M., ... & Haas, H. (2012). Mevalonate governs interdependency of ergosterol and siderophore biosyntheses in the fungal pathogen *Aspergillus fumigatus*. *Proceedings of the National Academy of Sciences*, 109(8), E497-E504.

Yu, S., Teng, C., Bai, X., Liang, J., Song, T., Dong, L., ... & Qu, J. (2017). Optimization of siderophore production by *Bacillus* sp. PZ-1 and its potential enhancement of phytoextraction of Pb from soil. *Journal of Microbiology and Biotechnology*, 27(8), 1500-1512.

Yue, Z., Chen, Y., Hao, Y., Wang, C., Zhang, Z., Chen, C., ... & Sun, Z. (2022). *Bacillus* sp. WR12 alleviates iron deficiency in wheat via enhancing siderophore-and phenol-mediated iron acquisition in roots. *Plant and Soil*, 471(1), 247-260.

UNDER PEER REVIEW