

Bacteriological Evaluation of Locally Prepared Smoothies Formulated from Different Fruit Combinations.

ABSTRACT

The pharmaceutical benefits of fruits to humans is unfolding daily, thus, ensuring its Microbiological quality for the safety of final consumers is of paramount importance. Smoothie combinations made of paw-paw, banana and bene-seed (PPBNBS), orange, water melon and banana (ORWMBN), orange, banana and pineapple (ORBNPN) and paw-paw, watermelon and banana (PPWMBN) were prepared and analyzed as fresh (analyzed immediately after preparation), room temperature (analyzed after 24 hrs of preparation on bench) and refrigerated (kept in the refrigerator for 24 hrs after preparation). The mean colony count for refrigerated smoothie drink was $1.0-9.0 \times 10^5$ Cfu/mL, $1.0-8.0 \times 10^5$ for smoothie kept at room temperature and $1.0-5.0 \times 10^5$ Cfu/mL for freshly analyzed drink. The high mean count observed among the refrigerated drink may be interpreted as high concentration of psychrophilic organisms. Nine (9) genera of Microorganisms were isolated and identified. *Rothia*, *Micrococcus*, *Paenibacillus*, *Klebsiella* and *Lactobacillus* had similar 11.11 % rate of occurrence. *Enterococcus* sp were 22.22 % abundant. *Bacillus* and *Staphylococcus* sp both had 33.3 % occurrence while 44.4 % being the most abundant was recorded for *Streptococcus* sp. Among the identified organisms, Gram positive were the most frequent isolates occurring 8 (88.9%) compared to 1 (11.11%) of Gram negative organisms. The distribution of isolated organisms were not influenced ($P=0.05$) by the fruits combinations of smoothie neither the analyzed conditions. Although, the orange, banana and pineapple (ORBNPN) combination tend to have more 40% isolates occurrence compared to other three (PPBNBS, PPWMBN and ORWMBN) with even distribution of isolates at 20% occurrence rate. Though *Streptococcus* sp were the most abundant 44.4 % strain of isolated organisms, *Staphylococcus* sp were presence in all smoothies' combination. The results of the index study report higher than acceptable colony count in fruit drink according to the Microbiological Criteria (GULF standard, 2000). And the isolated organisms are prominent indicators pathogens of foodborne infection. This therefore implies that, smoothie's drink that are not properly handled during preparation can serve as a potential threat and a salience vehicle for foodborne intoxication and possible outbreak.

Key words: Foodborne indicators pathogens, *Klebsiella*, *Staphylococcus*, *Bacillus* and *Streptococcus*, Smoothies drink, different fruits combinations, Microbiological quality.

Comment [BG1]: The context must be clearly defined and enrich the bibliography.

1. INTRODUCTION

Fruits are great sources of phytochemicals essential for human health and are consumed as an important component of a healthy human diet. Their beneficial effects are based on the composition of many vitamins, minerals, dietary fiber, phytochemicals and their protective role against diseases such as diabetes, obesity, hypertension or coronary heart disease [1-4]. Recently, fresh fruits, vegetables as well as smoothies is in high demand. This is because consumption of fruits, fresh and healthy food is a preferred healthier lifestyle of most populace.

Smoothies are thick beverage products prepared from a wide range of fruits and vegetables using a blender. It may include other ingredients such as water, crushed ice, fruit juice and sweeteners (such as honey, sugar, syrup), dairy products (such as milk, yoghurt, low fat or cottage cheese), plant milk (such as coconut milk, tiger nut milk, almond nut milk, soy milk), seeds (such as celery seeds), spices (such as ginger, garlic), tea, chocolate, herbal supplements and nutritional supplements [5]. In some developing Countries like Nigeria, smoothies are becoming a preferred choice of drink and are commonly prepared on demand and sold in big shops, hotels and other relaxation spots [6]. Silha *et al.*, 2022 [7] reported that smoothie drinks sold in fast food establishments of Eastern Bohemia are potential source of microbial pathogens. According to the study of Krahulcova *et al.*, 2021 [8], antibiotic-resistant coliforms bacteria proliferate more in green smoothies and juicy formulations containing more vegetable ingredient compare to others. None of the above mentioned studies on microbial safety of freshly made smoothies were carried out in Cross River State. Whereas, smoothie formulations of different fruit combinations are served in 50-70% homes, 50-80% renounced hotels and seat out-joints. Therefore the major objective of this paper is to evaluate and report the bacteriological contents of locally prepared smoothies formulated with different fruit combination which is limited in literature especially in Cross River State.

2. MATERIALS AND METHODS

2.1 Sample Collection and Preparation

The fruit samples used in this study were orange, banana, watermelon, bene-seed, pineapple and pawpaw. They were purchased within EkpoAbasi junction fruit market, Calabar south.

Samples were aseptically wrapped using a sterile polythene bag and conveyed immediately after purchased to Microbiology Laboratory, University of Cross River State. Different fruits combinations and milk were used as smoothie compositions in this study ((pawpaw, banana, liquid milk and been-seed), (orange, banana and pineapple), (orange, watermelon and banana) and (pawpaw, watermelon and banana))

The fresh fruits were sorted, thoroughly washed with running tap water, peeled with sterile knife, diced into smaller pieces and weighed on a weighing balance (Digital compact scale; Atom A123). 200 grams weight of each fruit item was homogenized using a Moulex juicer blender (Model JB-70B) with 400mL of water. The blending process was allowed for 5 minutes, and thereafter was turned into sterile bottles as stock contents.

The smoothie stock was divided into three different portions, one was analyzed fresh, the second portion was allowed at room while the third one was kept under refrigerating

Comment [BG2]: The methodology is very poor and deserves to be totally rewritten.

1) The fruits such as oranges used in this study, in what form did you use them? Don't forget that oranges contain seeds.

2) This study is oriented towards bacteriological evaluation, but nowhere in the method do you describe the various activities involved.

temperature and both were analyzed after 24 hours. 1mL of each homogenate was taken and suspended in 9 mL of distilled water and diluted down to 10^{-10} . 1mL diluent of 10^{-5} and 10^{-6} were pour-plated in already prepared molten semi-solid nutrient (Chaitanya, RDM-NA-01) and MacConkey (Chaitanya, RDM_MCA-02) agar respectively.

2.2 Data Analysis

The student unpaired T-test was used to compare and determine the significant bacterial mean count of different smoothie combinations at different analyzed conditions at $P=0.05$. Microsoft Excel 2006 was used for the pictorial representation of: Pie, Bar and Line graphs for better separation and understanding.

3. RESULTS

3.1 BACTERIAL CELL COUNT

The fresh portion of smoothie combinations made of pawpaw, banana and bene-seed had bacterial cell count of 2×10^5 CfU/mL reported only from Nutrient agar plate.

The colony count from room temperature ranged from 4.4 to 2.0×10^5 CfU/mL while that of refrigerating temperature range 8.0 to 6.0×10^5 CfU/mL.

Orange, banana and pineapple smoothie formulation yielded 1.0 to 2.0×10^5 CfU/mL for freshly prepared. The room temperature colony count was 3.0 to 7.0×10^5 CfU/mL while the refrigerated recorded 1.4 to 6.0 CfU/mL respectively.

Banana, watermelon and orange fresh smoothie combinations yielded 1.3 to 5.0×10^5 CfU/mL. Room temperature bacteria cell count was 1.8 to 8.0×10^5 CfU/mL, refrigeration, 9.0×10^5 CfU/mL.

Pawpaw, watermelon and banana fresh portion had 3.0×10^5 CfU/mL. Room temperature and refrigerated had 2.0 to 6.0×10^5 CfU/mL and 1.0 to 4.0×10^5 respectively (Table 1).

3.2 Isolated and Identified Microorganisms

Streptococcus sp, *Bacillus* sp, , *Staphylococcus* sp, *Enterococcus* sp, *Rothia* sp, *Micrococcus* sp, , *Paenibacillus* sp, *Lactobacillus* sp and *Klebsiella* sp were the nine prominent genera of microorganisms isolated and identified in this study with respective frequencies of 44.4 % for *Streptococcus*, 33.3 % for *Bacillus* and *Staphylococcus* and 22.22 % for *Enterococcus*. *Rothia*, *Micrococcus*, *Paenibacillus*, *Lactobacillus* spp all had percentage occurrence of 11.11 % respectively (Fig. 1). The occurrence rate of Gram positive isolates was higher 8(88.9 %) compare to 1 (11.11 %) for Gram negative organisms.

3.3. Variations in Isolated Organisms According to Smoothie Combinations

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Comment [BG3]: Is the percentage attributed to each isolated microorganism taken independently, or is it part of a sub-part of the whole? If it's a sub-part, how do you explain that the sum is greater than 100? But if it's taken independently, how did you calculate it?

The research records variations in the distributions of isolated organisms among smoothie formulations. The different analyzed conditions such as; fresh, room temperature and refrigerated had no influence on the frequency of isolated organisms. However, the freshly analyzed smoothie made from combination of ORBNPN had 6(60.0 %) followed by 3 (60.0 %) from ORWMBN while PPBNBS and PPWMBN combinations had 1 (20.0 %) isolated organisms.

Formatted: Highlight

Comment [BG4]: Based on these different figures, the values indicate a total of 10 microorganisms, whereas the previous results show 9. As for the percentages in brackets, I wonder how you calculated them..

The percentage occurrence of isolated organisms from different smoothie combinations analyzed at room temperature (RM) were 1(20.0 %) for both ORWMBN and PPWMBN, 3 (30.0 %) for ORBNPN and 2 (40.0 %) for PPBNBS respectively.

The refrigerated smoothie combinations of ORBNPN, ORWMBN, PPBNBS and PPWMBN yielded 10.0 %, 20.0 %, 40.0 %, and 40.0 % percentage frequencies of isolated organisms (Fig.2).

Comment [BG5]: The sum here gives 110%.

3.4 Sum-total of Isolates based on Gram Reactions

The study reports higher frequency 8 (88.9 %) of Gram positive organisms compared to 1 (11.1 %) Gram negative bacterial isolates (Fig. 3)

3.5 Sum-total of Isolates According to Smoothie Combinations and Conditions

The smoothie combination; ORWMBN, PPWMBN and PPBNBS fresh, room temperature and refrigerated conditions had similar 5 (20.0 %) isolates occurring rate while ORBNPN had the highest 10(40.0 %) percentage isolates (Fig. 4).

Table 1: Mean Bacterial Cell Counts in Colony Forming Unit (Cfu/mL) from Various Smoothies Analyzed Conditions

	Fresh	Room Temp	Refrigerated
ORWMBN	1.3-5.0 x 10 ⁻⁵ Cfu/mL	1.8-8.0 x 10 ⁻⁵ Cfu/mL	1.0-9.0 x 10 ⁻⁵ Cfu/mL
PPWMBN	1.0-3.0 x 10 ⁻⁵ Cfu/mL	2.0-6.0 x 10 ⁻⁵ Cfu/mL	1.0-4.0 x 10 ⁻⁵ Cfu/mL
ORBNPN	1.2-2.0 x 10 ⁻⁵ Cfu/mL	3.0-7.0 x 10 ⁻⁵ Cfu/mL	1.4-6.0 x 10 ⁻⁵ Cfu/mL
PPBNBS	1.3-2.0 x 10 ⁻⁵ Cfu/mL	2.0-4.4 x 10 ⁻⁵ Cfu/mL	6.0-8.0 x 10 ⁻⁵ Cfu/mL

Comment [BG6]: You get bacterial colony numbers followed by negative exponents. Are you sure of these values?

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Keys: ORWMBN = Orange, watermelon, Banana. PPWMBN = Pawpaw, watermelon and Banana. ORBNPN = Orange, Banana and pineapple. PPBNBS = Pawpaw, Banana and pineapple.

Table 2: Distribution of Isolates According to Different Smoothies Combinations Analyzed Under Different (Fresh, Room temperature and Refrigerated) Conditions.

ISOLATED ORGANISMS	ORWMBN			PPWMBN			ORBNPN			PPBNBS			TOTAL NO. OF APPEARANCE
	F	RM	R	F	RM	R	F	RM	R	F	RM	R	
<i>Bacillus sp.</i>	0.00	0.00	4.00	0.00	0.00	0.00	4.00	0.00	0.00	4.00	0.00	0.00	3
<i>Streptococcus sp.</i>	4.00	0.00	0.00	0.00	4.00	0.00	0.00	0.00	4.00	0.00	0.00	8.00	5
<i>Rothia sp.</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.00	0.00	1
<i>Micrococcus sp.</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.00	0.00	1
<i>Staphylococcus sp.</i>	8.00	4.00	0.00	0.00	4.00	12.0	8.00	4.00	0.00	0.00	0.00	0.00	9
<i>Paenibacillus sp.</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.00	0.00	0.00	0.00	0.00	1
<i>Klebsiella sp.</i>	0.00	0.00	0.00	0.00	0.00	0.00	8.00	0.00	0.00	0.00	0.00	0.00	2
<i>Lactobacillus sp.</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.00	0.00	0.00	0.00	0.00	1
<i>Enterococcus sp.</i>	0.00	0.00	0.00	4.00	0.00	0.00	4.00	0.00	0.00	0.00	0.00	0.00	2
Number of Times	3	1	1	1	1	3	6	3	1	1	2	2	25




Figure 1: Total number of bacteria Isolates and their percentage Occurrence in Different Smoothies Combinations.

Comment [BG7]: If these values are for each of the combinations, did you sum or average them? If you averaged them, where are the errors?




Fig.2: Different Smoothies Combinations and percentage frequencies of Isolates.




Fig.3: Sum-total of Isolates based of Gram Reactions




Fig 4: Sum-total of all Isolates According to Smoothies combinations

4. Discussion

The evaluation of bacterial quality of locally prepared smoothies formulated using four different fruits combinations (ORWMBN, PPWMBN, PPBNBS, ORBNPN) was carried out to ascertain the safety of smoothie drink to final consumers.

The cross sum of mean count for refrigerated smoothie was 1.0 to 9.0×10^{-5} Cfu/mL, 1.0 to 8.0×10^{-5} for room temperature and 1.0 to 5.0×10^{-5} Cfu/mL for freshly prepared. The presence of microorganisms in all the three analyzed smoothie conditions are unhealthy according to the microbiological criteria (GULF standard, 2000) [11]. However, high mean count observed among refrigerated smoothies may be interpreted as high concentration of psychrophiles.

Obasi and odoh [12] reported 1.3×10^{-4} and 8.0×10^{-3} Cfu/mL in smoothies made from two fruit combinations (pineapple and watermelon) in Makurdi, Benue state, Nigeria.

Bacillus was found in all smoothie combinations at frequency of 1 (4.0 %). The occurrence of *Streptococcus* sp was similar 1 (4.0 %) to that of *Bacillus* except that its occurrence in PPBNBS was higher 2 (8.0 %). *Rothia* sp and *Micrococcus* occurred just once 1 (4.0 %) in PPBNBS fruits combination. *Paenibacillus* and *Lactobacillus* sp occurred only in ORBNPN 1 (4.0 %). *Klebsiella* appeared only in ORBNPN at the rate of 2 (8.0 %) while *Enterococcus* sp had occurrence rate of 1 (4.0 %) in ORWMBN and ORBNPN respectively. *Staphylococcus* had the highest frequency of 9 (36.0 %) distributed at the rate of 2 (8.0 %), 1 (4.0 %) in ORWMBN and 1 (4.0 %), 3 (12.0 %), and 1 (4.0 %) in PPWM and in all analyzed conditions.

A total of 9 (36.0 %) of prominent strains of microorganisms were isolated and identified in this study. And comprised of; *Rothia*, *Micrococcus*, *Paenibacillus*, *Klebsiella*, *Lactobacillus*, *Enterococcus*, *Bacillus*, *Staphylococcus*, and *Streptococcus*. *Rothia*, *Micrococcus*, *Paenibacillus*, *Klebsiella* and *Lactobacillus* occurred within the same range of 11.11 %. *Enterococcus* was 22.22 %, *Bacillus* and *Staphylococcus* sp had similar occurrence rate of 33.3 % and *Streptococcus* sp was the most abundant isolate with percentage occurrence of 44.4 %.

Staphylococcus, *Bacillus* and *Klebsiella* are the most noticeable food borne pathogens isolated in this study. Their presence are serious public threat indicators and their major route of entry may be poor personal hygiene and handling during preparation. According to most published literature [12], *Staphylococcus aureus* is

Comment [BG8]: I've looked at this article (Organoleptic Properties and Microbial Quality of Smoothies Produced from Watermelon-Pineapple and Sweet Melon-Cucumber Blends), and the values you give are not in line with those obtained by these authors.

commonly found in smoothies formulated majorly from Watermelon, pineapple and sweet melon cucumber blends.

Streptococcus sp was the most abundant isolate in all smoothies' combinations. The report of Falkenhorst *et al.*, 2008 [13] pointed *Streptococcus* among the causative agents of food spoilage based of their outbreak experiment conducted on Group A *Streptococcus* species in Denmark.

Other microorganisms such as *Micrococcus*, *Rothia*, *Paenibacillus*, *Lactobacillus* and enterococcus occurred sparingly, however the potency in any pathogen recovery from diluent of 10^{-5} should not be undermined [14].

The index result reports higher frequency 8 (32.0 %) of Gram positive organisms compared to 1 (4.0 %) Gram negative bacterial isolates. This could be as a result of more gram positive bacteria being implicated in food poisoning [15].

The smoothies combination ORBNPN had the highest 10 (40.0 %) of isolated organisms to 5 (20.0 %) for ORWMBN, PPWMBN and PPBNBS respectively. Smoothie combinations have no significance output and or influence on bacterial occurrence, however, observing good hygiene practices during smoothie preparation is essential in reducing microbial load and ensuring safety of the final consumers.

Conclusion

The research report showed unacceptable bacterial colony mean count of 1.0-9.0, 1.0-8.0 and 1.0-5.0 x 10^{-5} Cfu/mL for smoothie analyzed after 24 hours of preparation and kept under refrigeration, the second kept on bench for 24 hours after preparation before analysis and the third analyzed fresh immediately after preparation. The isolated and identified organisms were majorly gram positive organisms of medical importance highly implicated in foodborne infections. They were: *Rothia*, *Micrococcus*, *Paenibacillus*, *Lactobacillus*, *Enterococcus*, *Bacillus*, *Staphylococcus*, and *Streptococcus* respectively. Improper handling and preparation of smoothie drink is a silence vehicle for the transmission of foodborne indicator pathogens. This implies that, the public and or the final consumers are the most at risk group. Hence, personal hygiene measures such as proper hand washing during food preparation, thorough washing of purchased fruits as well as using clean utensils during food preparation are important measures of microbial load reduction/infection mitigation.

REFERENCES

- Hall, J.N., Moore, S., Harper, S.B. & Lynch, J.W. (2009). Global Variability in Fruit and Vegetable Consumption. *Am. J. Prev. Med.*, 36, 402-409.
- Van Duyn, M. A.S., & Pivonka, E. (2000). Overview of the health benefits of fruit and vegetable consumption for the dietetics professional: selected Literature. *Journal of diet association*, 100, 1511-1521.
- Southon, S. (2000). Increased fruit and vegetable consumption within the EU: Potential health benefits. *Food Resolution International*, 33, 211–217.
- Oyebode, O.; Gordon-Dseagu, V.; Walker, A.; Mindell, J.S. (2014). Fruit and vegetable consumption and all-cause, cancer and CVD mortality: Analysis of Health Survey for England data. *Journal of Epidemiological Community Health*, 68, 856–862.
- Maksuda, M., Abu Torab, M.A., Rahim, M., Nazmu, H. & Shahjalal, H. K. (2016). Proximate and Water Soluble Vitamin Contents in Some Selected Bangladeshi Fruits and Vegetables. *Journal of Scientific Research & Reports*, 11(6), 1-8.
- Zavasta, T. (2009). Smoothie moves: Enjoy the benefit of green smoothies and puddings. *Raw Food and Hot Yoga*, pp. 39. In: <https://www.en.wikipedia.org/wiki/smoothie>; ISBN0-9742434-9-3. Accessed on 07-02-2019.
- Silha, D., Svarcova, K., Bajer, T., Kralovec, K., Tesarova, E., Mouckova, K., Pechalova, M. (2020). Chemical composition of natural hydrolytes and their antimicrobial activity on Arcobacter-Like Cells in comparison with other microorganisms. *Molecules*, 25, 5654.
- Krahulcova, M., Micajova, B., Olejnikova, P., Cverenkova, K., Birosova, L. (2021). Microbial safety of smoothie drinks from fresh bars collected in Slovakia. *Foods*, 10, 551.
- Sani, N.A.; Siow, O.N. Knowledge, attitudes and practices of food handlers on food safety in food service operations at the University Kebangsaan Malaysia (2014). *Food Control*, 37, 210–217.
- Sujeet K. M, & Vipin K. Fresh farm produce as a source of pathogens: A review. Research (2015). *Journal of Environmental Toxicology*, 9(2), 59-70.
- Gulf standards (2000). Microbiological criteria for foodstuffs, part 1. GCC, Riyadh, Saudi Arabia pp. 27-30.

Comment [BG9]: References in the text are numerical; these numbers must be included in the reference list.

Obasi, B. C., & Odoh, P.I. (2023). Organoleptic Properties and Microbial Quality of smoothies prepared from Watermelon-pineapple and sweet melon- cucumber Blends. *Greeener Journal of Agricultural Sciences*, 13(3), 186-197.

Falkenhorst, G., Bagdonaite, J., Lisby, M., Madsen, S. B., Lambertsen, K. E. P., & Molbak, K. (2008). Outbreak of Group A streptococcal throat infection: don't forget to ask about food. *Epidemiology infection*, 136(9), 1165-1171.

Cheesbrough, M. Microbiological tests: Biochemical test to identify bacteria. District laboratory practice in tropical countries. (2nd Edition) Cambridge: Cambridge University Press. 2006;62-70.

Humberto, H., Wilmer, E., Luera, P., Nayara, S., Anderson, S. (2014). Food Microbiology.

UNDER PEER REVIEW