

Physiological and Biochemical Basis of Thermotolerance in Selectively Fertilized Coconut Hybrids

ABSTRACT

Aims: Physiologically and biochemically assess selectively fertilized coconut hybrids for their tolerance to high temperature stress.

Study design: Two factorial Completely Randomized Design

Place and Duration of Study: Department of Plant Physiology, College of Agriculture, Vellayani, Trivandrum between 28th October 2023 to 28th December 2023.

Methodology: Six-month-old selectively fertilized hybrid seedlings of three coconut varieties, Kerasree, Keraganga, and Kerasankara, were cultivated in a polyhouse with a pre-set temperature of $40\pm 1^{\circ}\text{C}$ for two months. Various physiological and biochemical parameters were measured, including Relative Water Content, Chlorophyll Stability Index, Epicuticular Wax Content, Lipid Peroxidation Level, and antioxidant enzyme activity, all essential for assessing stress tolerance. The results were analyzed using analysis of variance, and the standard error was calculated accordingly.

Results: For the physiological parameters RWC and CSI, S.F hybrids varied significantly under high temperature from their corresponding N.F hybrids with average mean value of 88.53 % and 83.27% respectively. High temperature induced solute leakage was lowest in Kerasankara S.F with a mean value of 3.58 ± 0.63 and wax deposition values were on par for SF and N.F hybrids. MDA accumulation due to lipid peroxidation were significantly lowest in S.F hybrids with a mean value of $2.47\mu\text{gg}^{-1}$. Antioxidant enzymatic activity of S.F hybrids were remarkably higher with a significant percent increase of 97.1% for Peroxidase and 96.1% for SOD in Keraganga SF genotype.

Conclusion: Coconut hybrid accessions developed through pollen selection showed significant superiority over regular hybrids based on physiological and biochemical stress tolerance indices. Therefore, pollen selection followed by selective fertilization is a reliable, cost-effective, and time-saving approach for breeding stress-tolerant crops, including perennials.

Keywords: Selective fertilization, high temperature stress, Pollen selection, Stress tolerance

1. INTRODUCTION

The coconut palm, often referred to as the "tree of life," plays a vital role in the livelihoods of smallholders, providing cash income, nutrition, and materials. It primarily thrives in the humid tropics, located between 23°N and 23°S of the equator, and can grow at altitudes up to 600 meters above sea level for nut production. According to the FAO statistical database, global coconut production is about 63.6 million nuts, covering a planted area of 11.6 million hectares [1]. Optimal production requires well-distributed rainfall and an annual mean temperature of $27\text{-}29^{\circ}\text{C}$ [2]. A minimum temperature above 10°C promotes flowering, while temperatures below 10°C for a month cause nut fall. Conversely, temperatures above 40°C from April to July in the tropics reduce the functional leaf area index, dry matter production, and nut yield [3]

According to projections, the Earth's surface temperature would rise by 1.5°C to 11°C by 2100 as a result of an increase in greenhouse gases [4]. It is predicted that brief bursts of extreme weather, such as high temperatures, may occur more frequently, which will affect fruit set and production. It is imperative to develop types that can withstand high temperatures in order to maximize coconut production under these circumstances. In vitro pollen selection followed by selective fertilization is the most feasible and cost-effective approach. By applying the appropriate selective pressure during the

pollination and fertilization processes, only pollen grains that are resistant to the pressure will be able to germinate and fertilize the ovule. For a number of variables, including biotic [5] and abiotic stress tolerance [6], pollen genotype selection has shown promise.

Stephen et al.[7] developed water stress tolerance in coconut via invitro pollen selection. [8] evaluated the water stress tolerance of selectively fertilized coconut hybrids using a range of physiological and biochemical indices. The results showed that S.F hybrids fared better during dry spells. Therefore, the current study was carried out to confirm the effectiveness of the pollen selection technique in developing genotypes that are stress resistant and to investigate and assess the thermotolerance of the selectively fertilized coconut hybrids based on physiological and biochemical characteristics.

2. MATERIAL AND METHODS

2.1 Experimental Setup

To assess the resistance of the selectively fertilized coconut hybrids created through pollen selection to high temperature stress, a polyhouse-based experiment was carried out at the College of Agriculture, Vellayani from October 28, 2023, to December 28, 2023. In the polyhouse facility, where the temperature was set to a higher known value compared to the ambient condition, three varieties of hybrids—Kerasree (West Coast Tall x Malayan Yellow Dwarf), Keraganga (West Coast Tall x Ganga Bondam), and Kerasankara (West Coast tall X Chowghat Orange Dwarf)—were laid out under three replications. The hybrids were fertilized both normally and selectively. Selectively fertilized hybrids and normally fertilized hybrids used in this study were developed in a previous study by the author via pollen selection and conventional method of assisted pollination respectively. Six-month-old seedlings were sown in polybags of size 40x24x24cm filled with 1:2:1 potting mixture and maintained as per POP recommendation [9]

2.2 Physiological Evaluation

Following two months of high temperature stress ($40 \pm 1^\circ\text{C}$) on hybrid seedlings, an evaluation was performed for a range of physiological observations. Leaf samples were obtained from the third index leaf of the seedlings for each parameter that was taken into account.

2.2.1 Relative water content (Percentage)

A leaf's hydration status is assessed by comparing its actual water content to its maximum water-holding capacity when fully turgid (a process known as "relative turgidity," or relative water content, or RWC [10]. The sample's turgid weight was measured by immersing it in distilled water for three hours and then weighing it; the sample's dry weight was measured by drying the turgid leaf samples in a hot air oven at 70°C until they reached a constant weight.

RWC (%) was calculated using the formula:

$$\text{RWC} = (\text{Fresh weight} - \text{dry weight}) / (\text{turgid weight} - \text{dry weight}) \times 100$$

2.2.2 Membrane Integrity (Percentage of solute leakage)

The quality or state of the entire membrane in ideal condition is known as membrane integrity. Cell membrane thermostability can be effectively determined by measuring electrolyte leakage, which has been used as a direct heat injury indicator [11]. Using the following formula, the membrane integrity of leaf tissues was determined and expressed as a percentage of leakage.

$$\% \text{ leakage} = ((\text{EC}_b - \text{EC}_a) / \text{EC}_c) \times 100$$

where EC_a is the initial value of electrical conductivity caused by immediate leakage from leaf samples in distilled water, EC_b is the value of electrical conductivity caused by leakage from samples incubated for 30 minutes and EC_c is the value of electrical conductivity caused by leakage from samples treated in boiling water bath for 10 minutes.

2.2.3 Chlorophyll Stability Index(Percentage)

Plants' ability to withstand stress is indicated by the chlorophyll stability index, or CSI. A high CSI value indicates that the stress had little impact on the plants' chlorophyll content. Through improved chlorophyll availability, a higher CSI increases a plant's ability to withstand stress [12]. By calculating the ratio of the total chlorophyll content of the heated leaf samples to the control samples and multiplying the result by 100, CSI (%) was calculated.

$$\text{Chlorophyll Stability Index (\%)} = \left\{ \frac{\text{Total Chl. Content in Heated lot}}{\text{Total Chl. Content in Control lot}} \right\} \times 100$$

2.2.4 Epicuticular wax content (mg 10cm⁻²)

In order to protect plants from biotic stressors like excessive radiation, water loss, high temperatures, and drought, epicuticular wax (EW) coats aerial surfaces to form a barrier between the environment and the plant [13]. Leaf bits of 10cm² area was immersed in pre weighed beakers and chloroform solvent sufficient enough to immerse the leaf bits. Following a chloroform solvent wash, the wax that had been deposited on the leaf surfaces was collected by evaporating the solvent.

$$ECW \text{ (mg)} = W_2 - W_1$$

Where ECW is the Epicuticular wax content, W_1 = Initial weight of the beaker, W_2 = Final weight of the beaker.

2.3 Biochemical Evaluation

Evaluation for various biochemical observations were performed after two-month exposure of hybrid seedlings to high temperature stress. For all the parameters observed, leaf samples were taken from third index leaf of the seedlings

2.3.1 Malondialdehyde Content ($\mu\text{g g}^{-1}$)

The ROS-induced peroxidation of membrane lipid (measured by MDA content) is a sign of cellular damage brought on by stress. MDA content was determined by the Thiobarbituric acid (TBA) reaction as described by [14]. Absorbance of the assay extract were observed at 532 and 600 nm and MDA concentration ($\mu\text{g g}^{-1}$) was estimated by subtracting the non-specific absorption at 600 nm from the absorption at 532 nm.

2.3.2 Activity of enzymatic antioxidants

2.3.2.a Peroxidase (activity $\text{g}^{-1}\text{min}^{-1}$)

Plant cells' enzymatic defense system includes peroxidases, which initiate the transformation of H_2O_2 into water and oxygen [15]. The peroxidase activity in plants was estimated following Pyrogallol method described by [14]. One unit of peroxidase is defined as the change in absorbance/minute at 430nm.

2.3.2.b Superoxide dismutase (activity $\text{g}^{-1}\text{min}^{-1}$)

In plants, the enzyme SOD is directly related to stress, which initiates the first line of defense, converting O_2^- into H_2O_2 [16]. SOD activity of plants was quantified following the method described by [17]. One unit of enzyme activity is defined as the amount of enzyme that gave 50% inhibition of NBT reduction in one minute.

3. RESULTS AND DISCUSSION

The experimental data was statistically analyzed in two factorial completely randomized design using grapes Agri1 software [18]. Results for physiological and biochemical parameters are presented in Table 1 to 4 and Figures 1 to 3. In the following section, normally fertilized and selectively fertilized hybrids are referred as N.F and S.F respectively.

3.1 Relative Water Content (%)

The impact of ambient and high temperatures on the Relative Water Content (RWC) of N.F. and S.F. hybrid coconuts are illustrated in (Table 1). Under high temperature ($40 \pm 1^\circ\text{C}$), the mean values of RWC for S.F hybrids ranged from 85.73% to 94.67% and for N.F hybrids from 79.54% to 91.03%. Kerasankara S.F. distinguished itself significantly from the other hybrids with the highest mean value of 94.67%, a 15.9% increase over its N.F counterpart. At higher temperatures, the lowest RWC value (79.54%) was recorded in Kerasankara N.F. The highest RWC mean value (91.67%) under ambient conditions ($28 \pm 2^\circ\text{C}$) was found in Kerasankara N.F., which was comparable to the mean values of Kerasankara S.F. and Kerasree N.F. The Keraganga N.F., which differed significantly from its S.F. hybrid, had the lowest value. The RWC expresses the precise amount of water that a plant needs to reach full saturation, making it a valuable indicator of the state of the plant's water balance. Data on tomato cultivars also showed that, in stressed conditions, tolerant genotypes had significantly higher RWC than susceptible ones [19]. [20] observed higher leaf water status in selectively fertilized, drought-tolerant coconut genotypes.

Table 1: RWC of coconut hybrids under High and Ambient temperature conditions

Sl.No	Hybrid genotypes	Relative Water Content (%)					
		HT			AT		
		NF	SF	Percent difference	NF	SF	Percent difference
1.	Kerasree	91.03 ^b	85.73 ^c	5.8	90.35 ^{ab}	86.49 ^{cd}	4.3
2	Keraganga	85.22 ^c	85.19 ^c	0.04	82.50 ^e	83.33 ^e	0.9
3	Kerasankara	79.54 ^d	94.67 ^a	15.9	91.67 ^a	90.00 ^{ab}	1.82
Mean		85.26	88.53		88.17	86.60	

SE+	0.98	0.68
CD at 5%	2.95	2.03

CD=critical difference, SE=standard error, NF=Normally Fertilized, SF= Selectively Fertilized

3.2 Chlorophyll Stability Index(%)

The impact of both ambient and high temperatures on the Chlorophyll Stability Index (CSI) of N.F. and S.F. hybrid coconuts is displayed in Table 2. At high temperatures ($40\pm 1^{\circ}\text{C}$), the mean CSI for N.F and S.F hybrids is 80.04% and 83.27%, respectively. The Kerasree S.F genotype outperformed all other genotypes by a significant margin, and its NF hybrid had the highest CSI value (97.83%) in high temperatures. All S.F genotypes displayed greater CSI values in ambient conditions, except Keraganga, which displayed values comparable to its NF hybrid. A higher CSI makes a plant more resilient to stress by increasing the availability of chlorophyll. Chlorophyll resilience leads to greater dry matter production, higher productivity, and a higher photosynthetic rate in a salinity stressed rice plant. [12]. Similar information supporting the current investigation was shown for wheat cultivars [21]

Table 2: Chlorophyll Stability Index of coconut hybrids under High and Ambient temperature conditions

SI.No	Hybrid genotypes	Chlorophyll Stability Index(%)			
		HT		AT	
		NF	SF	NF	SF
1.	Kerasree	70.31 ^{de}	97.83 ^a	85.02 ^{de}	97.29 ^a
2	Keraganga	92.15 ^{ab}	65.19 ^c	92.68 ^{bc}	91.05 ^c
3	Kerasankara	77.65 ^{cd}	86.78 ^{bc}	72.43 ^t	82.49 ^e
Mean		80.04	83.27	83.38	90.28
SE+		3.05		1.53	
CD at 5%		9.15		4.58	

CD=critical difference, SE=standard error, NF=Normally Fertilized, SF= Selectively Fertilized

3.3 Epicuticular wax content (mg 10cm⁻²)

The impact of ambient and high temperatures on the Epicuticular wax content (ECW) of coconut hybrids is shown in Figure 1. In comparison to its N.F hybrid, the S.F hybrid of Keraganga displayed the highest wax content under high temperatures. Significantly, Kerasankara S.F had the highest ECW value in ambient conditions. Similar findings were reported in wheat cultivars [22] where significantly greater wax deposition was noted under higher temperature stress. The presence of ECW protects plants from environmental stresses and water loss [23].

3.4 Lipid peroxidation - Malondialdehyde content ($\mu\text{g g}^{-1}$)

Reactive oxygen species (ROS) including superoxide radical, hydrogen peroxide, hydroxyl radical, and singlet oxygen are produced in excess when there is heat stress. The lipid peroxidation caused by reactive oxygen species (ROS) generated in plant tissue under heat stress damages membranes and forms microscopic hydrocarbon fragments like malondialdehyde (MDA), one of the key markers of oxidative stress [24]. The amount of MDA accumulated in the N.F and S.F coconut hybrids at high and ambient temperatures is shown in Figure 2. All of the SF hybrids showed lower MDA levels than the corresponding N.F hybrids at higher temperatures, but the SF hybrids performed differently in terms of lipid peroxidation at ambient conditions. Mango cultivars exposed to high temperatures displayed higher MDA activity, according to studies on pollen germination [25].




Figure:1 Epicuticular Wax Content of Coconut hybrids under High and Ambient temperature conditions




Figure:2 Malondialdehyde Content of Coconut hybrids under High and Ambient temperature conditions

3.5 Activity of Peroxidase (g⁻¹min⁻¹)

Table 3 shows the impact of both ambient and high temperatures on the peroxidase activity of coconut hybrid seedlings. Under high temperatures, all genotypes of SF hybrids showed significant variation from their NF hybrids, with a mean value of 6.24 activity g⁻¹min⁻¹. In high temperatures, Kerasankara SF displayed the highest mean value of 11.13 activity g⁻¹min⁻¹, while Keranganga N.F recorded the lowest mean value of 0.12 activity g⁻¹min⁻¹. At high temperature, all of the SF hybrids showed a notable percentage increase in peroxidase activity (94.9,97.1,70.4%) (Figure 3). With a mean activity of 1.12g⁻¹min⁻¹, the S.F hybrids outperformed their N.F hybrids in ambient conditions, recording higher mean values. Peroxidase enzyme's activity was increased by high temperature stress, and it has been linked to the emergence of physiological injuries brought on by thermal stress in plants [26]. Plants increase their thermotolerance by recruiting the antioxidant enzymes superoxide dismutase, ascorbate peroxidase, catalase, glutathione reductase, and peroxidase [27].

Table 3. Peroxidase activity of coconut hybrids under High and Ambient temperature conditions

S.No	Hybrid genotypes	Peroxidase				
		HT		Percent Increase	AT	
		NF	SF		NF	SF
1.	Kerasree	0.18 ^f	3.53 ^d	94.9	0.56 ^{dc}	0.51 ^e
2.	Keraganga	0.12 ^f	4.07 ^c	97.1	0.25 ^f	0.99 ^c
3.	Kerasankara	3.30 ^e	11.13 ^a	70.4	0.49 ^e	1.87 ^a
Mean		1.2	6.24		0.43	1.12
SE±		0.04			0.04	
CD at 5%		0.12			0.12	

CD=critical difference SE=standard error NF=Normally Fertilized SF= Selectively Fertilized

3.6 Activity of Superoxide dismutase (activity $\text{g}^{-1}\text{min}^{-1}$)

Table 4 shows SOD activity in coconut hybrid genotypes affected by various temperature conditions. Under high temperatures ($40\pm 1^{\circ}\text{C}$), S.F hybrids demonstrated an average mean value of $5.49 \text{ activity g}^{-1}\text{min}^{-1}$ for SOD activity, with Keraganga demonstrating a 96.1 % increase in enzyme activity (Figure 3) in comparison to its N.F counterpart. In comparison to their N.F hybrids, none of the N.F hybrids displayed as much SOD activity at high temperatures. The enzyme activity's highest mean value ($7.41 \text{ activity g}^{-1}\text{min}^{-1}$) was found in Kerasankara S.F, while Keraganga S.F had the lowest mean value ($0.22 \text{ activity g}^{-1}\text{min}^{-1}$). According to [16] the enzyme SOD in plants is directly linked to stress and starts the first line of defense by changing $\text{O}_2^{\bullet-}$ into H_2O_2 . Tomato genotypes that can withstand higher temperatures have been shown to have higher SOD sensitivity [28].

Table 4. Superoxide Dismutase activity of coconut hybrids under High and Ambient temperature conditions

Sl.No	Hybrid genotypes	SOD				
		HT		Percent increase	AT	
		NF	SF		NF	SF
1.	Kerasree	0.36 ^f	3.47 ^c	89.6	0.44 ^e	3.58 ^b
2.	Keraganga	0.22 ^f	5.59 ^b	96.1	0.22 ^f	5.26 ^a
3.	Kerasankara	1.54 ^e	7.41 ^a	79.2	0.38 ^{ef}	2.79 ^c
Mean		0.71	5.49		0.35	3.88
SE±		0.12			0.07	
CD at 5%		0.35			0.21	

CD=critical difference SE=standard error NF=Normally Fertilized SF= Selectively Fertilized






Figure: 3 Percent difference of Peroxidase and SOD activity in coconut hybrids under high temperature

4. CONCLUSION

All genotypes of SF hybrids exhibited superior performance in the current study for all physiological parameters associated with stress tolerance, including epicuticular wax content, chlorophyll stability index, and relative water content. S.F hybrids outperformed their N.F hybrids in mean values for biochemical observations such as lipid peroxidation and enzymatic antioxidant activity. Together, these indices confirm that S.F. hybrids are more temperature-tolerant. In order to create high temperature tolerant hybrids in coconuts, the current evaluation experiment thus validates the effectiveness of the pollen selection technique followed by selective fertilization.

REFERENCES

1. FA OSTAT,2021. <https://www.fao.org/faostat>
2. Rajagopal V, Parthasarathy V., Naresh Kumar S, Reddy DVS, Iyer R. Coconut. In: Parthasarathy VA, Chattopadhyay PK, Bose TK, editors. Plantation Crops, Naya Udyog Pub., Kolkata, India. 2006;(2):1–178
3. Kumar NS, Bai KKV, Rajagopal V, Aggarwal PK. Simulating coconut growth, development and yield using Info Crop-coconut model. *Tree Physiol.* 2008; 28:1049–1058
4. Portnern HO, Roberts DC, Tignor M, Poloczanska ES, Mintenbeck K, Alegria A et al, editors. IPCC, 2022: Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press. Cambridge University Press, Cambridge, UK and New York, NY, USA. 2022;3056 pp. doi:10.1017/9781009325844.
5. Chikkodi SB, Ravikumar RL. Influence of pollen selection for *Alternaria helianthi* resistance on the progeny performance against leaf blight in sunflower (*Helianthus annuus* L.). *Sex Plant Reprod.* 2000;12: 222–226.
6. Frova C, Portaluppi P, Villa M, Sari-Gorla M. Sporophytic and gametophytic components of thermotolerance affected by pollen selection. *J. Hered.* 1995; 86: 50–54
7. Stephen R, Naik JB, Haridas. 2004. Invitro Pollen selection for water stress tolerance in coconut. *J. Plantation crops.* 2004; 32(suppl):45-49
8. Evaluation of Selectively Fertilized coconut hybrids (*Cocos nucifera*) for water use efficiency through stable isotope discrimination. M.Sc. (Ag) thesis, Kerala Agricultural University, Thrissur. 2015; 132
9. Package of Practices Recommendations: Crops (16 th Ed.). Kerala Agricultural University, Thrissur. 2024: 96
10. Nithinkumar K, Saikia N, Gairola A, Kumar V, Twinkle J, Kumari M, et al., Relative Water Content: Concept and Practical Implications. *Agriculture and Food:E Newsletter* . 2024;6(2): 280-284.www.agrifoodmagazine.co.in. ISSN:2581-8317
11. Saelim S, Zwiazek JJ. Preservation of thermal stability of cell membranes and gas exchange in high temperature-acclimated *Xylia xylocarpa* seedlings, *J. Plant Physiol.*2000; 156:380–385
12. Mohan MM, Subramanian LN, Ibrahim S. Chlorophyll Stability Index (CSI): Its impact on salt tolerance in rice. *IRRN.* 2000; 25:38-39.
13. Shepherd T, Griffiths DW. The effects of stress on plant cuticular waxes. *New Phytol.* 2006; 171: 469–499
14. Senthilkumar M, Amaresan N, Sankaranarayanan A. (2021). Estimation of Malondialdehyde (MDA) by Thiobarbituric Acid (TBA) Assay. In: *Plant-Microbe Interactions. Springer Protocols Handbooks.* Humana, New York NY. 2021. https://doi.org/10.1007/978-1-0716-1080-0_25
15. Pandey V, Awasthi M, Singh S, Tiwari S, Dwivedi U A. Comprehensive Review on Function and Application of Plant Peroxidases. *Biochem. Anal. Biochem.* 2017. <https://doi.org/10.4172/2161-1009.1000308>
16. Del Río LA, Corpas FJ, López-Huertas E, Palma JM. Plant superoxide dismutases: Function under abiotic stress conditions. In: Gupta D, Palma., Corpas F, editors. *Antioxidants and Antioxidant Enzymes in Higher Plants.* Springer; Cham, Switzerland: 2018; 1–26.
17. Sadasivam S, Manickam A. *Biochemical Methods.* New Age International (P) Ltd, New Delhi.2015

18. Gopinath P, Parsad R, Joseph B, Adarsh VS. grapesAgri1: Collection of Shiny Apps for Data Analysis in Agriculture. Journal of Open-Source Software. 2021; <https://doi.org/6.10.21105/joss.03437>
19. Lokesha AN, Shivashankara K, Hunashikatti L, Sadashiva AT, Shankar A. Response of Contrasting Tomato Genotypes under High Temperature Stress. Mysore J. Agric. Sci. 2019;53 (2): 9-14 (2019)
20. Afna AP, Stephen, Manohar V, Vanaja T, Manju RV, Vishnu R (2023). Physiological Assessment of Temperature Stress Tolerance in Selectively Fertilized Coconut Hybrids. Int. J. Environ. Clim Chang. 2023;13.:391-398.
21. Singh P, Prasad S, Verma A, Lal B, Singh R, Singh S, et al . Screening for Heat Tolerant Traits in Wheat (*Triticum aestivum* L.) Genotypes by Physio-biochemical Markers. Int. J. Curr. micro. Appl. sci. 2020; 9:2335-2343.
22. Mohammed S, Huggins TD, Beecher F, Chick C, Sengodan P, Mondal S, et al. The Role of Leaf Epicuticular Wax in the Adaptation of Wheat (*Triticum aestivum* L.) to High Temperatures and Moisture Deficit Conditions. Crop Sci.2018; 58: 679-689. <https://doi.org/10.2135/cropsci2017.07.0454>
23. Li L, Du Y, He C, Dietrich C, Li J, ma X, et al. A novel maize gene, glossy6 involved in epicuticular wax deposition and drought tolerance. 2018; 10.1101/378687.
24. Zafar SA, Hameed A, Khan AS, Ashraf M. Heat shock induced morpho-physiological response in indica rice (*Oryza sativa* L.) at early seedling stage. Pak. J. Bot. 2017; 49:453–463.
25. Liu X, Xiao Y, Zi J, Yan J, Li C, Du C, et al. (2023). Differential effects of low and high temperature stress on pollen germination and tube length of mango (*Mangifera indica* L.) genotypes. Sci. Rep. 2023; 13(1):1-14. <https://doi.org/10.1038/s41598-023-27917-5>
26. Gulen H, Eris. Effect of heat stress on peroxidase activity and total protein content in strawberry plants. Plant Sci. 2004; 166:739-744. 10.1016/j.plantsci.2003.11.014.
27. Jespersen D. Heat shock induced stress tolerance in plants: Physiological, biochemical, and molecular mechanisms of acquired tolerance. In: Hossain MA, Liu F, David J, Burritt D, editors. Priming-Mediated Stress and Cross-Stress Tolerance in Crop Plants. 1st ed. Academic Press Elsevier Inc.; London, UK: 2020;161–174.
28. Karkute SG, Ansari WA, Singh AK, Singh PM, Rai N, Bahadur A, Singh J. Characterization of high-temperature stress-tolerant tomato (*Solanum lycopersicum* L.) genotypes by biochemical analysis and expression profiling of heat-responsive genes. 3 Biotech. 2021;11(2):45. doi: 10.1007/s13205-020-02587-6. Epub 2021 Jan 11. PMID: 33489667; PMCID: PMC7801581.

ABBREVIATIONS

N.F= Normally Fertilized
 S.F=Selectively Fertilized
 HT=High Temperature
 AT=Ambient Temperature
 RWC= Relative Water Content
 CSI=Chlorophyll Stability Index
 ECW= Epicuticular Wax
 SOD=Superoxide dismutase
 ROS= Reactive Oxygen Species
 MDA= Malondialdehyde
 IPCC=Intergovernmental Panel on Climate Change

Term: Definition for the term