

Understanding Future Healthcare Providers' Knowledge, Attitudes, and Practices on Supplement Use at a Private Medical College in Lahore

Abstract

Background: The use of dietary supplements without healthcare provider recommendations is widespread, although many supplements should ideally be used under professional supervision.

Objective: This study aimed to evaluate the knowledge, attitude, and perception of future healthcare providers regarding the use of supplements.

Methods: A validated questionnaire was administered to future healthcare providers from various faculties, including medical, dentistry, and pharmacy. The questionnaire was divided into segments: demographics, attitude, knowledge, and perception regarding the use of supplements.

Results: The study revealed that female students exhibited a more positive attitude towards the use of supplements, with a mean score of 44.43 out of 50 (SD = 2.11), compared to males who scored 39.32 (SD = 4.65). Final-year students had a better attitude compared to pre-final-year students ($P = 0.001$, $\eta^2 = 0.213$). In terms of knowledge, females scored higher, averaging 8.9 out of 10. Pharmacy students demonstrated better knowledge compared to students from other faculties ($P = 0.35$, $\eta^2 = 0.14$). Perception results indicated that females had a more positive perception than males, and non-hostellers had a better perception compared to hostellers ($P = 0.008$, $\eta^2 = 0.18$). Overall, females, pharmacy students, and final-year students showed better knowledge, attitude, and practices regarding supplement use.

Conclusion: The study highlights the importance of targeted educational interventions to improve the knowledge, attitude, and perception of future healthcare providers regarding the appropriate use of supplements, with particular attention to gender and faculty differences.

Keywords: Malnutrition, COVID-19 pandemic, vitamin

1. Introduction

Nutrition is a fundamental requirement for a healthy body (1). Malnutrition and poor diet are primary risk factors for diseases and disabilities (2). A healthy diet provides essential nutrients for the proper functioning, growth, and development of living cells (3). Calcium and vitamin D are crucial for bone development (4). Additionally, vitamin D plays a vital role in the prevention and management of atherosclerosis, diabetes mellitus, metabolic syndrome, cardiac remodeling, and cardiovascular diseases (5). People often use supplements to meet the major nutritional requirements for maintaining a healthy body (6). In broiler chickens, *Spirulina platensis* is used as a nutritional supplement to improve the quality of meat, serving as a valuable source of human protein (7).

Supplements play a crucial role in boosting immunity, including essential vitamins (fat-soluble vitamins A and D, and water-soluble vitamins B12, B6, and folates) and minerals like zinc, selenium, iron, and copper (2,3). The usage rate of dietary supplements among gym members is 43.8% (4). During the COVID-19 pandemic, the prevalence of vitamin C use in Saudi Arabia was 68.4% (3). In Dubai, 87.9% of the general population used vitamins. Among university students in Kazakhstan, supplement use is highest among swimmers, with a rate of 55.0% (4). The use of dietary supplements among pharmacy students in Saudi Arabia is 46.8% (5, 6). In Japan, approximately 63.9% of the population uses nutritional supplements (7).

Dietary supplements are essential for individuals of all ages, tailored to the body's requirements for homeostasis and proper functioning, particularly in children and adolescents (1). Dietary fibers contribute significantly to the maintenance of pregnancy by ensuring optimal levels of iron, calcium, folate, and zinc in plasma (2). For pregnant women who avoid meat and dairy products, dietary supplements are particularly beneficial (3). The demand for nutrients like choline increases during pregnancy, as it is crucial for fetal development and helps prevent premature birth (4). To mitigate oxidative stress during pregnancy, antioxidants such as vitamins A, C, and E, along with selenium and zinc, play a vital role (5).

Vitamin D is a vital supplement, with cholecalciferol being the most effective form, while calcidiol and calcitriol are less suitable for use (1). In the management of osteoarthritis, hyaluronic acid (HA), chondroitin sulfate (CS), and glucosamine serve as chondroprotective dietary supplements (2). For patients with rheumatoid arthritis, dietary fibers and supplements such as probiotics, vitamin D, and the Mediterranean diet are beneficial, whereas red meat and salts have adverse effects (3). In menopausal women, decreased estrogen levels lead to reduced bone density; vitamin D is used to prevent bone fractures (4). People with cardiovascular problems benefit from supplements as well; coenzyme Q10 and vitamin D help reduce myocardial dysfunction, statin myopathy, and the risk of coronary artery disease (5).

In post-discharge colorectal cancer patients, oral nutritional supplements enhance the effectiveness of chemotherapy and reduce the risk of sarcopenia and muscle loss (1). Nutritional supplements also benefit brain health by improving memory, cognitive performance, and overall brain function (2). Creatine, a neuroprotective supplement, is effective in cases of traumatic brain injury and helps improve focus and sleep during phases of insomnia (3). Among various therapies, dietary supplements are considered one of the safest treatments for obesity and have become particularly appealing to people in the post-COVID era (4).

The objective of the current study is to evaluate the knowledge, attitude, and practice regarding supplements among future healthcare providers. Understanding supplements is crucial for guiding the general population about the basic requirements of nutrition. Nutritional deficiencies can lead to serious health issues, while an overdose of nutrients can also adversely affect health. Healthcare providers play a key role in advising on the maintenance of proper nutrition for a healthy body.

2. Methodology

This research study employed a cross-sectional design to evaluate the knowledge, attitudes, and practices regarding supplement use among students at a private medical college in Lahore. The study focused on final-year and pre-final-year students from the faculties of pharmacy, medicine, and dentistry. The participants were diverse in terms of age, gender, race, and living status. Conducted from August 15, 2022, to December 10, 2022, the study utilized a pre-validated

questionnaire. The survey aimed to include a minimum of 60 students from each faculty, with a total of 286 students participating.

Inclusion Criteria

The study included all pre-final and final-year medical, dental, and pharmacy students who were willing to participate and complete the distributed form.

Exclusion Criteria

1. The exclusion criteria include first-year, second-year, and third-year students of medicine, dental, and pharmacy at the private medical college.
2. Students of physiotherapy, biotechnology, and nutrition sciences at the private medical college are also excluded from the study.

Data Collection Method

The questionnaire was designed to evaluate the knowledge, attitudes, and practices regarding supplement use among students at a private medical college in Lahore. Experts in clinical pharmacy and pharmacy practice verified it. After validation and necessary revisions, the final survey questionnaire comprised 30 questions categorized into knowledge, attitude, and practice sections.

Section A (Attitude-based questions): Utilized a Likert scale (strongly disagree, disagree, neutral, agree, strongly agree).

Section B (Knowledge-based questions): Required selecting the correct answers.

Section C (Practice-based questions): Involved yes or no answers.

The first part of the questionnaire included an informed consent form and demographic details such as gender, age, faculty, year of study, and place of living. The second part contained 10 attitude-based, 10 knowledge-based, and 10 practice-based questions.

Likert scale questions (attitude-based): Scored from 1 to 5, with strongly disagree scoring 1, disagree scoring 2, neutral scoring 3, agree scoring 4, and strongly agree scoring 5.

Knowledge and perception-based questions: Correct answers scored 1 and incorrect answers scored 0.

Practice-based questions: Consisted of yes or no options, with only one correct answer to choose from.

Statistical Analysis

The collected data were entered into SPSS version 21 for analysis. Descriptive statistics were conducted to determine the percentage, frequency, mean, and standard deviation for the quantitative data. The normality of the data was assessed using the Kolmogorov-Smirnov test and by checking the skewness and kurtosis values. Upon confirming the normality of the data, parametric tests, such as the one-way ANOVA, were performed. A p-value of less than 0.05 was considered statistically significant.

3. Results:

The study comprised 286 participants who completed the questionnaire. The majority of respondents were female (59%), with the remaining 41% being male. Among the participants, 119 were from the Faculty of Medicine, 120 from the Faculty of Pharmacy, and 47 from the

Faculty of Dentistry. A total of 208 students were in their pre-final year, while 78 students were in their final year. The largest proportion of participants (78%) were in the age group of 21–25 years. Additionally, 76% of the 286 students were non-hostel residents.

Table 1 indicates that faculty, age, and place of residence do not significantly influence students' attitudes toward supplement use ($p > 0.05$), however females are having more positive attitude as compared with males. Table 2 summarizes students' practices regarding supplement use, revealing a significant difference among faculties with a p-value of 0.035 with effect size 0.014. According to Cohen's classification, the η^2 value of 0.014 indicates a weak positive association. In contrast, year of study, age, and place of residence showed no significant differences. Table 3 demonstrates a statistically significant difference in knowledge about supplements among two genders ($p = 0.008$, $\eta^2 = 0.0024$), year of education ($p = 0.008$) with an η^2 value of 0.047, suggesting a small positive association. For the remaining variables, the p-values are considerably higher than 0.05, indicating no statistically significant differences in knowledge about supplements with respect to gender, faculty, year of study, and place of residence.

Table 1: Attitude of students on supplement use(Total Maximum score was 50 and minimum was 10 for 10 questions).

Outcome Variable	Mean (SD)	95% Confidence Interval (C.I)		t- statistic (df)	p- value	Effect size (η^2)
		Lower bounds	Upper bounds			
Gender						
Male	39.87±4.85	38.56	41.63	7.762	0.003	0.034
Female	44.43±2.11	42.67	45.86	(1, 283)		
Faculty						
Medicine	41.39±3.67	39.78	41.98	18.67	0.261	-
Pharmacy	45.54±5.79	41.34	43.82	(1, 283)		
Dentistry	41.61±2.87	39.76	41.32			
Age						
18-20	40.61±4.61	38.86	41.78	32.48	0.357	-
21-25	41.69±4.51	40.97	42.16	(1, 283)		
>25	43.38±3.96	39.76	46.18			
Year of study						
Pre-final Year	40.82±4.96	40.54	41.90	8.679	<0.001	0.213
Final Year	44.92±3.34	41.68	45.57	(1, 283)		
Place of living						
Hostellar	42.31±4.61	41.56	42.41	7.561	0.431	-
Non-hostellar	42.41±5.52	41.54	42.32	(1, 283)		

Female	8.98±1.91	6.89	9.06	(1,283)		
Faculty						
Medicine	6.36±2.61	4.41	7.04	6.723	0.035	0.014
Pharmacy	8.36±1.94	5.98	9.04	(2,283)		
Dentistry	4.61±3.75	4.82	5.92			
Age						
18-20	5.92±1.73	5.74	6.09	5.382	0.506	-
21-25	5.56±1.88	5.15	5.98	(2,283)		
>25	6.68±2.88	6.11	7.32			
Year of study						
Pre-final Year	5.98±2.61	4.89	5.89	0.567	0.009	-
Final Year	8.13±1.91	7.36	9.13	(1,283)		
Place of living						
Hostellar	5.52±2.82	4.98	6.01	7.495	0.189	-
Non-hostellar	5.95±2.90	5.05	6.22	(1,283)		

Table 2 Practice of Students on supplement use(Total Maximum score was 10 and minimum was 0 for 10 questions).

Outcome Variable	Mean (SD)	95% Confidence Interval (C.I)		<i>t- statistic (df)</i>	<i>p- value</i>	<i>Effect size (η²)</i>
		Lower bounds	Upper bounds			
Gender						
Male	17.23±2.54	16.45	17.76	0.431 (1,283)	0.008	0.124
Female	19.12±2.67	18.79	19.84			
Faculty						
Medicine	18.45±2.47	17.97	18.92	13.556 (2,283)	0.653	-
Pharmacy	19.09±2.18	18.32	19.11			
Dentistry	17.29±3.62	17.85	19.93			
Age						
18-20	17.41±3.89	17.04	18.08	43.831 (2,283)	0.231	-
21-25	19.20±2.13	18.41	19.48			
>25	18.00±1.87	17.65	18.02			
Year of study						
Pre-final Year	16.28±2.97	15.70	16.76	6.782 (1,283)	0.003	0.161
Final Year	19.65±1.56	18.26	19.85			
Place of living						

Hostellar	15.69±2.74	14.69	15.98	8.397	0.008	0.198
Non-hostellar	18.51±2.82	18.17	19.83	(1,283)		

Table 3:
Knowledge of
students on
supplement

use (Total Maximum score was 20 and minimum was 0 for 20 questions).

UNDER PEER REVIEW

4. Discussion:

The present study represents a novel evaluation of the use of supplements among medical students at a university in Pakistan. Previous research in Pakistan has not assessed medical students' knowledge, attitudes, and practices regarding the self-use of supplementary medicines despite their future roles as healthcare providers. To evaluate the attitudes of these medical students, a total of 10 questions were administered using a five-point Likert scale.

The findings revealed that female students exhibited a more positive attitude compared to their male counterparts, with a significant P value of 0.007. However, the effect size was found to be 0.024, indicating a weak positive association between gender and student attitudes. These results align with several previous studies that have reported more positive attitudes among females than males(5). This trend may be attributed to females' greater health consciousness and concern for health-related issues, or it could be due to a generally more positive disposition among females(6).

The findings of the current study indicate that final-year students across all healthcare provider faculties have a more positive attitude towards the use of supplementary medicines compared to pre-final-year students. This is supported by a P value of 0.001 and an effect size of 0.213, suggesting a moderate positive association between the year of education and attitudes towards supplementary medicine use. A similar study conducted in Malaysia by Iqbal et al. reported that final-year students exhibited a more favorable attitude towards various health-related questions compared to pre-final-year students(7). The probable explanation for this trend could be the increased professionalism of final-year students regarding the use of supplementary medicines, as they are on the verge of graduating and becoming healthcare providers(8). This impending transition likely contributes to the more positive attitude observed among final-year students.

None of the other variables showed a significant relationship with the use of supplementary medicine. The study found no significant difference in attitudes among students from the faculties of medicine, pharmacy, and dentistry. However, pharmacy students exhibited a more positive attitude compared to students from the other two faculties. Similarly, when comparing age groups, students older than 25 years had a more positive attitude than those aged 18 to 25 years. These findings are supported by a study conducted in Malaysia, which also reported that pharmacy students had a more positive attitude compared to students from

other faculties. A probable reason for this could be the higher number of pharmacy students involved in research activities compared to students from the other faculties(9).

Another aspect of the current study was to evaluate the practices of future healthcare providers regarding the use of supplementary medicines. The findings indicate that only gender and faculty variables have a positive relationship with the practice of using supplementary medicines, while age, year of study, and place of living do not show any direct relation. The data revealed that female students had better practice scores compared to male students. Specifically, female students scored an average of 8.98 with a standard deviation of 1.91, whereas male students scored an average of 5.87 with a standard deviation of 1.83. The P value was 0.003, and the effect size (η^2) was 0.054, indicating a weak positive association between gender and the practice of using supplementary medicines among future healthcare providers. The probable reason for this difference could be that female students tend to have better practices regarding the use of supplementary medicines compared to males(10). These findings are supported by a study conducted by Iqbal et al., which also indicated that females had better practice scores compared to males, possibly due to similar reasons(11).

On the other hand, students from the pharmacy faculty demonstrated better practices regarding the use of supplementary medicines compared to those from the medical and dentistry faculties. Additionally, dentistry students exhibited the lowest practice scores in response to questions about the use of supplementary medications. A probable reason for this could be the higher use of supplements by pharmacy students and their tendency to recommend them. In contrast, dentistry students are less likely to consume and refer supplementary medicines to their patients. A statistically significant association was observed, with a P value of 0.035 and an effect size (η^2) of 0.014, indicating a weak positive association between faculty affiliation and the practice of supplementary medicine use. These findings are supported by previous studies conducted in different countries, which also showed that pharmacy students have better practices compared to students from other faculties(8,12). The probable reason for this could be the more frequent practical training and emphasis on the use of supplements within the pharmacy curriculum(13).

In a recent study assessing the knowledge of future healthcare providers, a total of 20 questions were used for evaluation. The findings revealed that females demonstrated better knowledge than males. The mean score for females was 19.12 with a standard deviation of 2.67, whereas the mean score for males was 17.23 with a standard deviation of 2.54. A

statistically significant association was observed between gender and knowledge, with a 95% confidence interval and an effect size of 0.124, indicating a positive weak association according to Cohen's classification of effect size(14). These results align with a study conducted in Malaysia by Iqbal et al., which also found that females had better knowledge compared to males(15). A plausible explanation for this difference is that females have a better understanding of their own health conditions compared to males. This justification has been supported and elaborated by various previous research studies(16,17).

The knowledge levels of final-year students were found to be significantly higher compared to pre-final-year students, with a p-value of 0.003 and an effect size of 0.161. This indicates that final-year students possess more extensive knowledge compared to pre-final-year students. These findings are consistent with the literature, which similarly reports that graduating students exhibit more advanced knowledge than all other students (18).

5. Conclusion:

This study provides a comprehensive evaluation of the attitudes, practices, and knowledge regarding supplement use among medical, pharmacy, and dentistry students at a university in Pakistan. Female students exhibited significantly more positive attitudes towards supplement use compared to males ($p = 0.007$, $\eta^2 = 0.024$). Final-year students had a more favorable attitude than pre-final-year students ($p < 0.001$, $\eta^2 = 0.213$). There were no significant differences in attitudes based on faculty, age, or place of residence. Gender and faculty affiliation significantly influenced practices regarding supplement use. Females demonstrated better practices compared to males ($p = 0.003$, $\eta^2 = 0.054$). Pharmacy students exhibited better practices than those from the medicine and dentistry faculties ($p = 0.035$, $\eta^2 = 0.014$). Final-year students possessed significantly higher knowledge than pre-final-year students ($p = 0.003$, $\eta^2 = 0.161$). Female students consistently showed more positive attitudes, better practices, and higher knowledge levels regarding supplement use compared to their male counterparts. This suggests a greater health consciousness and understanding of health-related issues among female students. Final-year students demonstrated more positive attitudes and higher knowledge levels, likely due to their advanced education and proximity to entering professional healthcare roles. This underscores the importance of education and professional training in shaping attitudes and knowledge. Pharmacy students had better practices related to supplement use, possibly due to the curriculum and practical training emphasizing supplement use more than other faculties.

Ethical Approval and Consent

The Lahore Medical and Dental College Human Ethics Committee approved the study protocol, including demographic authorization, on July 15, 2022. The Faculty Research Committee also reviewed and granted ethical clearance for the study. All respondents provided informed consent, and their information was kept confidential for research purposes.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of manuscripts.

REFERENCES

1. Lin Y. Effects of post-discharge oral nutrition supplement on nutritional status and body composition in malnourished patients with gastrointestinal cancer and the clinical implications. :1–19.
2. Whalley D, Globe G, Crawford R, Doward L, Tafesse E, Brazier J, et al. Is the EQ-5D fit for purpose in asthma? Acceptability and content validity from the patient perspective. *Health Qual Life Outcomes*. 2018 Aug 3;16(1).
3. Forbes SC, Cordingley DM, Cornish SM, Gualano B, Roschel H, Ostojic SM, et al. *Health*. 2022;1–16.
4. Bonetti G, Herbst KL, Donato K, Dhuli K, Kiani AK, Aquilanti B, et al. Dietary supplements for obesity. 2022;63:160–8.
5. Alowais MA, Selim MAE-H. Knowledge, attitude, and practices regarding dietary supplements in Saudi Arabia. *J Fam Med Prim Care [Internet]*. 2019 [cited 2024 Jul 12];8(2):365. Available from: /pmc/articles/PMC6436290/
6. Pop LM, Iorga M, Şipoş LR, Iurcov R. Gender Differences in Healthy Lifestyle, Body Consciousness, and the Use of Social Networks among Medical Students. *Med* 2021, Vol 57, Page 648 [Internet]. 2021 Jun 24 [cited 2024 Jul 12];57(7):648. Available from: <https://www.mdpi.com/1648-9144/57/7/648/htm>
7. Iqbal M, Rathi R, Prajapati S, Omar K, Bahari M, Rajan S, et al. Knowledge, attitude, and practice of

- oral hygiene among students of a private university. *J Pharm Bioallied Sci* [Internet]. 2020 [cited 2020 Nov 9];0(0):0. Available from: <http://www.jpbonline.org/preprintarticle.asp?id=298082;type=0>
8. Upadhyay D, Manirajan Y, Iqbal M, Paliwal N, Pandey S. A survey on medical, dental, and pharmacy students' knowledge, attitude, and practice about hepatitis B infection in a private medical University of Malaysia. *J Res Pharm Pract* [Internet]. 2020 [cited 2020 Nov 12];9(3):128. Available from: <https://www.jrpp.net/article.asp?issn=2319-9644;year=2020;volume=9;issue=3;spage=128;epage=134;aualast=Upadhyay>
 9. Zahid Iqbal M, Khan S-U-D, Shahid M, Iqbal M. Oral healthcare attitude among students of a medical university Medical Science. *Med Sci* [Internet]. 2020 [cited 2020 Nov 9];24(106):3891–9. Available from: http://discoveryjournals.org/medicalsecience/current_issue/v24/n106/A7.htm
 10. Budu HI, Abalo EM, Bam V, Budu FA, Peparah P. A survey of the genesis of stress and its effect on the academic performance of midwifery students in a college in Ghana. *Midwifery*. 2019 Jun 1;73:69–77.
 11. Iqbal MS, Iqbal MZ, Ahmed NJ. Evaluation of Community Pharmacists Practices towards Dengue Control and Management. *J Young Pharm* [Internet]. 2020 Jun 9 [cited 2020 Sep 19];12(1):90–3. Available from: <http://asiapharmaceutics.info/index.php/ajp/article/view/3620>
 12. Woelfel JA, Patel RA, Lee H, Chu S, Ranson C, Patel B, et al. An overview and study of beneficiaries' knowledge, attitudes, and perceptions of the Medicare Part D benefit. *Consult Pharm*. 2015 Feb 1;30(2):101–11.
 13. Iqbal MZ, Alqahtani SS, Mubarak N, Shahid S, Mohammed R, Mustafa A, et al. The influence of pharmacist-led collaborative care on clinical outcomes in type 2 diabetes mellitus: a multicenter randomized control trial. *Front Public Heal*. 2024 Feb 27;12:1323102.
 14. Sullivan GM, Feinn R. Using Effect Size—or Why the P Value Is Not Enough. *J Grad Med Educ* [Internet]. 2012 Sep 1 [cited 2024 Jul 13];4(3):279–82. Available from: <https://dx.doi.org/10.4300/JGME-D-12-00156.1>
 15. Iqbal M, Iqbal M, Deneshwary B, Rahul R, Muhammad Tahir M, Yeoh C, et al. Knowledge and Perception of Medicine, Dentistry and Pharmacy Students of Malaysian University Regarding Ebola Virus Disease (EVD). *Value Heal* [Internet]. 2016 Nov 1 [cited 2020 Aug 30];19(7):A913. Available from: <http://www.valueinhealthjournal.com/article/S1098301516306982/fulltext>
 16. Iqbal MZ, Ali AN, Bahari MB, Iqbal MS. Determinants Affecting Knowledge of Healthcare Students towards Mental Health Challenges. *Int J Pharm Investig*. 2020 Oct 10;10(3):413–6.
 17. Goldberg N, Rodriguez-Prado Y, Tillery R, Chua C. Sudden infant death syndrome: A review. *Pediatr Ann*. 2018 Mar 1;47(3):e118–23.
 18. Hisam A, Rana MN, Mahmood-Ur-Rahman. Knowledge and attitude regarding Ebola virus disease among medical students of Rawalpindi: A preventable threat not yet confronted. *Pakistan J Med Sci* [Internet]. 2016 Jul 1 [cited 2020 Aug 30];32(4):1015. Available from:

UNDER PEER REVIEW