

Original Research Article

Effect of vine management on yield, yield attributing and quality parameters of pumpkin (*Cucurbita moschata*)

ABSTRACT

The present study was carried out at the Instructional cum Research Farm, Department of Horticulture, Biswanath College of Agriculture, Assam Agricultural University, Biswanath Chariali with an aim to study the effect of vine management on yield, yield attributing factors and quality parameters (which included nutritional factors such as vitamin A, vitamin C) of pumpkin. The treatments were: T₁ (Trimming of growing tip of the primary vine at 8th node stage), T₂ (Trimming of growing tip of the primary vine at 10th node stage), T₃ (Trimming of growing tip of the primary vine at 12th node stage), T₄ (Trimming of growing tip of the secondary vine at 6th node stage), T₅ (Trimming of growing tip of the secondary vine at 8th node stage), T₆ (Removal of all tertiary vines), T₇ (Retention of two tertiary vines) and T₈ (control without pruning). T₃ further recorded the minimum days to appearance of first male (49.17 days), female (58.15 days) flowers, fruit set to harvest duration (51.43 days) and crop duration (110.55 days) while T₈ recorded the maximum days. Significant variation was observed in the yield parameters where T₃ recorded the maximum number of fruits (6.27), fruit weight (2.47 kg), fruit yield per plant (15.47 kg) and yield per hectare (27.88 t/ha). Fruit characters such as fruit length, fruit diameter and number of seeds revealed no significant differences among the different treatments but with respect to quality parameters pruning revealed significant differences in terms of vitamin A and ascorbic acid content. The experiment thus concluded that the treatment T₃ (Trimming of growing tip of the primary vine at 12th node stage) produced the highest yield with higher B: C ratio of 2.52 and it can be recommended to the farmers of Assam in order to maximize the production and productivity of pumpkin.

Keywords: trimming, nodes, fruit set, vitamin A, ascorbic acid, benefit-cost ratio

1. INTRODUCTION

The three species of the diverse genus *Cucurbita* L. (pumpkin, squash, gourd) with the highest agro-economic significance are *C. maxima* Duchesne, *C. moschata* Duchesne, and *C. pepo* L. The three are commonly grown for food, feed, and adornment. They are renowned for their excellent fruit polymorphism in terms of size, shape, and colour. Currently, the average annual production in Europe and the rest of the world is 3.38 and 23.70 million tonnes, respectively; this represents a growth of roughly 66 and 100% when compared to the final decade of the previous century. For the same time period globally, the higher total production is mostly due to an increase in harvested area (78%), followed by an increase in yield (13%). In contrast, the

increased yield (37%) and expanded crop area (24%) are to blame for the rise in European pumpkin production. Both the seeds and the flesh of the pumpkin fruit are excellent sources of nutrients; the seeds are a particularly good source of lipids, proteins, and minerals, and the flesh is rich in carotenoids, carbohydrates, flavonoids, minerals, amino acids, and phenolic compounds. The flesh has very low calorie content (about 20 kcal/100 g).

A key factor affecting the success of Cucurbitaceous crops is the quantity of vines per plant [1]. On the other hand, fruit size, mass, and yield are all strongly influenced by the number of fruits per vine [2]. In light of this, it is crucial to give these cucurbit management factors proper considerations. Cucurbits can be controlled for the quantity of vines per plant and fruits per vine using a variety of techniques, including fruit and vine trimming [3]. Pruning is a specialized horticulture technique that involves removing specific plant sections to encourage flowering and subsequent fruiting. As a result, the output increases while improving consumer-required characteristics such as fruit size and attractiveness [4]. According to [5], pruning's primary goal is to encourage a balance between vegetative growth and fruit load. Cucumber and butternut have both been shown to produce more when pruned [6,7]. In order to synchronize the harvesting period and produce homogeneous fruits, pruning is a frequent practice in the watermelon industry [4]. Pruning has an effect on a plant's ability to bear or produce fruit, which affects how well the plant functions. The plant's capacity to produce fruits is established and enhanced. By driving or directing the sap flow towards the section of the plant that bears fruit, pruning causes the plant or vine to forcefully produce fruits of greater quality. The pruning technique and its uses in pumpkin are quite uncommon since farmers have little understanding and inadequate information.

2. MATERIALS AND METHODS

The study was carried out at the Instructional Cum Research Farm, Department of Horticulture, Biswanath College of Agriculture, Assam Agricultural University, Biswanath Chariali (26.7° N latitude, 90.5° E longitude, and at 105 m above MSL) from October 2021 to April 2022.

Eight treatments, each with three replications, were used in the experiment. These treatments included T1 (trimming the growing tip of the primary vine at the eighth node stage), T2 (trimming the growing tip of the primary vine at the tenth node stage), T3, (trimming the growing tip of the primary vine at the twelfth node stage), T4, (trimming the growing tip of the secondary vine at the sixth node stage), T5 (Trimming of growing tip of the secondary vine at 8th node stage), T6 (Removal of all tertiary vines), T7 (Retention of two tertiary vines) and T8 (control without pruning) by utilizing a pumpkin of the same kind. Using secateurs, pruning was carried out when the plants reached their pruning stage in accordance with the various treatments. To prevent damaging the node, the pruning was done above the node. Standard agricultural techniques were followed to assure a robust crop stand, beginning with the full preparation of the experimental plot through thorough ploughing, followed by harrowing and leveling. The entire plot was then divided into 24 plots, each with 8 plots, in 3 replications. Each plot/bed was constructed with a 9 m x 4.5 m dimensions in mind. Then 30 cm³-sized trenches were dug and filled with a mixture of topsoil and cow manure. In the trenches, seeds were planted with a 3m x 1.5m spacing. Each pit was initially seeded with two to three seeds, and then thinning was done in order to keep the healthiest plant in each pit.

Yield attributing factors such as day to appearance of first male and female flowers, duration from fruit set to harvest, crop duration, number of fruits per plant and fruit weight, fruit yield

UNDER PEER REVIEW

per plant, fruit yield were studied. Also quality factors such as vitamin A, ascorbic acid content were examined to study the effect of different pruning treatments.

Vitamin A was estimated by using the formula

$$\beta\text{carotene}(\mu\text{g}/100\text{g}) = \frac{0.013.9 \times 10 \times 100}{\text{Wt of the sample } 560 \times 1000}$$

$$\text{Vitamin A} = \frac{\text{Betacarotene}(\mu\text{g})}{0.6 \times 100\text{g}}$$

For estimation of Ascorbic acid content

$$\text{Ascorbic acid}(\text{mg}/100\text{g}) = \frac{\text{Titre value} \times \text{Dye factor} \times \text{Volume made up}}{\text{Weight of the sample taken for estimation} \times \text{Aliquot of sample taken for estimation}} \times 100$$

$$*\text{Dye factor} = 0.5 / \text{Titre value}$$

Analysis of variance was used to compare data from laboratory tests and observations collected during field experiments. The approach outlined by [8] was used to calculate the corresponding "F" values in order to establish the significance or non-significance of the variation attributable to treatments. By computing critical difference (CD at 5%) estimations, the importance of the difference between the mean values of the treatment's parameters was examined.

3. RESULTS AND DISCUSSION

Effect of vine management on yield attributing factors

Days for appearance of first male flower (Table 1a)

The initial male flower appearance showed a notable variance with various pruning techniques. The treatment T8 had the longest time to produce the first male bloom, measuring 57.47 days, followed by T7, taking 54.42 days. However, under T3, the minimum number of days required for the first male flower to appear was noted to be 49.17 days.

The shortest duration for the emergence of the initial male (49.17 days) and female (58.15 days) flowers was noted for treatment T3, which involved trimming the primary vine's developing tip at the 12th node stage. This is in line with a bell pepper experiment conducted by [9], in which it was observed that the plants with pruning began to flower three days earlier than the plants without pruning. Comparing bottle gourd plants with and without pinching at the third node stage [10] showed similar results.

Days for appearance of first female flower (Table 1a)

Out of the eight distinct pruning treatments, treatment T3 took the fewest days (58.15 days) to produce female flowers, followed by treatment T4 (59.35 days). Treatment T8 took the longest (69.16) days to produce the first female flowers. The reduction in apical development that encouraged more lateral branches may have contributed to the early appearance of female flowers in the treatment T3. This is consistent with findings from [11] study on ridge gourd, which showed that early development of female flowers was observed on plants pruned to six principal vines. When plants were clipped on the main stem [12] observed that the cucumbers produced the most blooms and the bud break occurred earlier. Moreover, pepper plants showed a decrease in the length of time it took for female flowers to develop after pruning [13].

Durationfromfruitsettoharvest(Table1a)

UNDER PEER REVIEW

Significant variations in fruit set to harvest duration were found through pruning. The treatment T8 required the greatest time (58.95 days) from fruit set to harvest, followed by T1 (56.46 days) and T5 (55.43 days), while T3 (51.43 days) took the least time. When plants were clipped [14] found that the maximum number of female flowers and the minimum number of male flowers in long melons were recorded. According to [15], unpruned plants had the longest fruit set to harvest time. This confirms the current investigation's findings, which showed that treatment T3 (trimming the primary vine's growing tip at the 12th node stage) took the shortest amount of time (51.43 days) to harvest fruit, and treatment T8 (control without pruning) took the longest (58.95) days.

Crop duration (Table 1a)

The eight different pruning treatments demonstrated significant variations in the length of the crop. T3 had the shortest crop duration, measuring 110.55 days, followed by T4 (113.38 days) and T2 (116.59 days). Treatment T8 had the highest crop duration, measuring 127.96 days. Significant differences between the various pruning procedures were seen in the crop duration from the current study. T3 which involved trimming the major vine's growth tip at the 12-node stage recorded the shortest crop duration (110.55 days), while T8 which involved no pruning at all recorded the longest (127.96) days. One of the reasons for short crop duration of T3 might be because of flowering of both male and female flowers early. Pruning has been shown to be able to regulate fruiting and flowering, according to study published by [16]. The crop duration data in this study is consistent with tomato results that were republished, where plants that were cut to a single stem showed the bare minimum number of days needed for the first harvest as compared to the plants without being pruned [17,18].

Table 1a. Effect of vine management on yield attributing factors

Treatment	Days to first appearance of male flower	Days to first appearance female flower	Duration from fruit set to harvest	Total crop duration
T ₁	50.91	61.07	56.46	119.39
T ₂	52.44	62.24	53.55	116.59
T ₃	49.17	58.15	51.43	110.55
T ₄	50.98	59.35	52.42	113.38
T ₅	52.05	61.50	55.43	117.42
T ₆	53.03	64.53	52.56	120.47
T ₇	54.42	66.29	54.29	122.33
T ₈	57.47	69.16	58.95	127.96
SEd±	0.49	0.33	0.46	0.39
C.D(P=0.05)	1.06	0.71	0.98	0.85

T1: Trimming of growing tip of the primary vine at 8th node stage, T2: trimming of growing tip of the primary vine at 10th node stage, T3: trimming of growing tip of the primary vine at 12th node stage, T4: trimming of growing tip of the secondary vine at 6th node stage, T5: trimming of growing tip of the secondary vine at 8th node stage, T6: removal of

all tertiary vines, T7: retention of two tertiary vines and T8: control without pruning

UNDER PEER REVIEW

Effect of vine management on Quality Parameters

Vitamin A content (Table 2a)

The study examined the effects of various pruning procedures on the concentration of vitamin A. T1 had the highest recorded vitamin A content at 2216.49 IU, followed by T3 at 2150.85 IU and T7 at 2144.07 IU as shown in Table 2a. T8 had the lowest recorded vitamin A content at 1921.58 IU. Significant variations in vitamin A content were found in relation to trimming techniques. T1 (trimming the primary vine's developing tip at the eighth node stage) had the highest vitamin A content (2216.49 IU), whereas T8 (control without pruning) had the lowest (1921.58 IU). In a comparable manner [19] found that sweet potatoes with up to five secondary vines pruned had higher β -carotene content. Additionally, [20] found considerable variations in the vitamin A concentration of pointed gourds grown using various pruning techniques.

Ascorbic acid content (Table 2a)

Significant differences were observed in the ascorbic acid content between the various pruning treatments. T2 had the highest ascorbic acid content (7.42 mg/100g), followed by T6 (6.55 mg/100g). T8 had the lowest ascorbic acid content (5.40 mg/100g), statistically comparable to T5 (5.51 mg/100g) and T1 (5.71 mg/100g).

Several researchers have observed that pruning treatments positively boosted the level of ascorbic acid (vitamin C). When comparing trimmed plants to unpruned plants [21] found that the pruned plants had the highest levels of vitamin C. Comparably, sweet pepper plants that were trimmed to a single branch had higher levels of ascorbic acid [22]. The aforementioned results pertain to the current study, wherein the pruning treatments had an impact on the ascorbic acid concentration. The greater morpho-physiological characteristics brought about by pruning may be the cause of the higher ascorbic acid concentration observed under the pruning therapy. Additionally [23] found that when stem clipping was done, cherry tomatoes had a greater ascorbic content.

Effect of vine management on Fruity yield per plant and per hectare

Effect of vine management on number and weight of fruits (Fig. 1)

Regarding the quantity of fruits produced per plant, it was revealed that there were significant variations among the treatments. Following T1 (6.12) and T4 (5.41), treatment T3 (6.27) produced the highest amount of fruits per plant, while treatment T8 (4.18) produced the lowest fruits per plant. Between the various pruning methods, there were notable variations in the weight of each individual fruit. The lowest fruit weight (2.05 kg) was recorded by T8, which was comparable to T6 (2.16 kg), while the largest fruit weight (2.47 kg) was recorded by T3, which was comparable to T7 (2.37 kg) and T4 (2.33 kg).

With the most secondary vines produced, treatment T3 (trimming of the growing tip of the primary vine at 12th node stage) may have produced more female flowers on the secondary vines, which in turn may have produced more fruits. Various factors, such as fruit size and quantity, affect a larger yield [14]. This aligns with the findings obtained from the current study. While T8 (control without pruning) recorded the lowest fruit weight (2.05 kg), which proves that there is some effect of pruning on fruit weight. Fruit size and quantity are correlated with morpho-physiological factors. Pruning increased the number of branches, leaves, and LAI, which may have increased photosynthetic activity and the accumulation of assimilates, which in turn increased the number and weight of fruits [24].




Fig.1. Effect of vine management on number and weight of fruits

Effect of vine management on Fruit yield per plant and per hectare (Table 2a) The fruit production per plant showed significant differences across the pruning treatments; T3 had the highest output (15.47 kg), followed by T1 (13.40 kg) and T4 (12.61 kg), while T8 had the lowest yield (8.57 kg).

Significant variations were found between the levels of pruning treatments, as shown by the results in Table 2a. T3 recorded the highest yield (27.88 t/ha), followed by T1 (24.19 t/ha) and T4 (22.78 t/ha), while T8 recorded the lowest yield (15.48 t/ha). Additionally, a yield increase of 80.10% was noted from T3 over T8.

The current study showed that, in comparison to the control, trimming helped to increase the overall yield percentage. Table 2a makes it evident that the treatment T3 (trimming of the primary vine at 12 node stage) produced a yield increase of 80.10% in comparison to treatment T8 (control without pruning). It is possible that the pruned plants produced the maximum yield because they produced more and larger fruits. This is consistent with [24] findings with cucumber. By allowing proper light exposure, pruning raised the rate of photosynthesis and, in turn, the source to sink ratio, which raised the marketable output. Research showed that plants with pruning experienced greater fruit yields than those with no pruning [25] for bitter melon, and [26] for tomatoes. According to reports by [27], pruning increased the amount of eggplant that could be sold by reducing the number of waste fruits. When plants were trimmed to four stems in greenhouse-cultivated sweet pepper, the fruit yield was highest compared to unpruned plants [28]. Similar results were found by [29] in chilli and [30] in capsicum.

Table 2a. Effect of vine management on quality parameters, fruit yield per plant (kg) and fruit yield per hectare (t/ha)

UNDER PEER REVIEW

Treatments	Quality parameters		Fruityield (kg/plant)	Fruityield (t/ha)
	Vitamin A content(IU)	Ascorbic acid content (mg/100g)		
T ₁	2216.49	5.71	13.40	24.19
T ₂	1984.59	7.43	11.85	21.36
T ₃	2150.85	5.77	15.47	27.88
T ₄	1965.77	6.42	12.61	22.78
T ₅	1987.35	5.51	11.16	20.18
T ₆	2067.96	6.55	11.10	20.05
T ₇	2144.07	5.75	10.28	18.55
T ₈	1921.58	5.40	8.57	15.48
SEd±	0.81	0.23	0.03	0.02
C.D(P=0.05)	1.70	0.50	0.06	0.05

T1: Trimming of growing tip of the primary vine at 8th node stage, T2: trimming of growing tip of the primary vine at 10th node stage, T3: trimming of growing tip of the primary vine at 12th node stage, T4: trimming of growing tip of the secondary vine at 6th node stage, T5: trimming of growing tip of the secondary vine at 8th node stage, T6: removal of all tertiary vines, T7: retention of two tertiary vines and T8: control without pruning.

4. CONCLUSION

From the study conducted on different pruning treatments it was revealed that it had a significant effect in terms of yield attributing, quality and yield parameters of pumpkin. The treatment T3 was revealed to be best among the different pruning treatments.

REFERENCES:

1. Gomes RF, Santos LDS, Braz LT, Andrade FLDN, Monteiro SMF. Number of stems and plant density in mini watermelon grown in a protected environment. - Pesq. Agropec. Trop. 2019; 49: e54196, pp. 1-8.
2. Lins HA, Queiroga RCF, Pereira ADM, Silva GD, Albuquerque JRT. Growth, yield and quality of fruits of watermelon in function changes in relation sink-source. - Revista Verde de Agroecologia e Desenvolvimento Sustentável. 2013; 8(3): 143-149.
3. Campos AMD, Luz JMQ, Santana DG, Marquez GR. Influences of plant density and fruit thinning on watermelon hybrid production cultivated in different seasons. - Hort. Bras. 2019; 37(4): 409-414.
4. Oga IO, Umekwe PN. Effects of pruning and plant spacing on the growth and yield of watermelon (*Citrullus lanatus* L.) in Unwana-Afikpo. - Int. J. Sci. Res. 2016; 5(4):

110-115.

UNDER PEER REVIEW

5. Anwar NA, Gad AA, Bardisi A, Zyada HG. Effect of plant spacing and apical shoot pinching on growth and productivity of watermelon plants under sandy soil conditions. *Zagazi J. Agric. Res.* 2019; 46(2): 357-365.
6. Nayak SR, Parmar VK, Patel AN, Suchismita J, Lathiya JB, Tandel YN. Efficacy of pinching and plant growth regulators in enhancing yield characters of cucumber (*Cucumis sativus* L.). - *Int. J. Chem. Stud.* 2018; 6 (1): 1804-1807.
7. Eve B, Tuarira M, Moses M. The influence of pinching on the growth, flowering pattern and yield of butternuts (*Cucurbita moschata*). - *Int. J. Hort. Orn. Pl.* 2016; 2(1): 19-25.
8. Panse VG, Sukhatme PV. *Statistical Method for Agricultural Workers*. Indian Council of Agricultural Research, New Delhi; 1985.
9. Awal S, Shahjahan M, Roy AC, Akter A, Kabir MH. Response of bell pepper (*Capsicum annuum*) to foliar feeding with micronutrients and shoot pruning. *J. of Agri. and Eco. Res. Intern.* 2017; 11(3): 1-8.
10. Naafe M, Nabi G, Irshad M, Khan MN, Ali S, Hayat R. Influence of pinching on growth and yield of bottle gourd (*Lagenaria siceraria*). *Pure and App. Bio.* 2022; 11(4): 891-901.
11. Arora SK, Malik IJ. Effect of pruning and spacing levels on growth, flowering, earliness and fruit yield in Ridge gourd. *Haryana J. Hort. Sci.* 1989; 18(1-2): 99-105.
12. Ekwu LG, Nwokwu GN, Utobo EB. Effect of mulching materials and pruning on growth and yield of cucumber (*Cucumis sativus* L.). *Nig. Agri. J.* 2017; 48(2): 51-59.
13. Dongre SM, Mahorkar VK, Joshi PS, Deo DO. Effect of micronutrients spray on yield and quality of chilli (*Capsicum annuum* L.) var. Jayanti. *Agri. Sci. Digest.* 2000; 20(2): 106-107.
14. Singh AK, Sabir N, Jat GS, Singh J, Singh V, Singh A, Kumar A, Kumar J. Effect of spacing and pruning on growth, yield and economics of long melon (*Cucumis melo* var. *utilissimus*) under naturally ventilated polyhouse. *The Ind. J. of Agri. Sci.* 2021; 91: 885-89.
15. Kumar S, Singh BD. Pointed Gourd: Botany and Horticulture. *Horticultural Reviews*, 2012; 39: 206-226.
16. Mnzava NA. Influence of cluster and spacing on set, growth, yield and quality of early fruit of the West Virginia 63 tomato (*Lycopersicon esculentum* Mill.). Dissertation for Award of M.Sc. Degree at Sokoine University of Agriculture, Morogoro Tanzania. 1984; pp. 54.
17. Alam MS, Islam N, Ahmad S, Hussen MI, Islam MR. Effect of different staking methods and stem pruning on yield and quality of summer tomato. *Bangla. J. of Agric. Res.*, 2016; 41(3): 419-432.
18. Yadav S, Ameta K.D, Sharma SK, Dubey RB, Rathore RS, Kumar, Kapuriya VK. Effect of spacing and training on vegetative growth characteristics and yield of tomato (*Solanum lycopersicon* L.) Grown in polyhouse. *Int. J. Curr. Microbio. App. Sci.*, 2017; 6(5): 1969-1976.
19. Rahman MS, Hossain MM, Mian MAK, Hossain MF. Effect of vine pruning on yield and quality of sweet gourd (*Cucurbita moschata*), 2013.
20. Gupta KK, Buragohain N, Gautam BP, Langthasa S, Goswami RK, Kalita MK. Impact of sprout management on growth, quality and yield of pointed gourd (*Trichosanthes dioica* Roxb.). *Int. J. Curr. Microbiol. Appl. Sci.* 2020; 9: 1030-1037.

21. PatilSD,ShindeVN,MaherMB,DahatondeKA.EffectofPruningandMicronutrients on Growth, Quality and Chlorophyll Content in Sweet Pepper. Intern. J. of Current Microbio. and App. Sci., 2020; 11: 825-829.
22. CebulaS,KaliszaA,KunickiE.Canopyformationofsweetpepperplantsprunedtoone main shoot in greenhouse production. Folia Horticulturae. 1998; 10(2): 35-44.
23. Razzak HA, Ibrahim A, Wahb-Allah M, Alsadon A. Response of cherry tomato (*Solanum lycopersicum* var. *cerasiforme*) to pruning system and irrigation rates under green house condition. Asian J. Crop Sci. 2013; 64: 321-332.
24. Shivaraj D, Lakshminarayana D, Prasanth P, Ramesh T, Studies on the Effect of Pruning on Cucumber cv. Malini Grown Under Protected Conditions. Int. J. Curr. Microbio. App. Sci. 2018; 7: 2019-2023.
25. Palada MC, Chang LC. Suggested cultural practices for bitter gourd. AVRDC. No. 3; 2003: pp 547.
26. Hesami A, Khorami SS, Hosseini SS. Effect of shoot pruning and flower thinning on quality and quantity of semi-determinate tomato (*Lycopersicon esculentum* Mill.). Notulae Scientia Biologicae. 2012; 4: 108-111.
27. PaksoyM,AkellaM.Theeffectsofdifferentpruningsontheyieldandqualityof eggplant cultivars grown in the greenhouse conditions 2nd Symp. of Prot. Cultiv. of Solanacea in Mild Winter Climates, Turkey;1993.
28. JovicichE,CantliffeDJ,HochmuthGJ.PlantDensityandShootPruningManagement on Yield of a Summer Greenhouse Sweet Pepper Crop. Hort. Sci. 1999;34:532E-532.
29. Laxman S, Mukherjee S.Effect of foliar application of urea and NAA on yield and yield attributes of chilli (*Capsicum annuum* var: longum). Agri. Sci Digest. 2000; 20: 116-117.
30. Shetty GR, Manohar RK. Influence of pruning and growth regulators on flowering, fruit set and yield of coloured capsicum (*Capsicum annuum* L.) cv. Orabelle under naturally ventilated greenhouse. Asian J. of Hort. 2008; 3(2): 213-216.

UNDER PEER REVIEW