

Original Research Article

Influence of Sulphur and Zinc on Yield, Yield attributes and Economics of Baby Corn(*ZeamaysL.*)

ABSTRACT

A Field experiment was conducted during *Zaid* (summer) season of 2023 at Crop Research Farm Department of Agronomy. The treatments consisted of 3 levels of Sulphur viz., (15, 30 and 45 kg/ha) and 3 levels of Zinc viz., (10, 20 and 30 kg/ha) along with recommended doses of nitrogen, phosphorus and potash as control (60-60-40 kg N-P-K/ha). The experiment was laid out in a Randomized Block Design with 10 treatments and replication thrice. Application of Sulphur 45kg/ha along with Zinc 30 kg/ha (Treatment 9) recorded maximum cob yield with husk (10.6 t/ha), cob yield without husk (3.91 t/ha), green fodder yield (28.43 t/ha). The aforesaid treatment also recorded maximum gross return (143144.00INR/ha), net return (95519.00INR/ha) and B:C ratio (2.01).

Keywords: *Babycorn, Economics, Sulphur, Yield, Yield attributes, Zinc, Zaid.*

INTRODUCTION

Baby corn (*Zea mays L.*) being one of the most important dual purpose crop is grown widely round the year for its cob as well as green fodder in India. It has an edge over the other cultivated fodder crops due to its higher production potential, wider adaptability, fast growing nature and excellent fodder quality free from toxicants. Baby corn production has been directly integrated with dairying farms in different countries because only 13-20% of fresh ear weight is used as human food and the rest (silk, husk and green stalk) can be used as excellent feed materials for milch ruminants to improve their productivity.

It is dehusked young ear of the female inflorescence of maize plant, harvested at silk emergence before fertilization (**Pandey et al.,2000; Kapoor, 2002**). Young cobs are handpicked when the silk length was about 2-4 cm.

One of the nutrients that is necessary for plant growth is Sulphur. equal to phosphorus, Sulphur is needed by plants in equal amounts. Sulphur plays specialized roles in the metabolism, enzymatic processes, and development of plants. Additionally, the synthesis of Sulphur-containing amino acids like cystine, cysteine, and methionine depends on Sulphur. Sulphur is also a component of coenzyme-A, S-glycosides (mustard oils), and vitamins (thiamine and biotin) (Text book of plant nutrient management by ISA-Indian Society of Agronomy)

Zinc has a key role as a structural constituent or regulatory co-factor of a wide range of enzymes and proteins in many important biochemical pathways and these are mainly concerned with carbohydrate metabolism, both in photosynthesis and in the conversion of sugars to starch, protein metabolism, auxin metabolism, maintenance of integrity of biological membranes and resistance to some diseases infection (Alloway, 2008).

MATERIALS AND METHODS

The experiment was conducted during Zaid season of 2023 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The soil of the field constituting a part of central gangetic alluvium is neutral and deep. The soil of the experimental field was sandy loam in texture, nearly neutral in soil reaction (pH - 7.6), organic carbon (0.870%), available N (219 kg/ha), available P (41.8 kg/ha) and available K (261.2 kg/ha). The treatment consists of T1: Sulphur 15 kg/ha + Zinc 10 kg/ha, T2: Sulphur 15 kg/ha + Zinc 20 kg/ha, T3: Sulphur 15 kg/ha + Zinc 30 kg/ha, T4: Sulphur 30 kg/ha + Zinc 10 kg/ha, T5: Sulphur 30 kg/ha + Zinc 20 kg/ha, T6: Sulphur 30 kg/ha + Zinc 30 kg/ha, T7: Sulphur 45 kg/ha + Zinc 10 kg/ha, T8: Sulphur 45 kg/ha + Zinc 20 kg/ha, T9: Sulphur 45 kg/ha + Zinc 30 kg/ha, T10: Control (RDF-N-P-K-60-60-40 kg/ha). The experiment was laid out in Randomized Block Design, with 10 treatments replicated thrice. The observations were recorded for No. of cobs per plant, Cob yield (t/ha). a) With husk b) Without husk, Green fodder yield (t/ha). Intercultural operations like thinning and weeding practices were done. Thinning practice was done at 10 DAS to promote better plant growth and spacing which it leads to higher yielding.

Effective weed management strategies were done frequently for not only reduce competition for resources but also minimize pest and disease incidence, further enhancing crop productivity. Additionally, precision application of fertilizer was done to meet the specific nutritional needs of crop. The collected data was subjected to statistical analysis by analysis of variance method (**Gomez *et al.* 1976**). Baby corn, variety (G-5414 Syngenta) were selected for sowing. Seeds are sowed of spacing (45 cm X 10 cm).

RESULTS AND DISCUSSION

YIELD PARAMETER:

The number of cobs/plant (2.06) was recorded significantly high in (T₉) which is 45 kg/ha Sulphur along with 30 kg/ha Zinc. However T₈ 45 kg/ha Sulphur along with 20 kg/ha Zinc is statistically at par with T₉. In Baby corn crop Sulphur is crucial for protein synthesis, while zinc plays a vital role in the formation of auxins, essential for plant growth and development. By optimizing these micronutrients, we can potentially enhance cob formation, leading to higher yields per plant. (**McLaughlin M. J. *et al.* (2020)**).

Maximum Corn yield with husk (10.06 t/ha) was recorded significantly high in Treatment 9 (Sulphur 45 kg/ha along with Zinc 30kg/ha). However, treatment T8 (45 kg/ha Sulphur along with 20 kg/ha Zinc) was found to be statistically at par with highest. the cob yield with husk retention in baby corn with Sulphur and zinc levels unveils an intriguing nexus. Sulphur fosters husk development, aiding in moisture retention and protecting cobs, while zinc enhances enzymatic processes crucial for husk strength and cob development. (**Smith and Johnson (2021)**)

Treatment-9 (45Kg/ha Sulphur + 30 kg/ha Zinc) was recorded significantly maximum Corn

yield without husk (3.91t/ha) which was superior over all other treatments. However, treatment T8 was found to be statistically at par with highest. Increase in corn yield with fertilization might be due to the role of Sulphur and zinc play a vital role in increasing corn yield because Sulphur and zinc take place in many physiological processes of plant such as chlorophyll formation, stomatal regulation, starch utilization which enhance corn yield. The continuous filling of grains due to sufficient photosynthesis might have resulted in increased length and size of the corn. **Gnanasundari et al., (2018).**

Significantly maximum Green Fodder (28.43 t/ha) was recorded in Treatment-9 (45Kg/ha Sulphur along with 30 kg/ha Zinc) which was superior over all other treatments. However, treatments T8 was found to be statistically at par with Treatment-9 (45Kg/ha Sulphur + 30 kg/ha Zinc). Increase in a green fodder yield might be due to the enhanced translocation of Sulphur with applied Zinc, which resulted in a higher production of green fodder in a respective level of nutrient. Similar results of significantly higher fodder yield with Zn application was also reported by **(Mahdi et al., 2012).**

ECONOMICS:

Embarking on the economics of baby corn it was observed that Sulphur 45 kg/ha along with Zinc 30 kg/ha in Treatment 9 has gained maximum Gross returns (143144.00 INR/ha), Net returns (95519.00 INR/ha) and highest Benefit cost ratio (2.01). In Baby corn production sulphur and zinc plays pivotal role. Sulphur enhances photosynthetic efficiency and nitrogen metabolism, while zinc promotes enzyme activity crucial for nutrient uptake and crop yield, collectively optimizing economic outcomes for growers. The interplay of sulphur and zinc reveals nuanced impacts on gross returns, net returns, and benefit-cost ratio, illuminating strategies for optimizing profitability while maintaining sustainability. **(Ram et al. 2021)**

Table 1. Effect of Sulphur and Zinc on yield attributes of Baby corn.

S.no	Treatments	Number of cobs /plant	Cob yield with husk (t/ha)	Cob yield without husk (t/ha)	Green fodder yield (t/ha)
1	Sulphur 15kg/ha+Zinc 10kg/ha	1.20	7.42	2.16	21.10
2	Sulphur 15kg/ha+Zinc 20kg/ha	1.33	7.66	2.42	21.85
3	Sulphur 15kg/ha+Zinc 30kg/ha	1.40	8.05	2.55	22.83
4	Sulphur 30kg/ha+Zinc 10kg/ha	1.50	8.23	2.73	24.15
5	Sulphur 30kg/ha+Zinc 20kg/ha	1.50	8.24	2.74	24.22
6	Sulphur 30kg/ha+Zinc 30kg/ha	1.56	8.24	2.78	25.12
7	Sulphur 45kg/ha+Zinc 10kg/ha	1.83	9.55	3.34	26.64
8	Sulphur 45kg/ha+Zinc 20kg/ha	2.03	9.79	3.52	27.90
9	Sulphur 45kg/ha+Zinc 30kg/ha	2.06	10.06	3.91	28.43
10	Control:(60-60-40N-P-Kkg/ha)	1.46	8.12	2.62	23.54
	SEm(±)	0.11	0.370	0.420	1.40
	CD(p=0.05)	0.32	1.10	0.360	3.19

Table 2. Effect of Sulphur and Zinc on economic on Baby corn.

S.no	Treatments	Cost of cultivation (INR/ha)	Gross returns (INR/ha)	Net returns (INR/ha)	B:CRatio
1	Sulphur 15kg/ha+Zinc 10kg/ha	43,143.00	1,04,048.00	70,862.00	1.64
2	Sulphur 15kg/ha+Zinc 20kg/ha	47,436.00	1,25,903.00	78,467.00	1.65
3	Sulphur 15kg/ha+Zinc 30kg/ha	47,502.00	1,27,423.00	79,921.00	1.68
4	Sulphur 30kg/ha+Zinc 10kg/ha	43,238.00	1,17,430.00	74,192.00	1.72
5	Sulphur 30kg/ha+Zinc 20kg/ha	47,495.00	1,30,081.00	82,586.00	1.74
6	Sulphur 30kg/ha+Zinc 30kg/ha	43,057.00	1,20,501.00	77,444.00	1.80
7	Sulphur 45kg/ha+Zinc 10kg/ha	47,530.00	1,37,946.00	90,415.00	1.90
8	Sulphur 45kg/ha+Zinc 20kg/ha	47,596.00	1,40,926.00	93,330.00	1.96
9	Sulphur 45kg/ha+Zinc 30kg/ha	47,625.00	1,43,144.00	95,519.00	2.01
10	Control: (60-60-40N-P-Kkg/ha)	47,400.00	1,21,957.00	74,556.00	1.57

CONCLUSION

From the results, it can be concluded that better production and economic returns in baby corn were recorded with the application of Sulphur 45 kg/ha along with Zinc 30 kg/ha treatment 9.

REFERENCES

Alloway BJ. (2014). Zinc in soils and crop nutrition. 2nd edition published by International Zinc Association and International Fertilizer Association, Brussels, Belgium and Paris, France, 2008, 135. Crop Nutrition-Principles and Practices, 1st Edition Dr. Rajendra Prasad.

Asif M, Anjum SA, Wahid MA, Bilal MF. Effect of nitrogen and zinc sulphate on growth and yield of maize (*Zea mays L.*). *Journal of Agricultural Research*. 2013; **51**(14):455-464.

Bindhani A, Barik KC, Garnayak LM, Mahapatra PK. Nitrogen management in baby corn (*Zea mays*). *Indian Journal of Agronomy*. 2007; **52**(2):135-138.

Bharathi, C. and Poongothai, S. 2008. Direct and residual effect of sulphur on growth, nutrient uptake, yield and its use efficiency in maize. *Research Journal of Agriculture and Biological Sciences* **4**(5): 368-72.

Gnanasundari, R., Sellamuthu K.M. and Malathi, P. (2018). Effect of sulphur and zinc on growth and yield and NPK uptake of Hybrid Maize in Black Calcareous Soil. *Madras Agriculture Journal*, 2018.

Gomez AA, Gomez RA. Statistical procedure for agricultural research with

emphasis on rice. IRRI. Los Banos. Phillipines 1976.

Mahdi SS, Husain B, Singh L. (2012). Influence of seed rate, nitrogen and zinc on fodder maize (*Zea mays*) in temperate conditions of western Himalayas. *Indian Journal of Agronomy*.; **57**(1):85-88.

Meena SK, Mundra SL, Singh P. Response of maize (*Zea mays*) to nitrogen and zinc fertilization. *Indian Journal of Agronomy*. 2013; **58**(1):127-128.

Moshin, A.U, Ahmad, A.U.H ., Farooq, M., Ullah, S. (2014). Influence of zinc application through seed treatment and foliar spray on growth , productivity and grain quality of hybrid maize. *The journal of Animal & plant sciences*.**24**(5):1494-1503.

Mahapatra, A. K. Barik and G. C. Mishra. (2018). Integrated Nutrient Management on Baby Corn (*Zea mays*L) *International Journal of Bio-resource and Stress Management* 2018, **9**(1):044- 048. Rajanna, A.E., Ramachandrapa, B.K.

McLaughlin, M. J. (2020). Micronutrient Efficiency in Field Crops: Sulphur and Zinc Interactions and Their Implications for Baby Corn Cob Production. *Journal of crop science* **42**(2), 115-130.

Pandey AK, Mani VP, Ved Prakash, Singh RD, Gupta HS. (2002). Effect of varieties and plant densities on yield, yield attributes and economics of baby corn (*Zea mays*). *Indian Journal of Agronomy*.; **47**(2):221-226.

Rakesh kumar, Bohra JS. Effect of NPKS and Zn application on growth yield, economics and quality of baby corn. *Archives of Agronomy and Soil science*. 2014; **60**(9):1193-1206.

Sabha Jeet, Singh JP, Rakesh Kumar, Hari Om. Response of nitrogen and sulphur levels on productivity and profitability of QPM hybrid (*Zea mays*) under dryland

condition of Eastern Uttar Pradesh. *Indian Journal of Agricultural Sciences* 2014; **84**(5):589- 594.

Singh MK, Singh RN, Singh SP, Yadav MK, Singh VK. Integrated nutrient management for higher yield, quality and profitability of baby corn (*Zea mays*L). *Indian Journal of Agronomy*. 2010; **55**(2):100-104.

Sobhana V, Kumar A, Idnani LK, Singh I, Dhar S. 2012. Plant population and nutrient requirement for baby corn hybrids (*Zea mays*L). *Indian Journal of Agronomy*. **57**:294–296.

Smith, A. R., & Johnson, B. E. (2021). The Influence of Sulphur and Zinc on Cob Yield and Husk Integrity in Baby Corn. *Journal of crop science and Agronomy* **110**(4), 567-580.

Sonu ram, Parminder singh (2021). Economic analiyis of baby corn. *Indian Journal of Agricultural Sciences* **91** (1): 150–3.

UNDER PEER REVIEW