

# Standardising The Different Levels of NPK on Yield and Yield Attributes of Potato (*Solanum tuberosum* L.) for North Zone of Bihar

---

## Abstract

This field research was carried out on a potato crop of the cultivar *Kufri Lalit* in the Rabi season of 2021–2022 at research farm of TCA, Dholi (Muzaffarpur), Bihar. 13 treatments and 3 replications were used in this experiment's randomised block design. The experimental location had a sandy loam texture, a reaction pH of 8.47 that is alkaline, a low organic carbon content of 0.43%, and accessible N ( $223 \text{ kg ha}^{-1}$ ). However, P ( $16.95 \text{ kg ha}^{-1}$ ) and K ( $131.46 \text{ kg ha}^{-1}$ ) availability is moderate. The varying levels of key nutrients had a substantial impact on all growth and yield parameters, including percent emergence, plant height, shoot and leaf number  $\text{plant}^{-1}$ , dry matter accumulation, tuber bulking rate and yield. Among all treatments, treatment T<sub>10</sub> ( $240\text{N}$ ,  $120\text{P}_2\text{O}_5$  and  $150\text{K}_2\text{O} \text{ kg ha}^{-1}$ ) recorded highest per cent emergence (94.00 %), plant's height (44.27), shoot's number  $\text{plant}^{-1}$  (5.60), leaf's number  $\text{plant}^{-1}$  (56.30), tuber yield ( $26.53 \text{ t ha}^{-1}$ ) and treatment T<sub>10</sub> also recorded statistically at par with treatment T<sub>6</sub>.

(Key words: Potato, treatments, varying levels and alkaline)

## Introduction

Potato (*Solanum tuberosum* L.) is classified under family Solanaceae and in terms of human consumption world's III<sup>rd</sup> most significant food crop after wheat and rice. Potato appears to have developed via isolation in both geography and ecology. In general, potatoes needed an acidic soil with a pH of 5.5 for optimum growth and development. A number of many types of soils Mollisols, Inceptisols, Alfisols, Entisols, and Vertisols, are also used to grow potatoes. In terms of texture, sandy loam soils with a pH 5.50 to 8.0 are said to be the best for growing potatoes. Because of its sensitivity to alkalinity, potato cultivation should be avoided on soils with a pH of more than 8.2. (Pandey, 2007). The most significant dietary component of the potato, which has excellent nutritional value and contains crucial essential dietary elements, is carbohydrates. Proteins, minerals including calcium (Ca), phosphorus (P), and iron (Fe), as well as vitamins B<sub>1</sub>, B<sub>2</sub>, B<sub>6</sub>, and C are other essential components. It is made up of 70-82 percent water, 11-23 percent carbs, 0.8-3% protein, 0.1-% fat, 1.1-% mineral, and 17-29 percent dry matter. Ash content is 0.90 percent (Pandey, 2007).

Nitrogen is mostly contained in plants in the form of the plant cell wall, chlorophyll, protein content, DNA, and RNA. In general, a plant employs nutrients to carry out tissue growth, cell expansion, and the development of organs and systems. Nitrogen makes up between 2 and 7 percent of the Plant. The amount of nitrogen that plants need varies depending on the species. Plants that need a lot of water have large vegetative development, food reserves, or reproductive structures. Nitrate and ammonium in soils are the two main nitrogen sources for plants. (Owen and Jones, 2001). A lack of nutrients can cause stunted growth, tissue death, or yellowing of the leaves because less chlorophyll, a pigment vital to photosynthesis, is produced.

After nitrogen, phosphorus (P) is the mineral nutrient that has the greatest impact on crop output in the potato plant. P is the most significant primary nutrient limiting factor for potato development after nitrogen and potassium, according to Jasim *et al.* (2020). Reduced carbon absorption results from a lack of a deficiency in P that directly impacts photosynthesis by lowering the quantity of inorganic P available in the chloroplast. a deficiency in P in leaf mesophyll cells. According to enhanced photoassimilation to the roots is promoted by phosphorus deficit. At a yield of 29 t ha<sup>-1</sup>,

potatoes lose around 91 K<sub>2</sub>O kg ha<sup>-1</sup>, with potassium enhancing both marketable and total tuber weight by increasing average tuber diameters. (Moinuddin *et al.*, 2004). More so than on yield, potassium fertilisation has a beneficial impact on the quality of the tubers. The most popular techniques for recommending potassium for crops are based on soil testing, and while they occasionally work well for directing fertiliser applications, it is important to establish the crucial level of soil test potassium. The alternative strategy is to base fertiliser recommendations on agronomic effectiveness and yield response. Similar to how nitrogen absorption by potato crops is influenced by meteorological conditions, soil type, fertility, farmed variety, and crop management techniques. According to Bhattarai and Swarnima (2016) Maintaining osmotic potential enhances root permeability and water absorption, regulates ionic equilibrium controls stomata in plants and stimulates enzymatic reactions. Potassium aids in all of these processes.

### **Materials and Methods**

The current research trial investigated in plot number 11 at TCA Farm, Dh0li, Muzaffarpur, Dr. RPCAU, Pusa (Samastipur), Bihar, during the RabiseasOn of 2021-2022.

The experimental site situated at 25°98' North (N) latitude and 85°60' East longitude on the Burhi Gandak river's southern bank, which is 52.2 metres above mean sea level. Experimental soil was calcareous-alluvium and somewhat alkaline in nature having pH 8.47 and soil was moderately fertile-being low organic carbon (0.43%), nitrogen (223 kg ha<sup>-1</sup>), potassium (131.46 kg ha<sup>-1</sup>) and medium phosphorus content (16.95%). The crop was sown in November eighteen at 60x20 row to row and plant to plant spacing. Experiment was laid out in RCBD statistical design with 3 replications involving following treatments T<sub>1</sub>-(Control), T<sub>2</sub> (150N, 90P<sub>2</sub>O<sub>5</sub> and 100K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>3</sub> (0N, 80P<sub>2</sub>O<sub>5</sub> and 150K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>4</sub> (120N, 80P<sub>2</sub>O<sub>5</sub> and 150K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>5</sub> (180N, 80P<sub>2</sub>O<sub>5</sub> and 150K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>6</sub> (240N, 80P<sub>2</sub>O<sub>5</sub> and 150K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>7</sub> (300N, 80P<sub>2</sub>O<sub>5</sub> and 150K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>8</sub> (240N, 0P<sub>2</sub>O<sub>5</sub> and 150K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>9</sub> (240N, 40P<sub>2</sub>O<sub>5</sub>, and 150K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>10</sub> (240N, 120P<sub>2</sub>O<sub>5</sub> and 150K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>11</sub> (240N, 80P<sub>2</sub>O<sub>5</sub> and 0K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>12</sub> (240N, 80P<sub>2</sub>O<sub>5</sub> and 50K<sub>2</sub>O kg ha<sup>-1</sup>), T<sub>13</sub> (240N, 80P<sub>2</sub>O<sub>5</sub> and 100K<sub>2</sub>O kg ha<sup>-1</sup>).

The variety *Kufri Lalit* was developed by clonal selection from the hybrid 85-P-670 x CP-3192 with selection number 2001-P-55 at CPRI, Shimla (HP.) in 2014. It is released for cultivation mainly in North-eastern plains of India.

**% Plant emergence** = total number emerged plants/ total planted tubers x 100

#### **Plant's height**

Randomly chosen 5 plants from every plot were measured in height extending from the soil to the plant neck. The height was averaged using centimeter measurements.

#### **Total shoot's number plant<sup>-1</sup>**

The potato's shoots number plant<sup>-1</sup> counted at an interval of two weeks in each plot from randomly selected plant.

#### **Total leaf's number plant<sup>-1</sup>**

The number of plant's leaves plant<sup>-1</sup> was counted from randomly selected five plant of net plot area of each plot at 45DAP, 60DAP, 75DAP, and at harvest.

#### **Accumulation of dry matter (g plant<sup>-1</sup>)**

Measurements of dry matter accumulated (g plant<sup>-1</sup>) where two plants plucked from every plot of each treatment's gross plot area. The weight of the entire plant (tuber and shoots) was measured and averaged. The plants were then cut into bits, and 100g of the

total 'homogenous samples' were stored in paper bags, dried by oven, and then baked at 70<sup>0</sup>degrees Celsius in hot air oven till a consistent mass was achieved. As a result, the dry weightplant<sup>-1</sup> treatment<sup>-1</sup> ratio was calculated.

#### **Tuber's bulking rate(gday<sup>-1</sup>plant<sup>-1</sup>)**

Tuber's bulking rateobservation was taken at 45DAP, 60DAP, 75DAP and at harvest. On fresh weight basis of tubers, an increase tuber weight at 15-day intervals was calculated splitting the weightof tubers between 15 days intervals in gday<sup>-1</sup>plant<sup>-1</sup>.

#### **Tuber's yield (tha<sup>-1</sup>)**

All the collectedtubers from the net plot area of each where shadow dried tuber weights was calculated by using electronic balance into tha<sup>-1</sup>.

#### **Haulm yield (tha<sup>-1</sup>)**

Fifteen plants selected from each plotof net plot area were removed and properly Dried, and the mass of the vines was documented in kilogramsplot<sup>-1</sup>.

#### **Harvest index (%)**

$$= \frac{\text{Tuber's yield (tha}^{-1}\text{)}}{(\text{Tuber's yield} + \text{Vine's yield (tha}^{-1}\text{)})} \times 100$$

## **Result and Discussion**

### **Growth Parameter**

Data presented in Table No.- 1 revealed thatResponse of different nutrients level onPlant emergence (%) at 30 DAP,Plant height(cm), Number of Shoots plant<sup>-1</sup>and Number leaves plant<sup>-1</sup>reading were significantly increased by the application of different levels of NPK compare with control. Generally, the maximum value of growth parameter was observed with the was noted in treatment T<sub>10</sub>(240N, 120P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>) as compared to other treatments and these are statistically at par with the T<sub>7</sub> (300N, 80P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>) and T<sub>6</sub> (240N, 80P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>). In plant emergence different treatments had a non-significant effect on germination in percent at 30 DAP.Fertilizer levels generally had almost no impact on germination as compared to alternative treatments, NPK kg ha<sup>-1</sup> had uneven germination at 30<sup>th</sup> day after planting.Additionally, it was shown that increasing fertiliser levels caused a delay in the development of tubers (Adhikari 2009). The same

conclusions were reached Marthha *et al.* (2017) and Barman *et al.* (2014). The height of the dominant (44.27 cm) plant was measured in treatment T<sub>10</sub> (240N, 120P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>) at several growth stages. Statistics showed that treatment T<sub>10</sub> was comparable at par with T<sub>7</sub> (300N, 80P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>) but in T<sub>6</sub> (240N, 80P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>) at only 60 days after planting at par with T<sub>10</sub>. This is due to the fact that applying a relatively large amount of nitrogen causes the plant to develop quickly (Adhikari 2009) similar findings was documented by Thirupal *et al.* (2020). The maximum shoot's (5.60) number plant<sup>-1</sup> was recorded with T<sub>10</sub> (240N, 120P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>) it is statistically at par with treatment T<sub>6</sub> (240N, 80P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>) deliberately more than remained treatments. When applying different chemical fertiliser dosages to potatoes, the number of shoots per plant did not alter significantly Marthha *et al.* (2017) These kinds of results as reported Singh and Gupta (2005) and Kumar *et al.* (2008). Number of leaves plant<sup>-1</sup> (56.3) are affected by different level of fertilizers markedly varied between treatments respectively the maximum leaves number plant<sup>-1</sup> was found in T<sub>10</sub>. It was statistically similar with T<sub>6</sub> and T<sub>7</sub>. The response of different potato level to NPK treatment varies significantly, demonstrating a significant interaction between varieties and Application of NPK based on the number of leaves at distinct development stages. The greatest and smallest leaf counts per plant were measured using the application of NPK Thirupal *et al.* (2020) The current findings were similar to research conclusions of Marthha *et al.* (2017).

**Table 1.** Response of different nutrients level on Plant emergence (%) at 30 DAP, Plant height(cm), Number of Shoots plant<sup>-1</sup> and Number leaves

Treatments				Plant emergence (%) at 30 DAP	Plant height(cm)				Number of Shoots plant <sup>-1</sup>				Number leaves plant <sup>-1</sup>			
N (kg ha <sup>-1</sup> )	P <sub>2</sub> O <sub>5</sub> (kg ha <sup>-1</sup> )	K <sub>2</sub> O (kg ha <sup>-1</sup> )			45 DAP	60 DAP	75 DAP	At harvest	45 DAP	60 DAP	75 DAP	At harvest	45 DAP	60 DAP	75 DAP	At harvest
T <sub>1</sub>	Control			87.79	18.9	22.5	24.63	27.96	2.53	2.97	3.27	3.27	27.33	32.33	36.23	21.8
T <sub>2</sub>	150	90	100	92.45	27.4	34	36	38.6	3.65	4.44	4.9	4.9	39.83	45.64	47.56	33.04
T <sub>3</sub>	0	80	150	90.48	23.2	24.04	28.6	33.33	2.6	3.33	3.78	3.78	32.54	35.78	37.8	24.56
T <sub>4</sub>	120	80	150	92.33	26.6	34.57	34.54	37.8	3.53	4.2	4.83	4.83	39.3	43.9	46	32
T <sub>5</sub>	180	80	150	93.2	28.17	35.07	38.9	41	4	4.9	5.04	5.04	40.6	47.6	51.6	34.27
T <sub>6</sub>	240	80	150	93.78	29	35.9	39.63	41.8	4.27	5.33	5.42	5.42	42.7	50.77	54.6	35.33
T <sub>7</sub>	300	80	150	92.01	30.9	37	41.5	43.1	4.07	5.23	5.19	5.19	41.67	49.93	53.7	34.37
T <sub>8</sub>	240	0	150	88.23	24.74	31.46	32.44	36	2.97	3.74	4.17	4.17	36.67	40.8	43.8	29.43
T <sub>9</sub>	240	40	150	92	25.7	33.9	34	37	3.23	3.97	4.78	4.78	38	43.33	45.8	31.8
T <sub>10</sub>	240	120	150	94	32.3	37.8	42	44.27	4.5	5.47	5.6	5.6	43.2	52.43	56.3	37.5
T <sub>11</sub>	240	80	0	89.52	24.6	28.38	31	34.8	2.8	3.6	3.9	3.9	34	37.27	39.9	27.8
T <sub>12</sub>	240	80	50	91.86	25	32.5	33.3	36.4	3.08	3.8	4.47	4.47	37.6	42	45.12	30.8
T <sub>13</sub>	240	80	100	92.9	27.9	34.87	37.6	39	3.8	4.78	5	5	40.36	46.33	49.9	33.9
SEm (±)				3.14	1.11	0.93	0.91	1	0.15	0.18	0.13	0.13	0.92	1.37	1.56	0.88
LSD (p=0.05)				NS	3.23	2.71	2.65	2.92	0.43	0.53	0.39	0.39	2.69	3.99	4.56	2.57

plant<sup>-1</sup>

UNDER PEER REVIEW

## Yield Components

As shown in Table (2), in the most cases treatment T<sub>10</sub> (240N, 120P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>) exhibited high significant increases of yield components included Dry matter accumulation (g plant<sup>-1</sup>), Bulking rate (g day<sup>-1</sup> plant<sup>-1</sup>), Harvest index (%) haulm yield, (t ha<sup>-1</sup>) and Tuber yield (t ha<sup>-1</sup>) tested seasons compared to control treatment. Dry matter accumulation (g plant<sup>-1</sup>) was recorded maximum (85.5) with T<sub>10</sub> and it was statistically similar with treatment T<sub>5</sub> (81.90 g plant<sup>-1</sup>), T<sub>6</sub> (83.95 g plant<sup>-1</sup>) T<sub>7</sub> (83.05 g plant<sup>-1</sup>). The total DMA rose as crop ages increased, while the total DMP was significantly influenced by NPK levels (Banerjee *et al.* 2016). Chemical fertilizer supply plant nutrients easily that can be quite significant in photosynthetic activity of and dry matter accumulation activity of plants. These research's conclusions concur with those of Kushwah *et al.* (2005), Alam *et al.* (2007), Lal and Khurana (2013), and Patel (2013). tuber bulking rate difference that is statistically significant between all of the treatments The bulking rate was registered maximum (7.87) at 60 DAP with treatment T<sub>10</sub> and lowest (0.97) bulking rate at 75DAP was observed with T<sub>1</sub> (control). This was primarily caused by plants satisfying dosages diverting more sugars and photosynthates for crop growth and development, increasing the amount of radiation captured, an increase in the production of photosynthates. this outcome comparable to the end of Meena *et al.* (2016) and Banerjee *et al.* 2016). The observed mean data for harvest index showed that, despite statistical analysis of the no effects of various fertilizers level on various treatments, The partitioning of photosynthates to sink and the greater photosynthetic rate during the tuberization stage may be to blame for this. This result was supported by Nag (2006) and Patel (2013). The quintessential haulms yields were considered and recorded maximum (11.85 t ha<sup>-1</sup>) with T<sub>10</sub> (240N, 120P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>). These are statistically at par with treatments T<sub>6</sub> and T<sub>7</sub>. finding of Shaaban and Kisetu (2014) and Uzatunga *et al.* (2021). the key factor contributing to the rise in the haulms yield seems to be the availability of nitrogen at increasing at a particular level. The tuber yield (tha<sup>-1</sup>) data gathered showed that different level of fertilizes treatments had a significant impact on the tuber yield (tha<sup>-1</sup>). best tuber yield (26.53) with all possible treatments was produced with treatment T<sub>10</sub>. Higher growth and production may be attributable to the crop's improved nutrient availability, which may have raised the plant's photosynthetic capacity and metabolic activity with an up to a particular amount of NPK dose increase. (Adhikari 2009) the current research finding conform to findings of Sharma and Singh (1988) and Rykbost *et al.* (1993).

**Table 2.** Response of different nutrients level on Dry matter accumulation ( $\text{g plant}^{-1}$ ), Bulking rate ( $\text{g day}^{-1} \text{plant}^{-1}$ ), Harvest index (%), haulm yield ( $\text{t ha}^{-1}$ ) and Tuber yield ( $\text{t ha}^{-1}$ )

Treatments				Dry matter accumulation ( $\text{g plant}^{-1}$ )				Bulking rate ( $\text{g day}^{-1} \text{plant}^{-1}$ )			Harvest Index (%)	Haulm yield ( $\text{t ha}^{-1}$ )	Tuber yield ( $\text{t ha}^{-1}$ )
	N ( $\text{kg ha}^{-1}$ )	P <sub>2</sub> O <sub>5</sub> ( $\text{kg ha}^{-1}$ )	K <sub>2</sub> O ( $\text{kg ha}^{-1}$ )	45 DAP	60 DAP	75 DAP	At harvest	45 DAP	60 DAP	75 DAP			
T <sub>1</sub>	Control			19.64	22.64	29	11.28	2.52	3.43	0.97	63.62	6.45	11.28
T <sub>2</sub>	150	90	100	26.23	22.24	45.43	77.19	3.25	68.31	2.94	68.31	10.32	22.24
T <sub>3</sub>	0	80	150	21.33	15.94	33.76	56.91	2.63	68.15	1.42	68.15	7.45	15.94
T <sub>4</sub>	120	80	150	25.46	21.54	43.96	75.05	3.14	68.61	2.61	68.61	9.86	21.54
T <sub>5</sub>	180	80	150	27.77	23.64	49.32	81.9	3.68	69.09	3.37	69.09	10.58	23.64
T <sub>6</sub>	240	80	150	29.25	25.23	52.23	83.95	3.95	68.43	3.64	68.43	11.64	25.23
T <sub>7</sub>	300	80	150	28.57	24.63	51.09	83.05	3.83	68.9	3.56	68.9	11.12	24.63
T <sub>8</sub>	240	0	150	23.65	17.05	36.07	65.05	2.76	63.86	1.86	63.86	9.65	17.05
T <sub>9</sub>	240	40	150	24.79	20.18	39.9	71.1	3.04	66.98	2.39	66.98	9.95	20.18
T <sub>10</sub>	240	120	150	30.83	26.53	55.74	85.5	4.33	69.12	3.92	69.12	11.85	26.53
T <sub>11</sub>	240	80	0	22.25	16.45	35.7	61.43	2.7	63.83	1.71	63.83	9.32	16.45
T <sub>12</sub>	240	80	50	24.11	19.85	37.6	67.06	2.96	66.97	2.05	66.97	9.79	19.85
T <sub>13</sub>	240	80	100	26.86	23.17	47.54	77.65	3.42	68.89	3.1	68.89	10.47	23.17
SEm ( $\pm$ )				0.73	0.8	1.43	0.93	0.11	0.17	0.07	1.95	0.42	0.93
LSD ( $p=0.05$ )				2.13	2.34	4.18	2.69	0.34	0.48	0.2	NS	1.24	2.69

## CONCLUSION

Out of thirteen treatments tested during the research work, treatment T<sub>10</sub> (240N, 80P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O kg ha<sup>-1</sup>) was shown that the higher number of leaves, shoots, dry matter accumulation, plant height, tuber bulking rate and tuber, haulm yield and harvest index. However, variety *Kufri Lalit* responded significantly to the levels of NPK application and resulted better responses and trends were observed with the use of 240N, 80P<sub>2</sub>O<sub>5</sub> 150K<sub>2</sub>O level of fertilizer. The germination percentage at 30 days after planting and harvest index showed no discernible variation between the treatments. At several growth stages, including 45, 60, 75 and at harvest the number of leaves, shoots, plant height dry matter accumulation, tuber bulking rate on a plant was significantly impacted by different treatments.

## References

- Adhikari, R. C. (2009). Effect of NPK on vegetative growth and yield of *Desiree and Kufri Sindhuri* potato. *Nepal Agriculture Research Journal*, **9**, 67-75.
- Alam, M. N., Jahan, M. S., Ali, M. K., Ashraf, M. A., & Islam, M. K. (2007). Effect of vermicompost and chemical fertilizers on growth, yield and yield components of potato in barind soils of Bangladesh. *Journal of Applied Sciences Research*, **3**(12), 1879-1888.
- Banerjee, H., Rana, L., Ray, K., Sarkar, S., Bhattacharyya, K., & Dutta, S. (2016). Differential physiological response in potato (*Solanum tuberosum* L.) upon exposure to nutrient omissions. *Indian Journal of Plant Physiology*, **21**(2), 129-136.
- Barman, K. S., B. Ram and R. B. Verma (2014). Effect of Integrated Nutrient Management on Growth and Tuber Yield of Potato (*Solanum tuberosum*) cv. *Kufri Ashoka*. *Trends in Biosciences*, **7**(9): 815-817.
- Bhattacharai, B., & Swarnima, K. C. (2016). Effect of potassium on quality and yield of potato tubers-A review. *International Journal of Agriculture & Environmental Science*, **3**(6), 7-12.

- Kumar, M., Jadav, M. K., & Trehan, S. P. (2008, December). Contributing of organic sources to potato nutrition at varying nitrogen levels. In *Global potato conference* (Vol. 9, p. 12).
- Kushwah, V.S., Singh, S.P. and Lal, S.S. 2005. Effect of manures and fertilizers on potato (*Solanum tuberosum*) production. *Potato Journal***32(3-4)**: 157-158.
- Lal, M., & Khurana, S. C. (2013). Effect of organic manure, biodynamic compost and biofertilizers on potato. *Potato Journal*, **34(1-2)**.
- Marthha, D., Sahu, G. S., Sahu, P., & Mishra, N. (2017). performance of potato cv. *kufri asoka* as influenced by graded levels of N, P and K. *Plant Archives*, **17(2)**, 1435-1438.
- Meena, B. P., Kumar, A., Dotaniya, M. L., Jat, N. K., & Lal, B. (2016). Effect of organic sources of nutrients on tuber bulking rate, grades and specific gravity of potato tubers. *Proceedings of the national academy of sciences, India section B: biological sciences*, **86(1)**, 47-53.
- Moinuddin, Singh, K., Bansal, S. K., & Pasricha, N. S. (2004). Influence of graded levels of potassium fertilizer on growth, yield, and economic parameters of potato. *Journal of plant nutrition*, **27(2)**, 239-259.
- Nag, G.P., Sarnaik, D.A., Verma, Satish K. and Tamrakar S.K. 2006. Integrated nutrient management in potato for Chhattisgarh plains. *The Orissa Journal of Horticulture*. **36(2)**: 158-161.
- Owen, A. G., & Jones, D. L. (2001). Competition for amino acids between wheat roots and rhizosphere microorganisms and the role of amino acids in plant N acquisition. *Soil biology and biochemistry*, **33(4-5)**, 651-657.
- Pandey, S. K. (2007). Vegetable Science. *Central Potato Research Institute, Shimla, 171001*.
- Patel, B. 2013. Effect of different levels of NPK on growth, development and yield of potato cv. Kufri Ashoka under Chhattisgarh plain condition. M. Sc. (Ag) Thesis IGKV, Raipur. pp. 70-71.
- Rykbost, K. A., Christensen, N. W., & Maxwell, J. (1993). Fertilization of Russet Burbank in short-season environment. *American Potato Journal*, **70(10)**, 699-710.

- Shaaban, H., & Kisetu, E. (2014). Response of Irish potato to NPK fertilizer application and its economic return when grown on an Ultisol of Morogoro, Tanzania. *Journal of Agricultural and Crop Research*, **2(9)**, 188-196.
- Sharma, U. C., & Singh, K. (1988). Response of potato to NP and K in acidic hill soil of Meghalaya. *Journal Indian Potato Association*, **15(1)**, 40-44.
- Singh, S. K. and V. K. Gupta (2005). Influence of farm yard, nitrogen and biofertilizer on growth, tuber yield of potato under rain fed condition in east khasi hill district of Meghalaya. *Agriculture Science Digest*, **25(4)**: 281-283
- Thirupal, D., Ramanandam, G., Jyothi, K. U., Umakrishna, K., Sujatha, R. V., Rao, A. D., & Rao, M. P. (2020). Potato growth as influenced by planting date, spacing and NPK levels under Godavari conditions. *International Journal of Chemical Studies*, **8(4)**, 3638-3643.
- Uzatunga, I. (2021). Effects of fertilizer application on yield and yield related parameters of low yielding potato varieties in Uganda. *African Journal of Agricultural Research*, **17(12)**, 1540-1546.