

**Original Research Article**  
**Innovative seed treatment for *Cassia siamea*  
Lam. germination and carbon stock analysis in  
varied stem sizes for sustainable land  
management and climate change mitigation**

---

**ABSTRACT**

This study investigates seed treatment effects on *Cassia siamea* germination and assesses carbon stock potential across stem size classes. The germination test involved six treatments, including immersion in various water temperatures and chemical solutions. Carbon stock potential across eleven stem size classes of *Cassia siamea* was assessed through air drying, oven drying, and ash weight measurements following standardized methods. Results highlight significant variation in germination rates among treatments, with immersion in concentrated H<sub>2</sub>SO<sub>4</sub> showing the highest efficacy. Carbon stock in wood varies significantly across stem size classes, though no direct relationship with size is observed. Findings suggest non-linear growth patterns in carbon accumulation, influenced by factors beyond stem size. Insights from this research aid in optimizing seedling establishment and maximizing carbon sequestration potential in *Cassia siamea*, supporting climate change mitigation efforts through afforestation initiatives. Moreover, implications for sustainable land management, particularly in contexts like roadside tree planting in India, underscore the practical relevance of these findings in combating rising atmospheric carbon concentrations.

*Keywords: Cassia siamea, seed germination, afforestation, wood carbon stock, climate change mitigation*

**1. INTRODUCTION**

The escalation in carbon dioxide emissions, primarily from burning fossil fuels and deforestation, is a major driver of global warming [1]. This process, exacerbated by human activities, leads to a sustained rise in Earth's average surface temperature, intensifying the dynamics of climate change[2]. It is crucial to prioritize initiatives aimed at bolstering carbon stocks and increasing carbon absorption capacity to effectively mitigate climate change[3].

Trees play a vital role in the global carbon cycle as natural carbon sinks, absorbing atmospheric carbon dioxide through the process of photosynthesis [4]. The planting of trees is widely recognized as a crucial strategy in mitigating climate change by reducing the concentration of greenhouse gases in the atmosphere. Despite this understanding, there remains a notable gap in our understanding of the carbon accumulation potential of exotic tree species, highlighting the need for further research in this area [5]. One such exotic species of interest is *Cassia siamea* Lam., commonly known as Siamese cassia or Siamese senna, native to Southeast Asia. This fast-growing evergreen tree species, belonging to the family Fabaceae (Caesalpinioideae), is valued for its versatility, with applications ranging from timber production to soil improvement [6, 7]. Notably, it exhibits resilience in degraded environments, making it a promising candidate for agroforestry and reforestation initiatives.

The study by Mukhopadhyay and Masto [5] sheds light on the potential of *C. siamea* in carbon sequestration and mine site reclamation. Their findings indicate a significant increase in biomass carbon stock over a 16-year period,

underscoring the species' ecological value. However, accurately quantifying tree carbon stock remains a challenge due to the limitations of direct measurement methods. While direct methods involve cutting down trees and weighing their components, providing precise results, they are often costly, labor-intensive, and ecologically disruptive. Consequently, researchers often turn to indirect methods, such as allometric equations, to estimate biomass without the need for tree destruction [8, 9, 10]. Allometric equations establish empirical relationships between tree biomass and easily measurable structural parameters such as diameter, height, and crown area. However, the choice between using generic or species-specific allometric equations is subject to ongoing debate, with proponents and detractors for each approach [11, 12].

In India, Ragula and Chandra [4] conducted a study on the carbon stock of roadside trees in Bilaspur, emphasizing their role in urban carbon sequestration and climate mitigation efforts. Given India's extensive road network, which is experiencing rapid expansion to accommodate the increasing demands of passenger traffic, the importance of roadside tree plantations cannot be overstated [13]. Trees along roadways serve as crucial carbon sinks, absorbing pollutants such as carbon dioxide emitted by vehicles and mitigating the adverse effects of urbanization on local climate and air quality.

In light of the aforementioned context, understanding the seed germination characteristics and carbon stock potential of *C. siamea* is of paramount importance. Therefore, this study aims to investigate seed germination under various conditions and assess the carbon stock potential of *C. siamea* across various stem size classes. By elucidating these aspects, the research seeks to contribute to the effective management and utilization of *C. siamea* in environmental conservation efforts, agroforestry initiatives, and reforestation programs. Moreover, insights gained from this study can inform policymakers and practitioners on the role of *C. siamea* in enhancing carbon sequestration and climate resilience, particularly in regions experiencing environmental degradation and land-use changes. Ultimately, advancing our knowledge of *C. siamea* towards ecological attributes and carbon sequestration potential can facilitate informed decision-making and promote sustainable land management practices in the face of climate change.

## 2. MATERIAL AND METHODS

We have selected the tree species *Cassia siamea* for the present study. The seeds of *C. siamea*, collected from Bharathiar University campus (located in Coimbatore, India, Fig. 1) were subjected to germination tests under varied treatment conditions. Also, the carbon stock potential of wood samples of *C. siamea* across different stem size classes was assessed using standardized methodologies.

### 2.1 Experiment on seed germination with different treatments

The germination test was done by sowing the seeds in plastic pots (9 cm height and 7 cm diameter). The pots were filled with mixture of topsoil, coarse sand and fine sand. Six seed treatments (T) were set for the experiment, i.e., T1: control (normal water), T2: immersion in cold water (4°C) for 12 h, T3: immersion in hot water (80°C) for 10 min, T4: seeds immersed in hot water (80°C) for 1 min, followed in cold water soaking for 12 h, T5: immersion in concentrated H<sub>2</sub>SO<sub>4</sub> (80%) for 20 min, T6: instead of water KNO<sub>3</sub> was sprayed once in 2 days in the concentration of (0, 3, 6, 9, 12, 15 mg/L). One seed was sown in each pot and pots were kept in shade throughout the experiment. Twenty such pots were done for all the six treatments. The seeds were sown in the depth of 0.5-1.5 cm. *Cassia* species generally require less water, watering can be limited to once in 2 days. Temperature and humidity were recorded in the pot soil every day.

### 2.2 Carbon stock potential across different stem size classes

The wood part of *Cassia siamea* belonging to eleven stem size class (in cm) viz. 5-10, 10-15, 15-20, 20-25, 25-30, 30-35, 35-40, 40-45, 45-50, 50-55, 55-60, were collected and were air dried for two days. Then, the samples were dried in the hot air oven for 24 hours at 105°C to get dried weights. Then, 1 gram of oven dried grind samples leaf and wood were taken separately in pre-weighted crucibles. The crucibles were placed in the furnace at 550±5°C for 2 hours. Then, the crucibles were cooled slowly inside the furnace. After cooling, the crucibles with ash were weighted for calculation of percentage of carbon stock using the following equations [14, 15].

Estimation of carbon stock (C):

$$C (\%) = (100 - \text{ash } \%) \times 0.58$$

Calculation of ash (%) for estimation of carbon stock:

$$\text{Ash } (\%) = (W_3 - W_1) / (W_2 - W_1) \times 100$$

Where W1 is the weight of crucibles

W2 the weight of oven dried grind samples + Crucibles

W3 is the weight of ash + Crucibles

### 2.3 Statistical analysis

One-way analysis of variance (ANOVA) was used to check the variation in seed germination of *C. siamea* among the six treatments, and carbon stock potential of *C. siamea* belonging to different stem size classes. Also, regression analysis was performed to check the relationship between the wood carbon stock and stem size class of *C. siamea*.




Fig. 1. Location map of Bharathiar University campus, Coimbatore, India

### 3. RESULTS AND DISCUSSION

#### 3.1 Seed germination

Among the six seed treatments, the percentage of seed germination was maximum recorded for T5 (95 %) followed by T4 (Table 1). The seed germination results show that the seeds of *C. siamea* treated with immersion in concentrated  $H_2SO_4$  (80%) for 20 min had the maximum (19) seedlings out of the 20 seeds sown (Fig. 2). Statistically, one-way ANOVA showed that seed germination of *C. siamea* varied significantly ( $F_{(5,114)} = 44.345$ ,  $p < 0.001$ ) among the six treatments (Table 1).

Table 1. Percentage of seed germination for *Cassia siamea* among the six treatments

Treatment	Seed germination (at the end of 20th day)	(%)
T1	3	15
T2	3	15
T3	2	10

T4	7	35
T5	19	95
T6	2	10





Fig. 2. Results of seed germination experiment of *Cassia siamea*

### 3.2 Carbon stock potential

Table 2 provides the carbon stock in wood of *C. siamea* categorized by the eleven stem size classes. The carbon stock value was recorded maximum ( $56.43 \pm 0.34\%$ ) at the 55-60cm size class and it was minimum recorded for 40-45cm size class (Table 2). One-way ANOVA showed that the carbon stock potential in wood of *C. siamea* among the eleven stem size classes varied significantly ( $F_{(10,22)} = 7.196$ ,  $p < 0.001$ ). However, regression analysis revealed that there is no relationship ( $R^2 = 0.020$ ) between the wood carbon stock and stem size class of *C. siamea* (Fig. 3).

Table 2. Carbon stock potential of wood of *Cassia siamea* categorized by 11 stem size classes

Stem size class (in cm)	Wood carbon stock (%)	±S.D.
5-10	55.93	0.22
10-15	56.12	0.11
15-20	56.18	0.04
20-25	56.14	0.20
25-30	55.50	0.26
30-35	56.19	0.14
35-40	56.41	0.14
40-45	54.38	0.81
45-50	55.74	0.69
50-55	55.80	0.26
55-60	56.43	0.34



**Fig. 3. Carbon stock potential of wood of *Cassia siamea* categorized by different stem size class**

*Cassia siamea*, primarily displays seed coat-induced dormancy, due to intricate seed coat structures [16]. Findings of the present study underscore the significance of treatment conditions in influencing seed germination. Immersion in concentrated  $H_2SO_4$  emerged as the most effective treatment, yielding a remarkable 95% seedling emergence. Further, one-way ANOVA revealed that seed germination of *C. siamea* varied significantly ( $p < 0.001$ ) among the six treatments. These results suggest the importance of targeted pre-germination treatments to enhance seedling establishment in reforestation efforts. Earlier, Ogungbesan et al. [16] carried out a study on the effect of hot water on the germination rate of *C. siamea* seeds, and reported that hot water ( $100^\circ C$ ) has shown to effectively break seed dormancy in *C. siamea* seeds, indicating a brief treatment time (12 min) is sufficient for dormancy removal. Whereas, in the present study, treatment T3 immersion in hot water ( $80^\circ C$ ) for 10 min had just 10% of seed germination for *C. siamea*. This can be attributed to the variation in hot water temperature and seed soaking time.

Further, in this study we found that the carbon stock potential in wood of *C. siamea* across 11 stem size classes had significant variation ( $p < 0.001$ ). However, no relationship was observed between the wood carbon stock and stem size class of *C. siamea*. We point that *C. siamea* might exhibit non-linear growth patterns concerning carbon accumulation. While larger trees generally have more biomass and thus more carbon stock [17], the rate of carbon accumulation might not directly correlate with stem size.

Factors such as age, health, and environmental conditions could influence carbon stock rates independently of stem size. As trees age, their root systems often experience increased growth, enabling them to delve into fresh soil volumes and extract nutrients from deeper layers [5]. Trees allocate resources differently at various growth stages. Smaller trees might allocate more resources to vertical growth and foliage development rather than radial growth, which contributes more significantly to wood carbon stock. This allocation strategy could lead to a lack of correlation between stem size and carbon stock. Understanding these dynamics is crucial for accurately estimating forest carbon stocks and informing sustainable forest management practices.

Worldwide lots of measures have been taken to curb the rising atmospheric carbon concentration, such as Clean Development Mechanism (CDM) forestry projects began from 2006 in light of carbon offsetting targets, then, the Climate Action Reserve (CAR) projects, Verified Carbon Standard (VCS) and the American Carbon Registry (ACR) came into action [18]. Although, several policies (carbon tax and subsidy) and carbon market business were taken, the rise in atmospheric carbon concentration and its consequence are at alarming rate [17].

India has the second-largest road network globally, following the United States [4]. It is reported that, India's road network spans 5.5 million kilometers, serving 90% of total passenger traffic, and is expanding at a rate of 26.93 kilometers per day[13]. Hence, planting roadside trees, particularly *C. siamea* is one of the best options in India to reduce the atmospheric carbon loads, besides afforestation and reforestation programs. Although *C. siamea* is an exotic tree, it is proved to be a good candidate for restoration in mining area [19], even it is better performer than the native tree species *Albizia lebbbeck*(L.) Benth. (commonly known as Indian siris) in terms of biomass carbon stock, soil carbon stock and soil available nitrogen stock in post mining lands of Jharia Coalfield in Jharkhand, India[5].

Overall, this research contributes valuable insights into optimizing seedling establishment and enhancing carbon stock potential in *Cassia siamea*, supporting efforts to mitigate climate change through afforestation initiatives for the nation.

#### 4. CONCLUSION

The present study concludes that seed germination of *C. siamea* was found high (95%) for the treatment, T5 (immersion in concentrated H<sub>2</sub>SO<sub>4</sub> (80%) for 20 min). Among the eleven stem size classes, carbon stock potential was high for the 55-60 cm size class (56.43±0.34%) for the wood sample. This study contributes valuable insights into optimizing seed germination protocols for *C. siamea* and understanding its carbon stock potential. The findings of this study offer invaluable guidance to policymakers and practitioners regarding the pivotal role of *C. siamea* in bolstering carbon sequestration and fortifying climate resilience, especially in areas grappling with environmental degradation and shifts in land use. By elucidating the mechanisms through which *C. siamea* contributes to carbon storage and ecosystem stability, this research equips decision-makers with actionable insights to implement targeted strategies for sustainable land management and mitigation of climate-related challenges. Ultimately, embracing the potential of *C. siamea* stands as a promising avenue towards fostering ecological health and resilience in vulnerable regions worldwide.

#### REFERENCES

1. Ganesh KP, Pragasan LA. Effects of nitrogen addition on Eucalyptus globulus growth and carbon sequestration potential under various CO<sub>2</sub> climatic conditions. *Geology, Ecology, and Landscapes*. 2024;8(2):185-93.
2. Fernández-Martínez M, Vicca S, Janssens IA, Ciais P, Obersteiner M, Bartrons M, Sardans J, Verger A, Canadell JG, Chevallier F, Wang X. Atmospheric deposition, CO<sub>2</sub>, and change in the land carbon sink. *Scientific Reports*. 2017;7(1):9632.
3. Frianto D, Sutrisno E, Wahyudi A, Novriyanti E, Adinugroho WC, Yuniarto AS, Kurniawan H, Khotimah H, Windyoningrum A, Dharmawan IW, Tata HL. Carbon stock dynamics of forest to oil palm plantation conversion for ecosystem rehabilitation planning. *Global Journal of Environmental Science and Management*. 2024 May 1.
4. Ragula A, Chandra KK. Tree species suitable for roadside afforestation and carbon sequestration in Bilaspur, India. *Carbon Management*. 2020;11(4):369-80.
5. Mukhopadhyay S, Masto RE. Comparative evaluation of *Cassia siamea* and *Albizia lebbbeck* for their potential to recover carbon and nutrient stocks in a chronosequence post-mining site. *Catena*. 2022;208:105726
6. Agus C, Putra PB, Faridah E, Wulandari D, Napitupulu RRP. Organic carbon stock and their dynamics in rehabilitation ecosystem areas of post open coal mining at tropical region. *Procedia Engineering*. 2016;159:329-337.
7. Mukhopadhyay S, Maiti SK, Masto RE. Use of Reclaimed Mine Soil Index (RMSI) for screening of tree species for reclamation of coal mine degraded land. *Ecological Engineering*. 2013;57:133-142.
8. Karthick A, Pragasan LA. Stand structure and above-ground biomass of two tree plantations at Bharathiar University, Coimbatore. *Indian Forester*. 2014;140: 29-33.
9. Pragasan LA. Assessment of tree carbon stock in the Kalrayan hills of the Eastern Ghats, India. *Walailak Journal of Science and Technology*. 2015;12(8): 659-670.
10. Kuyah S, Muthuri C, Wakaba D, Cyamweshi AR, Kiprotich P, Mukuralinda A. Allometric equations and carbon sequestration potential of mango (*Mangifera indica*) and avocado (*Persea americana*) in Kenya. *Trees, Forests and People*. 2024;15:100467.
11. Fayolle A, Doucet JL, Gillet JF, Bourland N, Lejeune P. Tree allometry in Central Africa: Testing the validity of pantropical multi-species allometric equations for estimating biomass and carbon stocks. *Forest Ecology and Management*. 2013;305:29-37
12. Basuki TM, Van Laake PE, Skidmore AK, Hussin YA. Allometric equations for estimating the above-ground biomass in tropical lowland Dipterocarp forests. *Forest Ecology and Management*. 2009;257(8):1684-94.
13. Solanki HK, Ahamed F, Gupta SK, Nongkynrih B. Road transport in Urban India: Its implications on health. *Indian Journal of Community Medicine*. 2016;41(1):16-22.
14. Allen SE, Grimshaw HM, Rowland AP. Chemical analysis. *Methods in plant ecology*. Blackwell Scientific Publications. 1986:285-344.

15. Vidhya, R. and Pragasan, L.A. Age effects of millet crops on phytolith carbon sequestration. *Applied Ecology and Environmental Sciences*. 2022;10(12):747-753.
16. Ogungbesan AM, Ogunsola OR, Lamidi A, Ishiaku YM. Effect of Hot Water on the Germination Rate of *Cassia siamea* L. Seeds. *Agricultural and Food Science Journal of Ghana*. 2017;10(1):780-6.
17. Pragasan LA. Assessment of carbon stock of tree vegetation in the Kolli Hill forest located in India. *Applied Ecology and Environmental Research*. 2016;14(2):169-83.
18. Pearson TR, Brown S, Sohngen B, Henman J, Ohrel S. Transaction costs for carbon sequestration projects in the tropical forest sector. *Mitigation and Adaptation Strategies for Global Change*. 2014;19:1209-22.
19. Ilyas S. Carbon sequestration and growth of stands of *Cassia siamea* Lamk. in coal mining reforestation area. *Indian Journal of Science and Technology*. 2013;6(11):5405-10.

UNDER PEER REVIEW