

Polymorphic survey of KMR-3R and Kasalath (*Oryza sativa* L.) by using SSR markers and morphological traits.

Comment [11]: PARENTAL POLYMORPHISM SURVEY BETWEEN RICE VARIETIES FOR PHOSPHOROUS STARVATION TOLERANCE GENE USING SSR MARKERS

Abstract: ~~Low phosphorus stress~~ In sufficient soil phosphorous has ~~been the most~~ become more important abiotic factor affecting rice yield under ~~present changing climate~~ rice grown conditions. KMR-3R is ~~one of the a~~ popular restorer line used ~~for~~ commercial rice hybrids production ~~in India~~. ~~For majority of the rice ecosystems, M~~most of the rice hybrids released ~~for the diverse ecosystem conditions~~ using three-line systems ~~were suited~~ for normal phosphorus conditions ~~only~~. ~~Consequently, these derived hybrids~~ ~~which exhibits severe yield reduction~~ ~~suffer drastic yield decline~~ under low phosphorus conditions. To facilitate the efficient use of complex polygenic traits in hybrid rice, molecular breeding ~~research, approaches we developed~~ ~~the were developed using~~ KMR-3R BILs with *Pup1* QTL by introgression of major QTL *Pup1* controlling grain yield under low phosphorus stress ~~into KMR-3R~~. To begin marker assisted back cross breeding for introgression of a desirable trait into a variety from the donor genotype, study of parental polymorphism is a pre-requisite. Unless the parents are polymorphic for the traits of interest, the further selection of plants carrying the traits of interest is not possible in the progenies. We screened all reported SSR primers linked to low phosphorus tolerance and observed K-46-1 and K-46-2 linked to *Pup1* QTL was polymorphic between Kasalath and KMR-3R. Simultaneously we screened 300 full-genotype SSR markers covering entire genome and observed 86 SSR primers as polymorphic between Kasalath and KMR-3R. After polymorphic survey, we evaluated Kasalath and KMR-3R under low phosphorus stress and normal phosphorus conditions and recorded significant differences among the parents. Kasalath performed well when compared to KMR-3R for all the characters studied under low phosphorus stress conditions. Construction of a Linkage Map could be ensued procuring the generated genotyping data which could further avail QTL analysis and identification of markers linked to low phosphorus stress.

Comment [12]: Plz modify the sentence

Comment [13]: Plz modify the sentence

Comment [14]: Pl change the sentence

Comment [15]: Need sentence modification

Keywords: Rice, KMR-3R, low phosphorus stress, Yield, BILs and Marker assisted backcross breeding.

INTRODUCTION

Rice is one of the most important food crops for more than half of the world's population. Over 90% of the global rice production is in Asia, ~~and it which~~ contributes 20% of total caloric intake on a daily basis. The green revolution has revolutionized the world's rice production in the last six decades, and the production of rice is equivalent to the population growth in India. With an average annual population growth rate (1.5%) and an average daily rice consumption of 250g per person, the demand for rice is expected to reach at least 140 million tonnes (Mt) by 2050. By 2050, India's population is going to need production of rice to rise by 2.5 M T annually (<http://www.fao.org/rice2004/en/pdf/khush.pdf>). Worldwide, the rice cultivation is affected due to the scarcity of natural resources like land, water, and nutrients coupled with climatic change. The global problem of climate change is currently affecting rice cultivation worldwide by influencing resources like land, water, and nutrients to become scarcer. The development of rice cultivars resistant to climate change is urgently needed. As of now, as many as 8000 QTLs governing various traits have been identified and flanking/tightly linked/functional markers for these traits can be used to transfer into elite varieties through MAS. However, identification of background markers for each pair of donor and recipient parental varieties is very laborious and time consuming ~~also~~.

Comment [16]: Plz modify the sentence

Comment [17]: Plz modify

Comment [18]: Need major modification

Hence, the current study was carried out to identify ready to use SSR markers for background selection of two widely used donor and recipient varieties. One of the readily available and economically viable technological options for meeting the projected global demand of rice is exploitation of heterosis through large-scale cultivation of rice hybrids. To increase the nation's rice yield, hybrid rice technology is essential. According to the AICRIP progress report 2021, 127 rice hybrids have been certified for commercial cultivation in India as of now. In Kharif 2017, 3.5 million tonnes of hybrid rice were produced on 3 million hectares of land. Majority of the rice hybrids were developed for irrigated ~~conditions—ecosystems~~. Currently, three line hybrid seed production using CMS system was widely used for exploitation of hybrid vigour in rice. Phosphorus (P), one of the most important macronutrients necessary for healthy rice plant growth and development, is one of the most deficient elements in rice soils, and its low availability in the soil is one of the major yield-limiting factor in rice production (Wissuwaet *al.*, 1999). Global demand for P fertilizer is steadily rising, despite the fact that commercial phosphate supplies are expected to run out in a few decades. In India, soils are either low (49.3% of soils) or medium (48.8% of soils) in terms of available P necessitating the

Comment [19]: Pl modify

Comment [110]: Use latest data

country's importation of phosphorus-based fertilizers on a massive ~~scale~~. As a result, it is critical to be prepared with alternate solutions to this problem, such as improved crop residue management, adoption of integrated nutrient management, and ~~creation-development of~~ of low soil P tolerant rice cultivars to handle low soil P levels. Genetic enhancement of rice ~~tolerance~~ to P-limiting soils should be one of the focus areas of rice research and development in order to reduce the use of phosphatic fertilizers, which is required not only to increase farmers' income by lowering fertilizer costs, but also to maintain rice production. Therefore, creating hybrids resistant to low phosphorus stress will aid in preventing losses in quantum yield. Though conventional plant breeding made significant impact on increasing the yield coupled with resistance to various biotic and abiotic stresses for target environments, these efforts are slow, need expertise labour and resource intensive.

Comment [111]: Reference plz

Before initiating marker assisted selection or marker assisted backcross breeding for transferring a desired trait into a variety from the donor parent, an analysis of parental polymorphism is required. Further selection of plants possessing the traits of interest is not achievable in the progenies unless the parents are polymorphic for the traits of interest. Polymorphism is an index of genetic variation that shifts with different sets of parents. According to the survey of parental polymorphism, there is demonstrable variation between parents. Due to their many advantageous features, microsatellites (SSRs) have been used more than any other kind of molecular marker. These include having a large genomic distribution, being co-dominant and highly polymorphic, and being amplified by PCR. Through their ability to facilitate precise transfer of specific genomic regions, these markers have also demonstrated significant promise in enhancing breeding efficiency.

SSRs have been widely utilized ~~by-in~~ most crop ~~plants-species~~ for which sequence information is available, including rice. ~~Many genetics related domains, which have wider applications like such-as~~ gene tagging, genetic diversity research, population genotyping mapping, linkage map creation, tracking marker-trait associations, single marker analysis, and quantitative trait loci (QTL) mapping ~~etc., can benefit from these~~. Finding the relevant polymorphic SSR markers for the donor and recipient parents was the goal of the current investigation. After that, these markers were included to a backcross breeding experiment using ~~marker support~~, which was used to ~~insert-introgress~~ low phosphorus deficiency tolerance QTLs

Comment [112]: ?

from the donor parent Kasalath into the background of the extensively used restorer line of hybrid rice *i.e.*, KMR-3R.

MATERIALS AND METHODS

The parents used for the study were KMR-3R and Kasalath and their characteristics were mentioned below.

Comment [113]: The details of the parents used for the study were presented below

1. **KASALATH**: Kasalath is a low P tolerant indica rice genotype harboring major effect QTL *Pup1* controlling grain yield under phosphorus stress.
2. **KMR-3R**: KMR-3R is the restorer line of the popular hybrid KRH2.

Comment [114]: Need not upper case

Comment [115]: italics

Leaf sample collection and DNA isolation

Total genomic DNA was isolated from the parental lines using the method of Zheng et al. (1991) with modifications. Healthy leaf pieces (2-3 cm) from the youngest leaves of a 20-25 day old rice seedlings were taken into the well of a spot testplate (M/s Thomas Scientific, USA). 400 µl of extraction buffer (50 mM Tris HCL, pH 8.0, 25 mM EDTA, 300 mM NaCl and 1 % SLS) was added to the well. The leaf tissues were macerated for about 15-20 seconds till it was completely homogenized. Another 400 µl of extraction buffer was added to the well containing the leaf sample. Using a micropipette of 1 ml capacity, the entire content from the well was transferred to a 1.5 ml capacity micro centrifuge tube. About 400-500 µl Phenol (pH 8.0) : Chloroform : Isoamyl alcohol (25:24:1) was then added to the micro centrifuge tube. The contents were mixed well by inversion for about 10 minutes and centrifuged at 12000 rpm for about 10 minutes at room temperature. After the centrifugation, the supernatant was aliquoted from the micro-centrifuge tube without disturbing the intermediate layer into a fresh 1.5 ml micro-centrifuge tube. To the clear supernatant, 5-10 µl of RNAase (10 mg per ml) was added and incubated for 45-60 minutes at room temperature (~ 37°C). After completion of the incubation period, about 500 µl of Chloroform: Isoamyl alcohol (24:1) was added to the micro-centrifuge tube and mixed well by inversion for 10 minutes. The content was centrifuged at 12000 rpm for 10 minutes and the supernatant was collected into another fresh 1.5 ml micro-centrifuge tube without

Comment [116]: degree

disturbing the intermediate layer. To the clear supernatant equal volume (~500-600 ml) of chilled isopropanol was added. The contents were mixed gently and centrifuged at 12000 rpm for 10 minutes at room temperature. The supernatant was drained gently without disturbing the DNA pellet. About 150-200 µl of 70 % ethanol was added to the pellet at the bottom of the micro-centrifuge tube. It was ensured that the pellet was completely immersed in 70% alcohol. Then the content was centrifuged at 5000 rpm for 5 minutes at room temperature. After centrifugation, the supernatant was drained and 70% alcohol wash was repeated once again. Finally, after centrifugation and draining out the supernatant, the pellet was left overnight air drying at room temperature. After complete drying of the pellet, depending upon the size of the pellet, about 50-100 µl of sterile distilled water was added for dissolving the pellet. About 3 µl of the DNA sample was taken and loaded in 0.8 % ethidium bromide stained agarose-TBE (concentration of TBE) gel, electrophoresed for about an hour and the bands of genomic DNA were visualized and documented in a UV gel documentation system (M/s Alpha Innotech Corporation, USA).

Comment [117]: follow uniformly

Polymerase Chain Reaction (PCR) using SSR markers

The genomic DNA of parental lines subjected to PCR amplification as per the procedure described by Chen (1997). PCR amplification was carried out in a thermocycler using the temperature profile, followed by rapid cooling to 4°C prior to analysis. For electrophoretic studies, The SSR-PCR products were resolved in a 3% Agarose gel prepared in 0.5 X TBE buffer stained with Ethidium Bromide. The gel was run at a constant voltage of 90 V for 1 h and visualized under UV light and documented using gel documentation system. The temperature specifications for the denaturation of DNA strands, Annealing of primers and extension steps are as following .

S.No.	Steps	Temperature	Time
1	Initial denaturation	94 ⁰ C	5 min
2	Denaturation	94 ⁰ C	30 sec
3	Annealing	55 ⁰ C	1 min
4	Extension	72 ⁰ C	1 min
5	Final extension	72 ⁰ C	7 min
6	Cooling	4 ⁰ C	A

Agarose gel electrophoresis and image visualization

PCR products were analyzed by electrophoresis using a 3.0% agarose gel. About 9.0 g of agarose was weighed and applied to a conical flask and 400 ml of 1X TAE buffer was added and mixed well. The ingredients were boiled slowly in the microwave by stirring occasionally. Complete melting of the agarose and a clear solution were maintained during the process. To clean the gel cast plate, it was soaked in water and then wiped with ethanol. After cooling the agarose to room temperature, two milliliters of ethidium bromide (10 mg/ml) was added to the melt and the mixture was then poured onto a gel casting tray with the necessary gel combs and allowed to settle at 20°C for 30 minutes. The gel was transferred to an electrophoresis apparatus containing 1X TAE buffer. Before loading, PCR-amplified products were mixed with 1/6 volumes of gel loading dye (ethidium bromide) and loaded into wells. A 100 bp DNA ladder was added to one well to determine the size of the amplified fragments. Ethidium bromide-stained gels were documented in a UV gel documentation system (Syngene) and the subsequent banding pattern was evaluated.

Comment [118]: follow uniformly

Statistical Analysis

The percentage of polymorphism was calculated by using the formula:

$$\text{Polymorphism (\%)} = \frac{\text{Number of molecular markers showing polymorphism}}{\text{Total number of molecular markers run}} \times 100$$

Graphical representation of the genome

Graphical representation of molecular marker data was performed using Graphical Genotype (GGT2.0) software.

Phenotypic evaluation of IR58025B and Apo in field under control and moisture stress conditions

Comment [119]: why moisture stress condition

In normal phosphorus experiment, regular irrigation and flooded conditions were maintained until maturity of the crop and soil was maintained with P > 18.3 ppm. In low phosphorus experiment, regular irrigation and flooded conditions were maintained until maturity

Comment [120]: write the standard procedure and add reference

of the crop and soil was maintained with $P < 2$ ppm. The crop was harvested when the grains reached physiological maturity. Low 'P' level in the field was maintained without external application of P fertilizers. The levels of phosphorus in the field were estimated from the collected soil sample in each season through Olsen P method before taking up of planting of experimental breeding material in the field plot. The differences between the parents on yield and drought tolerated traits were tested using paired t-test.

Results and Discussion

Genotypic polymorphism by molecular markers

The presence of two or more discontinuous genotypes or alleles on a specific locus in a given population is referred to as genetic polymorphism. In the present study, clear polymorphism was observed between the parents both for full genome coverage markers and drought tolerant markers. A total of 300 full genome coverage SSR markers (forward and reverse) and 10 phosphorus uptake yield QTL markers reported by Chin *et al.* (2010) were analyzed for molecular polymorphism. Polymerase Chain Reaction was done to check each sample for parental polymorphism using specific Rice Microsatellite (RM) primers. Out of 300 full genome coverage primers used the parents showed polymorphic only to 86 RM primers listed in Table 1 were polymorphic whereas K-46-1 and K-46-2 markers linked to *Pup1* QTL listed in Table 2 were polymorphic between the parents for low phosphorus deficiency tolerant markers. These are used in foreground selection. This type of parental polymorphic survey was done by previous workers *i.e.*, Collard *et al.* (2006), Chin *et al.* (2010), Sarkar *et al.* (2011), Tyagi *et al.* (2012), Aluwihare *et al.* (2014), Goncharova and Kharitonov (2016), Neelam *et al.* (2017), Zhang *et al.* (2018) and Lapitan and Mercado (2023), before starting marker assisted backcross breeding for introgression of desired yield controlling *qDTYs* under moisture stresses in rice. Chromosome wise distribution of polymorphic markers used in this study was presented in the Table 3. Percentage of polymorphism was highest on 1 chromosome and least on 8 chromosomes. The average per cent polymorphism on all the chromosomes was 28.74. The lack of detectable polymorphism between two parents could be due to the fact that both parents, Kasalath and KMR-3 Rare indica lines. Lack of molecular marker polymorphism between indica genotypes has been earlier noticed in studies by Xu *et al.* (2002) and Biradar *et al.* (2004). In the present study dinucleotide repeats in the microsatellite region showed more polymorphism than tri and tetra nucleotide repeats. It may be due to the abundance of dinucleotide repeats distributed throughout

Comment [121]: aim is drought tolerance or low p tolerance
drought stress means follow IRRI SES Protocol
all drought parameters according to SES should be evaluated

Comment [122]: Plz modify

Comment [123]: Pl modify

the genome of rice. The prevalence of dinucleotide repeats in parental polymorphism was reported by previous workers. The usage of molecular marker technology in breeding programs has greatly increased its efficiency and fastens the transfer of desirable genes among varieties and to introgress novel genes from related wild species. Polygenic markers which were previously difficult to analyze using traditional breeding methods, would now be easily tagged using molecular markers. The screening of markers for parental polymorphism among the rice cultivars forms the basis for tagging of the desired gene, fine mapping of the gene in the rice chromosome and in the subsequent Marker Assisted Selection (MAS) programmes.




Fig 1 : Parental polymorphism shown by K-46-1 and K-46-2 for *Pup1 QTL*

Comment [124]: Moisture stress or p stress ??

Phenotypic polymorphism by morphological traits under control and moisture stress conditions

KMR-3R and Kasalath were assessed for grain yield and its attributing traits under low phosphorus stress and normal phosphorus conditions (Table 4). KMR-3R recorded significant differences for all traits studied between normal phosphorus and low phosphorus stress conditions. Similarly, Kasalath recorded significant difference for all the traits studied except for grain yield per plant between normal phosphorus and low phosphorus stress conditions. These findings are used for development of low P tolerant backcross inbred lines of KMR-3R by marker assisted backcross breeding.

CONCLUSION

Phosphorus is one of the major plant nutrients and plays a vital role in productivity of rice. It is one of the least abundant of all essential nutrients in soil. The present situation in India in both high and low input farming systems is unsustainable because P is a finite resource. It is difficult to recognize the deficiency of mild to moderate P in the field. As a result, phosphorus deficiency in rice is known as hidden hunger. Application of phosphorus fertilizer can solve this problem. But it is difficult for resource poor farmers in India to apply phosphorus fertilizer every year. Therefore selection of rice cultivars capable of extracting phosphorus from phosphorus limiting soils with higher phosphorus use efficiency is considered as a significant cost-effective solution.

Comment [125]: Increasing the

Fortunately, significant genetic variation has been reported in rice for tolerance to soil phosphorus deficiency (Fageria and Baligar, 1997). A large effect QTL known as *Pup1* controlling grain yield under phosphorus deficiency has been mapped on chromosome 12 of Kasalath rice cultivar (Wissuwaet *al.*, 1998). Marker Assisted Selection (MAS) programmes rely on the identification of parental polymorphism in rice cultivars through the screening of markers. This allows for the subsequent tagging and fine mapping of the target gene within the rice chromosome. Polymorphic markers discovered in this work were used to genotype the entire mapping population and facilitated the introgression of *Pup1* QTL involved in grain production under low phosphorus stress into KMR-3R. This study helped in the development of backcross inbred lines of KMR-3R by introgression of *Pup1* from Kasalath into KMR-3R by marker assisted backcross breeding. These BILs were utilized in development of hybrids tolerant to low phosphorus stress.

Comment [126]: Pl modify

Comment [127]: As mentioned in the title plz mention the morphological trait for P stress tolerance

References

1. Aluwihare, Y.C., Lelwala, R.V., Ishan, M., Sirisena, D.N., Samarasinghe, W.L.G and Sooriyapathirana, S.D.S.S. 2014. DNA marker genotyping for *pup1* locus in rice to identify tolerant varieties under low phosphorus soil conditions in srilanka. *Proceedings of post graduate institute of science research congress*, Srilanka.
2. Biradar S K, Sundaram R M, Thirumurugan T, Bentur J S, Amrudhan S, Shenoy V V, Mishra B, Bennett J and Sharma N P. 2004. Identification of flanking SSR markers for a

major rice gall midge resistance gene and their validation. *Theoretical and applied genetics* 109:1468-1473.

3. Chin, J.H., Lu, X., Haefele, Gamuyao, R., Ismail, A., Wissuwa, M and Heuer, S. 2010. Development and application of gene-based markers for the major rice QTL Phosphorus uptake 1. *Theoretical and Applied Genetics*. 120: 1073-1086.
4. Collard, B.C.Y., Thomson, M., Penarubiai, M., Lu, X., Heuer, S., Wissuwa, M., Mackill, D.J and Ismail, A.M. 2006. SSR analysis of rice near-isogenic lines (NILs) for P-deficiency tolerance. *SABRAO Journal of Breeding and Genetics*. 38(2): 131-138.
5. Goncharova, J.K and Kharitonov, E.M. 2016. Genetic control of traits associated with phosphorus uptake in rice (*Oryza sativa* L.) varieties. *Russian Journal of Genetics and Applied Research*. 6(3): 270-278.
6. <http://www.fao.org/rice2004/en/pdf/khush.pdf>.
7. Lapitan, V.C and Mercado, M.J.T. 2018. Genetic improvement of upland rice through marker assisted selection (MAS) for tolerance to phosphorus deficiency. <http://www.uplb.edu.ph>
8. Murray, H.G and Thompson, W.F. 1980. Rapid isolation of high molecular weight DNA. *Nucleic Acids Research*. 8: 4321-4325.
9. Neelam, K., Thakur, S., Neha, I., Yadav, S., Kumar, K., Dhaliwal, S.S and Singh, K. 2017. Novel alleles of phosphorus-starvation tolerance 1 gene (*PSTOL1*) from *Oryza rufipogon* confers high phosphorus uptake efficiency. DOI: 10.3389/fpls.2017.00509.
10. Sarkar, S., Yelne, R., Chatterjee, M., Das, P., Debanth, S., Chakraborty, A., Mandal, N., Bhattacharya, K and Bhattacharya, S. 2011. Screening for phosphorus(P) tolerance and validation of *Pup-1* linked markers in indica rice. *Indian Journal of Genetics*. 71(3): 209-213.
11. Tyagi, W., Rai, M and Dohling, A. 2012. Haplotype analysis for *pup1* locus in rice genotypes of north eastern and eastern india to identify suitable donors tolerant to low phosphorus. *SABRAO Journal of Breeding and Genetics*. 44 (2): 398-405.
12. Xu W, Virmani S S, Hernanadez J E, Sebastian L S, Redona E D and Li Z. 2002. Genetic diversity in the parental lines and heterosis of the tropical rice hybrids. *Euphytica*. 127: 139- 148.
13. Zhang, Y., Anis, G.B., Wang, R., Wu, W., Yu, N., Shen, X., Zhan, X., Cheng, L and Cao,

L. 2018. Genetic dissection of QTL against phosphate deficiency in the hybrid rice 'Xieyou9308'. *Plant Growth Regulators*. 84: 123-133.

Table 1 Microsatellite markers exhibiting polymorphism between KMR-3R and Kasalath

1	RM 10694	21	RM 250	41	RM 3558	61	RM 408	81	RM 7391
2	RM 8094	22	RM 29	42	RM17486	62	RM 331	82	RM 209
3	RM 3412	23	RM 3316	43	RM6909	63	RM 256	83	RM 457
4	RM 140	24	RM 262	44	RM2275	64	RM 447	84	RM 2136
5	RM 10794	25	RM 530	45	RM127	65	RM 126	85	RM 28048
6	RM 10825	26	RM 424	46	RM5270	66	RM 342	86	RM 571
7	RM 562	27	RM 569	47	RM 413	67	RM 23679	87	RM 28199
8	RM 10927	28	RM 3766	48	RM 169	68	RM 23911	88	RM 27100
9	RM 543	29	RM 15087	49	RM 5140	69	RM 24173		
10	RM 486	30	RM 231	50	RM 3638	70	RM 24179		
11	RM 315	31	RM 468	51	RM 3321	71	RM 24390		
12	RM 472	32	RM 16232	52	RM 204	72	RM 201		
13	RM 12023	33	RM 16236	53	RM 6836	73	RM 105		
14	RM 165	34	RM 523	54	RM 3827	74	RM 26076		
15	RM 8051	35	RM 15914	55	RM20546	75	RM 286		
16	RM 312	36	RM 15087	56	RM 20801	76	RM596		
17	RM 12611	37	RM 545	57	RM 21330	77	RM 6132		
18	RM 145	38	RM 551	58	RM 21427	78	RM 206		
19	RM 6374	39	RM 16616	59	RM 5481	79	RM 590		
20	RM 3275	40	RM 5757	60	RM 248	80	RM 286		

Table 2 Details of polymorphic primer used for foreground selection of *Pup1* QTL

QTL	Chromosome	Position	Primer
<i>PUP1</i>	12	523 bp	K-46-1
<i>PUP1</i>	12	227 bp	K-46-2

Table 3 Chromosome wise polymorphic primers between KMR-3R and Kasalath

Chromosome	No. of SSR markers used	No. of polymorphic markers on each chromosome	No. of monomorphic markers on each	Percentage of primers showing polymorphism on each

			chromosome	chromosome
1	35	14	21	40.00
2	30	6	24	20.00
3	28	9	19	32.14
4	25	5	20	20.00
5	25	7	18	28.00
6	28	8	20	28.57
7	25	7	18	28.00
8	27	5	22	18.52
9	20	8	12	40.00
10	19	5	14	26.32
11	20	6	14	30.00
12	18	6	12	33.33

UNDER PEER REVIEW




Table 4 Comparison of KMR-3R and Kasalath for grain yield and its attributes under normal P and low P stress conditions

S.No.	Trait	KMR-3R			Kasalath			S.E. of difference
		Normal P (N.P.) (n=5)	Low P stress (L.P.) (n=5)	Mean difference (N.P-L.P)	Normal P (N.P.) (n=5)	Low P stress (L.P.) (n=5)	Mean difference (N.P-L.P)	
1	Plant height (cm)	115.12	105.12	10.00**	94.24	83.19	11.05**	1.16
2	Days to 50% flowering	103.00	110.00	-7.00**	85.00	90.00	-5.00**	0.64
3	Days to maturity	128.00	130.00	-2.00**	115.00	120.00	-5.00**	0.58
4	No. of tillers per plant	15.00	8.00	7.00**	16.00	10.00	6.00**	0.37
5	No. of panicles per plant	12.00	7.00	5.00**	14.00	8.00	6.00**	0.35
6	Panicle length (cm)	25.10	19.92	5.18**	23.62	21.24	2.38**	0.64
7	1000 grain weight (g)	24.12	20.21	3.91**	30.56	21.90	8.66**	0.55
8	No. of filled grains per panicle	150.00	74.00	76.00**	135.00	98.00	37.00**	1.24
9	No. of unfilled grains per panicle	8.00	58.00	-50.00**	35.00	48.00	-13.00**	0.96
10	Spikelet fertility (%)	94.93	56.06	38.87**	79.41	65.77	13.64**	0.60
11	Grain yield per plant (g)	24.24	12.41	11.83**	19.12	18.04	1.08	0.71
12	Biological yield per plant (g)	60.14	50.14	10.00**	70.21	65.21	5.00**	0.98
13	Harvest index (%)	40.30	24.75	15.55**	27.23	27.66	-0.43**	1.25
14	Leaf phosphorus content (%)	0.50	0.30	0.20**	0.90	0.80	0.10**	0.03
15	Seed phosphorus content (%)	1.20	1.00	0.20**	1.60	1.50	0.10**	0.03

* Significant at 5% level;

** Significant at 1 % level

UNDER PEER REVIEW

