

IMPORTANCE OF MINERALS IN SILKWORM NUTRITION: A REVIEW

ABSTRACT

The amount of silk produced by silkworms is influenced by several of factors. The silkworm's nutrition is the most important element. Good quality cocoon and silk production is dependent on the healthiness of the silkworm. Health, growth and resistance to biotic factors are directly dependent on the nutrition of the silkworm which is derived from the food. Consuming a well-balanced nutritious food is particularly important for silkworms, since they only consume throughout their larval period. Among the different nutritional requirements by the silkworm minerals plays an important role. Minerals are not synthesized within insects, although they are essential elements and affect various metabolic processes. Minerals affect the growth and development of larva, economic cocoon characters and silk yield. The many functions of the neuromuscular, reproductive and hormonal systems among others can be efficiently controlled by adding micronutrients and trace elements such as minerals to the diet.

Key words: Silkworm; nutrition; mineral; cocoon characters and silk yield

1. INTRODUCTION:

Insects and other animals have similar nutritional needs. The prime component of a living being's is good growth and development and its dietary intake. One of the most important factors for the growth and development of insect species is its nutritional requirement and metabolism in the body [1]. The nutrition of silkworm has been of primary importance in sericulture, because not only good growth of the larva and survival but also cocoon and egg production has been known to be influenced by the nutritive value of the nourishment [2]. The quality of food directly influences the growth and development of the insect and ultimately the quantity and quality of product produced by them [3,4].

Mulberry silkworm is a monophagous in nature among the four commercial silkworms such as eri, muga and tasar. Silkworms are voracious feeders to fulfill their nutritional requirement and all of them solely depend on their nutritious host leaves for healthy growth and development. Any nutritional imbalance in their food results in poor larval development. Poor and unhealthy larval development is the result of nutritional imbalance in their diet.

Quantitative requirement for each nutrients and the required balance of nutrients for optimum level of nutrition may vary within and between species owing to many factors including synthetic ability of the insect and metabolic activities involving specific interactions between certain nutrients [5]. Mineral salts played an important role in the nutrition of silkworm [6]. In relation to other nutritional needs, mineral nutrition has received less attention and it is unknown how much of a quantity insects need [7]. Minerals are important components of the silkworm diet and play an important role in regulating the osmotic pressure of intracellular and extracellular fluids and participate as cofactors in various enzyme systems [8]. Micronutrients are required in small quantities but are essential for insect nutrition. Mineral affects not only yield components significantly, but also food consumption, co-efficient of utilization and larval development [9]. Potassium, sodium, phosphorus, magnesium, manganese, zinc, and copper are important for the growth and development of insect [10]. Lack or deficiency of a micronutrient has a significant negative impact on insect biology. Larvae of *Bombyx mori* L. are known to require appreciable amounts of potassium (8000-9000 ppm),

Comment [h1]: Add scientific names as well.

Comment [h2]: Add few lines on how silkworms derive essential nutrients from mulberry leaves and their impact on growth and development of silkworm, *B. mori*. Take reference from paper, "Impact of sericin fortified mulberry leaves on the cocoon traits of silkworm *B. mori*"

phosphate (2000-6000 ppm) and magnesium (1000 ppm) [11]. It has been reported that 28% of the silkworm larval structure at different stages includes absorbed minerals [12]. By supplementing the diet with micronutrients and trace elements *i.e.* minerals, the various functions of the hormonal system, neuromuscular system, and reproductive system can be modulated effectively [13]. Therefore, studies related to qualitative and quantitative requirement of minerals in silkworm is essential for improvement of the diet of silkworm.

In terms of nutrition, a mineral is a chemical element that an organism needs as an essential nutrient in order to successfully carry out functions which are important to life. They are found in ionized form in the body of an organism. Among the different minerals, metals and their salt cations play an important role as catalyst on the structural components of large molecules with a specific function which are crucial for the better functioning of life [14].

Minerals are classified as macro and micro minerals. Calcium, phosphorus, potassium, sodium, magnesium, sulfur and chlorine are the macro minerals. Macrominerals are present in larger quantities in the animal's body or are needed in larger quantities in food. Micro minerals such as iron, iodine, cobalt, copper, zinc, manganese, molybdenum, selenium, chromium and fluorine are also known as trace elements since they are needed in lesser amounts in an animal's food or are found in the human body in low proportions. In the present study, the necessity of several minerals for the nutritional needs of silkworms is reviewed.

Comment [h3]: Add these lines in 3rd paragraph initial to introduction of mineral nutrition in silkworm.

2. Utilization of different minerals and their salt cations in silkworm nutrition:

2.1 Calcium (Ca):

Calcium is important for silk formation in silkworm [15]. Feeding of calcium treated (0.1%) mulberry leaves gave the best results regarding larval development and cocoon weight [16]. Egg shell *i.e.*, egg chorion contains a large amount of calcium. *Antheraea assamensis* Helfer, the muga silkworm, used to eat its egg shell after hatching, which is regarded as a positive sign [17].

Comment [h4]: Explain the role of calcium in silkworm physiology and calcium constituents of silkworm. Take reference from paper "Calcium in edible insects and its use in human nutrition."

2.2 Phosphorus (P):

Phosphorus influences growth in most invertebrates and provide support for the growth rate hypothesis [18]. But in the environment phosphorus is extremely limited, often being 10-20 times lower in plants than what invertebrate herbivores require [18]. In this context it is an essential mineral for growth and development of the phytophagous silkworms. Phosphorus increases the growth and economic character of silkworm *B. mori* L [19].

2.3 Potassium (K):

Potassium is an unique element which promotes the growth of silkworm to maximum extent [20]. Haemolymph trehalose and fat body glycogen content increased after feeding with potassium sulphate [21]. The oral supplementation with potassium permanganate and potassium chloride to mulberry silkworm increase the fat body and haemolymph proteins and at the same time the weight of silk gland, cocoon and shell weight are also significantly increased [22]. Potassium nitrate supplementation of mulberry leaves markedly enhanced the silkworm *B. mori's* fat body glycogen, haemolymph trehalose, protein, and lipid levels [23,24]. Fecundity of adult moth is enhanced by feeding potassium iodide in *B. mori* [25].

2.4 Magnesium (Mg):

The overall process of trehalose production requires magnesium [26]. Mineral salt of magnesium chloride significantly increased the fat body glycogen, protein, total lipids and haemolymph protein and trehalose in the silkworm *Bombyx mori* [27]. Magnesium ion is important for silk formation in silkworm [15]. Supplementation of mulberry leaves with magnesium sulphate has been found to enhance the commercial characters of *B. mori* at 0.25 and 0.50% concentrations [28].

2.5 Iron (Fe):

Iron is needed by insects for several kinds of important physiological processes, including neuronal function, detoxification, DNA synthesis, and cellular respiration [29]. It increases the growth of silkworm as well as the economic characters of cocoon in *B. mori* [30].

2.6 Cobalt (Co):

Cobalt is essential for the various physiological processes in insects, including immune function and resistance against the pathogens [31]. Cobalt sulphate increases the rate of protein synthesis in early stages in the larvae of *Samia ricini* [32]. The mulberry silkworm's silk glands' lipid composition is influenced by cobalt chloride [33]. Higher silk yield has been recorded when cobalt was supplemented in the diet of silkworm [34,35].

2.7 Copper (Cu):

Supplementation with copper sulphate to the silkworm feed significantly increased the larval weight, fecundity, cocooning and moth emergence in *B. mori* [25,36]. Silkworm larvae when fed on mulberry leaves treated with 0.2N % + 0.03% K + 0.01% Ca + 0.05% copper concentrations consumed more food, gained more larval weight and produced heavier cocoons [37].

2.8 Zinc (Zn):

Zinc is the only heavy metal with amphoteric properties and dissolves freely in water and bases to form zincates. Since 1930s, zinc has been widely recognized as an important constituent of animal diet and is widely distributed in all animal tissues. Cofactor function of zinc is considered of almost universal importance to organisms and trace elements can be essential for insect in general [13]. It plays a vital role in synthesis of lipids, proteins and carbohydrates and also in reducing the duration of larval and pupal stages of mulberry silkworm, *B. mori* L [33]. The use of zinc positively influences the mass of the silk glands and also increases the weight of cocoon and cocoon shell [8]. It also plays an important role in augmenting the growth and antioxidant protection of the larvae of *Antheraea mylitta*, which may improve the larval fitness, quality and quantity of silk production [38]. Moreover, mulberry leaves fortified with zinc increases the fecundity, quality parameters of cocoons like higher cocoon shell ratio, silk-body ratio, raw silk percentage, denier of the silk. It lowers the floss-shell ratio by decreasing the floss protein synthesis in *B. mori* silkworm [39]. Zinc chloride ($ZnCl_2$), a binary salt of zinc is a potent modulator for overall growth and development of eri silkworm and positively influences the economic parameters of eri cocoon [40].

2.9 Nickel (Ni):

Nickel is also an important element in the nutrition of organisms [41]. Nickel supplementation shortened the larval period and increased significantly all economic parameters of silkworm *B. mori* L. [42,43]. Growth and development of the silkworm enhanced when they reared on host leaves supplemented with nickel chloride. It increases total lipid of the fat body in silkworm *B. mori* [44].

2.10 Selenium (Se):

Selenium is well known for its beneficial effects and it is an essential micronutrient for both human and animals [45]. It is a cofactor in many enzymes protecting the cell against free radicals and maintains high tissue antioxidant level [46]. Use of selenium in silkworm, leads to an improvement of the larval weight, silk gland weight, cocoon weight, shell weight, filament length [8,47].

3. Impact of mineral deficiency in silkworm:

As minerals are essential for healthy growth and development of silkworm, deficiency of these elements cause some negative impact on it. In *B. mori*, lower body weight is the consequence of lower consumption of potassium, calcium, magnesium, and phosphorus [48]. The cocoon and silk characteristics of mulberry silkworms are seriously affected by phosphorus imbalance [49]. Low levels of sulfur-containing amino acids result from a deficiency in sulfur, which lowers protein synthesis and

Comment [h5]: Add different studies conducted on feeding of mineral fortified food to silkworm mentioning their results as well. Present it in tabulated form, it will be useful for the readers to gather all the information pertaining to mineral nutrition in silkworm in this article.

Comment [h6]: This sub- topic is required to be in introduction part. That will be useful for the readers to understand the importance of mineral fortified nutrition to silkworm.

interferes with the synthesis of silk [20]. The altered iron content in mulberry leaf resulted in the reduced larval weight, cocoon weight and silk filament length [50].

4. CONCLUSION:

In pursuance of silkworm nutrition, mineral elements are indispensable to enhance the larval development and cocoon yields but the mode of complementation, right dosage, time of application, number of sprays all have to be examined to secure maximum benefit from such nutrition. If suitable dose and treatment are advocated and practiced at the research level, this can give much needed boost to overall production of the silk and seed. The quality of leaf has got direct influence on the health, growth and survival of silkworm. Though minerals are not synthesized within the insect body, it is very much essential for overall growth and development of the insect. Since the silkworms are the commercially important insect, to improve economic characters and better growth of silkworm, more research works should be done on silkworm food additives such as the minerals.

REFERENCES

1. Pattnaik A. and Pattnaik S. During 5th instar development a correlative study on changes in concentration of haemolymph ascorbic acid of bivoltine races of *Bombyx mori* L. International Journal for Scientific Research and Development. 2017;5(10): 77-81
2. Ito T. Nutritional requirements of the silkworm, *Bombyx mori* L. Proceedings of the Japan Academy. 1967;43:47-67
3. Dutta LC. Effect of castor varieties on growth, nutrition and cocoon characters of eri silkworm, *Samia ricini* Boisduval. Ph.D. Thesis, Assam Agricultural University, Jorhat, Assam. 2000.
4. Legay JM. Recent advance in silkworm nutrition. Annual Review of Entomology. 1958;3:75-86
5. House HL. Insect Nutrition. In: The Physiology of Insect. 1974;5(2): 1-53
6. Akhtar M. and Asghar A. Nutritional requirement of silkworm (*Bombyx mori*). Pakistan Journal of Zoology. 1972;4:101-107
7. Locke M and Nichol H. Iron economy in insects: Transport, Metabolism and Storage. Recent advances in silkworm nutrition. Annual Review of Entomology. 1992;3: 75- 86
8. Bentea M, Sara A, Marghitas L Al, Gabor E, Dezmirean D, Valic B and Creta C. Effect of selenium and zinc supplementation on production parameters and DNA of the mulberry silkworm (*Bombyx mori* L.), baneasa white variety. ARPN Journal of Agricultural and Biological Science. 2012; 7(11): 871-876
9. Javaid MA. Effect of treated mulberry leaves with mineral supplement (N, K) on the development of silkworm and silk yield. M.Sc. Thesis, University of Agriculture, Faisalabad, Pakistan. 1991.
10. House HL. Insect Nutrition. Annual Review of Entomology. 1961; 6:13-26
11. Muniandy S, Sheela M and Nirmala S. Effect of vitamins and minerals (Filibon) on food intake, growth and conversion efficiency in *Bombyx mori*. Environmental Ecology. 2001;13:433-443
12. Ito T. Silkworm nutrition: silkworm an important laboratory tool. (Eds) Tazima Y. Kodansha Ltd., Tokyo. 1978;121-157
13. Devi KL and Yellamma K. The modulatory role of zinc in the silkworm, *Bombyx mori* L. Bioscience Discovery. 2013;4(1): 58-68
14. Murugesha KA, Aruna R and Chozhan K. Effects of Minerals on Growth of Silkworm, *Bombyx mori* L. and Their Impact on Cocoon Economic Parameters. Madras Agricultural Journal. 2020; 108(1-3): 146-152
15. Ochi A, Hossain KS, Magoshi J and Nemoto N. Rheology and dynamic light scattering of silk fibroin solution extracted from the middle division of *Bombyx mori* silkworm. Biomacromolecules. 2002;3: 1187-1196
16. Maqbool S. Effect of feeding calcium and nitrogen treated mulberry leaves on larval development and silk yield of *Bombyx mori* L. M.Sc. Thesis, University of Agriculture, Faisalabad, Pakistan. 1991.

17. Chowdhury, SN. Muga Silk Industry. Directorate of sericulture & weaving, Government of Assam. 1982
18. Visanuvimol L, Bertram SM. How dietary phosphorus availability during development influences condition and life history traits of the cricket, *Acheta domesticus*. *Journal of Insect Science*. 2011;11:63
19. Chakrabarti S, Subramanyam MR, Singhal BK and Datta RK. Nutrient deficiency management in mulberry. Central Sericultural Research and Training Institute, Mysore. 1997; 5-15.
20. Mahadeva A. Nutritive elemental status in mulberry foliage under jassids (*Empoasca flavescens* F.) infection. *Indian Journal of Natural Science*. 2016;7(38): 11537-11539
21. Bhattacharya A and Kaliwal BB. Influence of mineral potassium permanganate on the biochemical constituents in the fat body and haemolymph of the silkworm, *B. mori* L. *International Journal of Industrial Entomology*. 2004;9(1): 131- 135.
22. Dasmahapatra AK, Chakraborti MK and Medda AK. Effect of potassium iodide, cobalt chloride, calcium chloride and potassium nitrate on protein, RNA and DNA content of silkgland of silkworm (*Bombyx mori* L.) nistari race. *Sericologia*. 1989;29: 355-359
23. Goudar KS and Kaliwal BB. Effect of potassium nitrate on the biochemical parameters of the silkworm, *Bombyx mori* L. *International Journal of Industrial Entomology*. 2001;3:93- 96.
24. Kochi SC and Kaliwal BB. The Effects of Potassium Bromide on Biochemical Contents of the Fat Body and Haemolymph of Crossbreed Races of the Silkworm, *Bombyx mori* L. *Caspian Journal of Environmental Science*. 2006;4 (1):17-24
25. Magadam SB. Studies on the effect of some chemicals and hormones on the gonads and the life cycle of the silkworm, *Bombyx mori* L. Ph.D. Thesis, Karnataka University, Dharwad, India. 1987
26. Murphy TA and Wyatt GR. The enzymes of glycogen and trehalose synthesis in silkworm fat body. *Journal of Biological Chemistry*. 1965;240: 1500-1508
27. Bhattacharya A and Kaliwal BB. Synergetic Effects of Potassium and Magnesium Chloride on Biochemical Contents of the Silkworm, *Bombyx mori* L. *Caspian Journal of Environmental Science*. 2005;3(1):15-21
28. Islam MZ and Khan AR. Growth and development of the mulberry silkworm, *Bombyx mori* L. (Lepidoptera: Bombycidae) on feed supplemented with manganese sulphate. *Journal of Biosciences*. 1993;1: 21-30.
29. Gorman MJ. (2023). Iron homeostasis in insect. *Annual Review of Entomology*, 68: 51-67.
30. Horie Y, Watanabe K. and Ito T. Nutrition of the silkworm *Bombyx mori* XVIII. Quantitative requirements for Potassium, Phosphorus, magnesium and zinc. *Bull. Sericul. Expt. Sta*. 1967;22: 181-193
31. Khan S and Lang MA. Comprehensive Review on the Roles of Metals Mediating Insect-Microbial Pathogen Interactions. *Metabolites*. 2023; 13(7):839
32. Padaki PR. Some aspects of physiology of the eri silkworm, *Philosamia ricini*. Ph.D. Thesis, Karnataka University, Dharwad, India. 1991
33. Bhattacharya A and Medda AK. Effect of cyanocobalamin and cobalt chloride on glycogen of silkgland of *Bombyx mori* L. Nistari race. *Science and Culture*. 1981;77:268-270
34. Narasimhamurthy CV and Govindappa S. Effect of cobalt on silkworm growth and cocoon crop performance. *Indian Journal of Sericulture*. 1988;27:45-47
35. Arnaudo M. Effect of cobalt on the life-cycle of the silkworm, *Atti della Societa Italiana delle Scienze Veterinarie*. 1954;8:279-82.
36. Magadam SB, Holli MA and Magadam VB. Effect of the feeding copper sulphate on the economic parameters of the polyvoltine silkworm *Bombyx mori* L. *Sericologia*. 1992;32: 395-399
37. Ashfaq M, Din R, Akram W, Saleem M. and Rasool KG. Effect of feeding optimum doses of nitrogen, potassium, calcium and copper on the development and silk yield of silkworm, *B. mori* L. *Pakistan Entomologist*. 1998;20:104-106

38. Sahu S, Dandapat J and Mohanty N. Foliar supplementation of zinc modulates growth and antioxidant defense system of tasar silkworm *Antheraea mylitta*. *Journal of Entomology and Zoology Studies*. 2015;3(2):25-35
39. Wani MY, Rather RA, Bashir M, Shafi S and Rani S. Effect of zinc on the larval growth and quality cocoon parameters of silkworm (*Bombyx mori* L.): A review. *International Journal of Fauna and Biological Studies*. 2018;5(4):31-36
40. Bora N, Singha A, Gogoi D, Kalita S and Saikia H. The impact of zinc chloride supplementation on larval growth and economic cocoon parameters of eri silkworm. *Journal of Entomological Research*. 2023;46:1108-1113
41. Spears JW. Nickel as a 'newer trace element' in the nutrition of domestic animals. *Journal of Animal Science*. 1984;59(3):823-835
42. Chamundeswari P and Radhakrishnaih K. Effect of zinc and nickel on the larval and cocoon characters of the silkworm, *Bombyx mori* L. *Sericologia*. 1994;34(2): 327-332
43. Islam R, Ali AO, Paul DK, Sultana S, Banu NA and Islam R. Effect of salt, nickel chloride supplementation on the growth of silkworm *Bombyx mori* L. (Lepidoptera: Bombycidae). *Journal of Biological Science*. 2004;4:170-172
44. Saha BN and Khan AR. Effect of nickel chloride, supplementation on the lipid content in *B. mori* L. *Indian Journal of Sericulture*. 1995;34: 156-158
45. Fleet JC. Dietary selenium repletion may reduce cancer incidence in people at high risk who live in areas with low soil selenium. *Nutrition Reviews*. 1997;55: 29-48
46. Beck MA, Handy J and Levander OA. Host nutritional status: the neglected virulence factor. *Trends in Microbiology*. 2004;12: 417-423
47. Smitha S. Impact of selenium on some biochemical and histological aspects of the silkworm *Bombyx mori* L. Ph.D. Thesis. Sri Krishnadevaraya University, Anantapur (A.P.). 2006.
48. Iwanari Y. Bulk farms, Tokyo University of Agriculture and Technology. 1976;7: 62.
49. Radha N, Natarajan T, Muthukrishnan TS and Oblisami G. Effect of antibiotics of mulberry silkworm. *Proceeding on Sericulture*. Cymp. Coimbatore. 1980;173-177
50. Shankar MA. Handbook of Mulberry Nutrition. Published by Shetty, G. P., Multiplex, Karnataka Agro Chemicals. Bangalore. 1997; 19-75