

Original Research Article

Assessing the Impact of Trap Nests on *Megachile* Bee Population in Mung Bean

Abstract

This study explores the impact of nest installation on *Megachile* bee populations, which are potential pollinators of mung bean crops in Uttarakhand, India. Field investigations over two seasons reveal that six *Megachile* species visit mung bean blooms regularly for their provisioning. Assessments performed pre- and post-nest installation show slight changes in *Megachile* abundance and species-specific responses to environmental factors. Post-nest installation *M.lanata* showed the highest abundance (5.13 ± 0.77 bees/m²/5min) followed by *M.umbripennis* (4.76 ± 1.05 bees/m²/5min), although nest occupancy rates were low. Correlation analyses of weather parameters show preferences for specific climatic conditions with temperature having a considerable influence. These findings highlight the intricate relationship between *Megachile* bees and environmental factors, underlining the need for specific pollinator management approaches. Further research into *Megachile* nesting behaviours is essential to enhancing crop pollination and ensuring sustainable agricultural practices.

Keywords: *Megachile* bees, Wood-nesting bees, Nesting behaviour, Pollinator management, Environmental factors.

Introduction

India stands as the leading producer and consumer of pulses on a global scale. This includes 25% of production, and 27% of consumption globally including more than two-thirds of pulse production accounted by the six states viz., Madhya Pradesh, Rajasthan, Maharashtra, Karnataka, Andhra Pradesh, and Uttar Pradesh. Apart from the key states in pulse production, Uttarakhand has shown an ongoing rise in pulse yield from 2015-16, with an expected production of 1069 kg/ha by 2021-22 (ASAG, 2022). Among pulses, mung bean (*Vignaradiata* L. Wilczek) is nutritionally valuable with high protein content and adaptability to diverse climates (Pratap, 2021; Miller, 1989). It is cultivated in broad rows, but reduced row spacing is more effective in weed control (Pratap, 2021). This plant is a member of the Leguminosae family, specifically the Papilionoideae subfamily, they bear tiny yellow or greenish-yellow flowers that cluster into groups of 10-15. Mung beans have a rare trait: they are cleistogamous, which implies that pollination can occur whilst the flowers are closed (Huppertz et al., 2023). However, pollinators facilitate cross-pollination, which benefits legume crops including mung beans, insect pollination has a substantial influence on mung bean production in smallholder systems of farming (Elisante et al., 2020). Family Megachilidae is a significant crop pollinator (Hall and Avila, 2016), among these, those in the genus *Megachile*, are excellent pollinators.

Megachile bees (leafcutter or mason bees) are a large, solitary species in the family Megachilidae, having unique nesting practices, and displaying a wide range in size, colour, and habits. Leafcutter bees build their nests by excising leaf pieces. In contrast, mason bees use resin or clay (Chaudhary and Jain, 1978) to create partitions within tunnels or cavities, resulting in distinct cells for their provisioning and egg-laying leading to progeny

development (Michener, 2007). These complex nesting behaviours show *Megachile* bees' persistence and resourcefulness, supporting pollination dynamics across various plant species. These bees interact with a variety of plant species, especially those in the Leguminosae (Fabaceae) and Compositae (Asteraceae) in Uttarakhand (Kunjwalet *et al.*, 2020), Malvaceae in Uttar Pradesh (Udikeri and Chandra, 2019), Lamiaceae, Umbelliferae, and Leguminosae in Jammu and Kashmir (Shankar and Abrol, 2011), Leguminosae in Tamil Nadu and Odisha (Rakesh *et al.*, 2019; Padhyet *et al.*, 2019), and Cucurbitaceae, in Haryana (Balinaet *et al.*, 2012).

For bees and wasps, which commonly create their nests in pre-existing cavities trap nests are built to provide artificial nesting sites. Trap-nesting wasps are important predators of crop pests while cavity-nesting bees are important pollinators of native plants and crops (Campbell *et al.*, 2017). Trap nests are artificial cavities used by bees and wasps to attract females seeking suitable nesting sites (Staabet *et al.*, 2018). Artificial trap nests are essential in providing a habitat for solitary leafcutter bees. These nests promote bee activity and nesting abundance (Udayakumar and Shivalingaswamy, 2022). Six *Megachile* bee species (subgenus: *Callomegachile*, *Pseudomegachile*, and *Eutricharaea*) have been found to nest in tunnels (5-10 mm diameter) in various nests, including drilled tunnels in posts, wooden logs, and hollowed tunnels in bamboo sticks and reed stems (Kunjwal and Khan, 2023). These artificial nests are used to study cavity nesters' groups and environmental impact on them, with the size of the nest crucial for success, progeny characteristics, and survival. Trap nesting is also a conservation tool in areas where farming practices destructively affect bee populations. Keeping the above-mentioned points in mind, the present study was planned to examine the influence of nest installation on *Megachile* bee abundance, revealing their significant role as potential mung bean pollinators in Uttarakhand, India, and highlighting the need for targeted pollinator management strategies based on environmental factors.

Materials and Methods

The study was conducted to understand the effect of nest installation on the *Megachile* population in the mung bean crop at Norman E. Borlaug Crop Research Centre at the G.B.P.U.A.T., Pantnagar (29.0222° N, 79.4908° E) from Sept. 2021 to Oct. 2022. *Megachile* bees were tallied on flowers from mung beans. Counts were made per square meter area for five minutes amid different day hours using a stopwatch. Overall visits were recorded three times a day. Data were averaged to get a daily count of bees. To study the impact of artificial nesting on *Megachile* abundance, the data on *Megachile* was recorded for two seasons viz., pre- and post-nest installation. Meteorological data from the Department of Agrometeorology, G.B.P.U.A.T., Pantnagar were also used to evaluate the correlation between weather parameters and *Megachile* abundance.

Results and Discussion

Impact of Trap Nests on *Megachile* Bee abundance

Six *Megachile* species [*M. (Creightonella) albironis*, *M. (Xanthosarus) anthracina*, *M. (Amegachile) bicolor*, *M. (Pseudomegachile) lanata*, *M. (Callomegachile) relata*, and *M. (Callomegachile) umbripennis*] were identified to be regular visitors to the experimental fields in both seasons, suggesting that these bees could be possible pollinators for mung

beans. Table 1 and 2 indicate the population abundance (bees/m²/5min) of *Megachile* bees visiting mung bean flowers pre- and post-nest installation. The results presented in Table 1 indicate that the abundance of all *Megachile* pollinators varied significantly during the pre-nest installation phase across the various weeks of the mung bean flowering period. *M. umbripennis* was abundant on most observation weeks, followed by *M. bicolor* and *M. relata*. In the 39th standard meteorological week, *M. umbripennis* had a higher abundance than *M. relata* and other species. *M. albicans* and *M. anthracina* had the lowest activity levels during the observation period. However, Table 2 data from the post-nest installation phase indicates the number of *Megachile* pollinators changed significantly across different standard meteorological weeks throughout the mung bean flowering period. *M. lanata* was the most abundant during the majority of the observation days, followed by *M. umbripennis* and *M. bicolor*. In the 40th standard meteorological week, *M. lanata* had the highest abundance than *M. umbripennis* and other species. Again, *M. albicans* and *M. anthracina* exhibited the lowest activity rates throughout the observation period. Only *M. lanata* nested in the trap nests provided in the fields, and even then, the nest occupancy rate remained low.

These findings point to a broader role for these species in legume pollination. Similar to this study different *Megachile* bees have been reported from all over the country. The importance of *Megachile* bees has been thoroughly studied in states like *M. anthracina* in Jammu and Kashmir (Abrol and Chatterjee, 2022), *M. bicolor* in the pigeon pea in Odisha and Arunachal Pradesh (Padhey et al., 2019; Bora et al., 2021), *M. cephalotes* in mung bean and pigeon pea in Uttarakhand and Odisha, respectively (Kunjwalet et al., 2016; Singh et al., 2017), *M. lanata* in Arunachal Pradesh and Odisha (Bora et al., 2021; Singh et al., 2017), *M. relata* in Odisha and Uttarakhand (Singh et al., 2017; Kunjwalet et al., 2016), and *M. umbripennis* in Uttarakhand (Kunjwalet et al., 2016). This broad spectrum of *Megachile* Bees pollinating legume crops warrants additional exploration into other crops as well.

Correlation with weather parameter

Megachile bees' relative abundance was affected by a variety of meteorological conditions, positive as well as negative. Correlation and regression studies were carried out in 2021 and 2022 to assess the connection between the dependent and independent variables [maximum temperature (T_{MAX}), minimum temperature (T_{MIN}), morning relative humidity (MRH), evening relative humidity (ERH), rainfall (RF), number of rainy days (NRD), sunshine hours (SSH), wind velocity (WV), and evaporation (EVP)]. The impact of different dependent variables on different independent factors on mung bean flowers is described below (Table 3 and Figure 1).

Megachile anthracina

The findings indicated that the pre-nest installation, T_{MAX} , T_{MIN} , MRH, and SSH showed non-significant and positive correlation ($r=0.048, 0.030, 0.185,$ and 0.072), however, ERH, RF, RD, WV, EVP showed non-significant and negative correlation ($r= -0.292, -0.596, -0.530, -0.419$ and -0.434) with *M. anthracina* population. The regression equation being $y = -67.85 + 1.48(T_{MAX}) + 0.12(RHM) + 0.19(RHE) - 0.15(RF)$. Whereas, post-nest installation, *M. anthracina* showed a non-significant and negative correlation with MRH and RF ($r= -0.404$ and -0.585) but a non-significant and positive correlation with T_{MAX} , T_{MIN} , ERH, NRD,

SSH, and WV ($r=0.531, 0.783, 0.378, 0.030, 0.187$ and 0.368). The regression equation being $y=8.48-0.21(\text{RHM})+0.19(\text{RHE})-0.01(\text{RF})+0.15(\text{NRD})$.

Megachilebicolor

The findings indicated that the pre-nest installation, T_{MIN} , ERH, WV, and EVP showed a non-significant and positive correlation ($r=0.680, 0.308, 0.348,$ and 0.213) however, T_{MAX} , MRH, RD, and SSH showed non-significant and negative correlation ($r= -0.459, -0.073, 0.037,$ and -0.469) with *M. bicolor* population. The regression equation being $y=127.47-2.79(T_{\text{MAX}})-0.53(\text{RHM})+0.26(\text{RHE})-0.16(\text{RF})$. Whereas, post-nest installation, *M. bicolor* showed a significant and negative correlation with EVP ($r= -0.919$). The regression equation being $y=24.66-0.41(\text{RHM})+0.22(\text{RHE})-0.37(\text{NRD})$.

Megachilecephalotes

The findings indicated that the pre-nest installation and sunshine hours showed a significant and negative correlation ($r=-0.936$) with *M. cephalotes* population. The regression equation being $y= 7.63 -0.27 (T_{\text{MAX}}) +0.02 (RHE) +0.01 (RF)$. Whereas, post-nest installation, *M. cephalotes* showed non-significant and negative correlation with T_{MAX} , T_{MIN} , MRH, ERH, NRD, SSH, WV, and EVP however non-significant and positive correlation with RF ($r= -0.188, -0.381, -0.127, -0.075, -0.201, -0.305, -0.293, -0.259$ and 0.565). The regression equation being $y=27.30-0.28(\text{RHM})-0.03(\text{RHE})+0.01(\text{RF})-0.11(\text{NRD})$.

Megachilelanata

The findings indicated that the pre-nest installation, T_{MAX} , T_{MIN} , ERH, WV, and EVP showed a non-significant and positive correlation ($r= 0.064, 0.455, 0.022, 0.083,$ and 0.055), however, MRH, RF, RD, and SSH showed non-significant and negative correlation ($r= -0.196, -0.327, -0.384, -0.040$) with *M. lanata* population. The regression equation being $y=-41.17+1.04(T_{\text{MAX}})-0.11(\text{RHM})+0.35(\text{RHE})-0.25(\text{RF})$. Whereas, post-nest installation, *M. lanata* showed a significant and positive correlation with T_{MIN} and a significant and negative correlation with EVP ($r= 0.950$ and -0.880). The regression equation being $y=11.66-0.14(\text{RHM})+0.11(\text{RHE})-0.36(\text{NRD})$.

Megachilerelata

The findings indicated that the pre-nest installation, T_{MIN} , ERH, RF, NRD, WV, and EVP showed a non-significant and positive correlation ($r= 0.849, 0.489, 0.120, 0.083, 0.473,$ and 0.306), however, T_{MAX} , MRH, and SSH showed non-significant and negative correlation ($r= -0.193, -0.159,$ and -0.467) with *M. relata* population. The regression equation being $y=12.78-0.27(T_{\text{MAX}})-0.30(\text{RHM})+0.46(\text{RHE})-0.27(\text{RF})$. Whereas, post-nest installation, *M. relata* showed a significant and negative correlation with EVP ($r= -0.936$). The regression equation being $y=7.32-0.18(\text{RHM})+0.18(\text{RHE})-0.39(\text{NRD})$.

Megachileumbripennis

The findings indicated that the pre-nest installation, T_{MAX} , T_{MIN} , ERH, WV, and EVP showed a non-significant and positive correlation ($r= 0.014, 0.708, 0.327, 0.343,$ and 0.237), however, MRH, RF, NRD, and SSH showed non-significant and negative correlation ($r= -0.220, -0.038, -0.091, -0.256$) with *M. umbripennis* population. The regression equation being

$y = -46.41 + 1.15(T_{MAX}) - 0.19(RHM) + 0.52(RHE) - 0.33(RF)$. Whereas, post-nest installation, *M. umbripennis* showed a significant and negative correlation with T_{MIN} ($r = 0.935$). The regression equation being $y = -29.29 - 0.28(RHM) + 0.17(RHE) - 0.01(RF) - 0.17(NRD)$.

The present study shows similarity with the higher preference of *Megachile* bees towards warmer temperature and higher activity during flowering seasons (Felicioli and Pinzauti, 2008). Temperature and intensity of light impact foraging, with low temperatures and high light intensity leading to foraging and lowering light intensity ceasing daily foraging activities (Sheblet *et al.*, 2008). Megachilid bees thrive in temperatures ranging from 26.9°C to 38.9°C, with peak activity between March-May and October-November, and activity dropping amid high humidity and winter/rainy seasons (Kunjwaletal., 2016). Similar environmental interactions have been identified for *M. lanata* (Abrol and Kapil, 1986). However, the prevalence and nesting of these pollinators in mung bean fields remain unresolved necessitating additional research.

Conclusion

This study highlights the vital role of *Megachile* bees as mung bean pollinators in Uttarakhand, India. Nest installation influences *Megachile* abundance and species responses to environmental conditions, as shown by correlation analysis between bee population and meteorological parameters.

Despite low pollinator population and nest occupancy rates in mung bean, *Megachile* bees were reported to frequently visit mung bean flowers. Trap nests substantially enhance pollinator populations, such as *M. lanata*. This shows the potential of trap nests for increasing bee numbers in mung bean crop as well as in other crops for managed pollination. Understanding the nesting behaviour and habitat management of *Megachile* bees is essential for improving pollination services while promoting sustainable agriculture.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

Abrol, D. P. and D. Chatterjee, (2022). Plant-Pollinator Interaction and Diversity of Megachilid Bees In Different Landscapes Of Jammu, India. *Journal of Palynology Vol, 58*, 15-21. <https://shorturl.at/GJKbh>

Abrol, D. P., & Kapil, R. P. (1986). Factors affecting pollination activity of *Megachile lanata* Lepel. *Proceedings: Animal Sciences*, 95, 757-769. <https://link.springer.com/article/10.1007/BF03179493>

Agricultural Statistics at a Glance, (2022). Directorate of Economics and Statistics, Ministry of Agriculture, Govt. of India (Retrieved from <https://eands.dacnet.nic.in>).

Balina, P., Sharma, S. and Rana, M. 2012. Diversity, abundance and pollination efficiency of native bee pollinators of bitter melon (*Momordica charantia* L.) in India. *J. Apic. Res.*, 51: 227 - 231. <https://ui.adsabs.harvard.edu/abs/2012JApiR..51..227B/abstract>

Bora, M. S., Kanwat, M., Kalita, H., Sasmal, D., Rahman, A., & Ngangom, B. (2022). Diversity of Insects in Pulses at Arunachal Pradesh. *Indian Journal of Entomology*, 409-411.

Campbell, J. W., Smithers, C., Irvin, A., Kimmel, C. B., Stanley-Stahr, C., Daniels, J. C., & Ellis, J. D. (2017). Trap nesting wasps and bees in agriculture: a comparison of sown wildflower and fallow plots in Florida. *Insects*, 8(4), 107. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5746790/>

Chaudhary, J. P. and Jain, J. L. 1978. Nesting and foraging behaviour of a mason bee *Megachile lanata* L (Megachilidae: Hymenoptera). *Indian J. Entomol.* 40(4): 405-411. <https://www.cabidigitallibrary.org/doi/full/10.5555/19810212387>

Elisante, F., Ndakidemi, P., Arnold, S. E., Belmain, S. R., Gurr, G. M., Darbyshire, I., ...& Stevenson, P. C. (2020). Insect pollination is important in a smallholder bean farming system. *PeerJ*, 8, e10102. <https://peerj.com/articles/10102/>

Felicioli, A. and Pinzauti, M. 2008. Pollination by *Osmia* Bees (Hymenoptera: Megachilidae). In JL Capinera (2nd ed.) *Encyclopedia of Entomology*. Springer. 4346p.

Hall, H. G., & Avila, L. (2016). *Megachile sculpturalis*, the giant resin bee, overcomes the blossom structure of sunn hemp (*Crotalaria juncea*) that impedes pollination. *Journal of Melittology*, (65), 1-11. <https://journals.ku.edu/melittology/article/view/5887>

Huppertz, M., Kachhap, D., Dalai, A., Yadav, N., Baby, D., Khan, M. A., ...& Panigrahi, K. C. (2023). Exploring the potential of mung bean: From domestication and traditional selection to modern genetic and genomic technologies in a changing world. *Journal of Agriculture and Food Research*, 100786. <https://www.sciencedirect.com/science/article/pii/S2666154323002934>

Kunjwal, N., & Khan, M. S. (2023). Trap nesting bees of genus *Megachile* and nesting biology of three species of subgenus *Callomegachile* (Hymenoptera; Megachilidae) from North India. *Animal Biology*, 73(2), 171-194. https://brill.com/view/journals/ab/73/2/article-p171_4.xml

Kunjwal, N., Khan, M. S., & Srivastava, P. (2016). Species richness and seasonal activity of the leaf cutter and resin bees (Hymenoptera: Megachilidae) at Pantnagar. *Int. J. Sci. Res*, 5, 972-977. <https://shorturl.at/Owcv0>

Kunjwal, N., Khan, M. S., Kumar, G. and Srivastava, P. 2020. Notes on the nesting ecology of the Megachile bees from North India. *J. Apic. Res.*, 60(5): 807-816. https://www.researchgate.net/publication/342280559_Notes_on_the_nesting_ecology_of_the_Megachile_bees_from_North_India

Michener, C. D. 2007. The Bees of the World. V.66. 2nd ed., Johns Hopkins University press, Baltimore. Maryland. 277-347p. <https://static1.squarespace.com/static/5a849d4c8dd041c9c07a8e4c/t/5ad3bc968a922d44a4728936/1523825933048/Michener+2007+The+Bees+of+the+World.pdf>

Miller, J. C., Barineau, S. M., Smallwood, D. G., Scheuring, D. C., Fernandez, G. J., Park, H. G., ... & Shanmugasundaram, S. (1989). 'TexSprout' Mungbean. *HortScience*, 24(4), 715-716.

Padhy, D., Satapathy, C. R. and Mohapatra, R. N. 2019. Insect fauna associated with pigeon pea (Cajanuscajan L.) in Odisha. *J. pharmacogn. phytochem*, 8(4): 766-768. <https://www.phytojournal.com/archives/2019.v8.i4.9019/insect-fauna-associated-with-pigeon-pea-ltemgtcajanus-cajanltemgt-l-in-odisha>

Pratap, A., Gupta, S., Rathore, M., Basavaraja, T., Singh, C.M., Prajapati, U., Singh, P., Singh, Y., & Kumari, G. (2021). Mungbean. *The Beans and the Peas*. <https://www.sciencedirect.com/science/article/abs/pii/B9780128214503000093>

Rakesh, S. R., Ganesan, N. M., Srinivasan, M. R. and Kumar, M. 2019. Studies on diversity, abundance and pollination efficiency of insect pollinators and pollination mechanism involved in sunnhemp (*Crotalaria juncea* L.). *Plant Breed.*, 10(2): 838-851. https://www.researchgate.net/publication/334202364_Studies_on_diversity_abundance_and_pollination_efficiency_of_insect_pollinators_and_pollination_mechanism_involved_in_sunnhemp_Crotalaria_juncea_L

Shankar, U. and Abrol, D.P. 2011. Pollinators Other Than Honey Bees Visiting Certain Medicinal Plants in Jammu. *J Apic Sci.*, 26: 229-231. https://www.researchgate.net/publication/269168255_Pollinators_other_than_honey_bees_visiting_certain_medicinal_plants_in_Jammu

Shebl, M. A., Kamel, S. M., Hashesh, T. A. A., & Osman, M. A. (2008). Seasonal abundance of leafcutting bees (Megachile minutissima, Megachilidae, Hymenoptera). *World Journal of Agricultural Sciences*, 4(2), 280-287. <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=fabf84132836b4964db2c511934ba782b0ebd5ae>

Singh, A. K. (2017). Abundance and Foraging of Insect Pollinators of Pigeonpea. *Indian Journal of Entomology*, 79(3), 295-299. <https://www.indianjournals.com/ijor.aspx?target=ijor:ije&volume=79&issue=3&article=014&type=pdf>

Staab, M., Pufal, G., Tschardt, T., & Klein, A. M. (2018). Trap nests for bees and wasps to analyse trophic interactions in changing environments—A systematic overview and user guide. *Methods in Ecology and Evolution*, 9(11), 2226-2239. <https://besjournals.onlinelibrary.wiley.com/doi/full/10.1111/2041-210X.13070>

Udikeri, A. and Chandra, U. 2019. Pollination efficiency of different insects on Phalsa *Grewia subinaequalis* DC. *J. Entomol. Zool. Stud.*, 7(1): 1061-1065. <https://www.entomoljournal.com/archives/2019/vol7issue1/PartQ/7-1-202-405.pdf>

Udayakumar, A., & Shivalingaswamy, T. M. (2022). Leafcutter bees as pollinators of pigeon pea: Artificial trap nests as a strategy for their conservation. *Sociobiology*, 69(1), e7202¹ <https://periodicos.uefs.br/index.php/sociobiology/article/view/7202>

UNDER PEER REVIEW

Table 1. The abundance of *Megachile* pollinators on mung bean fields in the blooming period at Pantnagar (2021)

<i>Megachile</i> species	Number of <i>Megachile</i> bees on different dates (bees/m ² /5 min)					Grand Mean
	37 th SMW	38 th SMV	39 th SMV	40 th SMV	41 th SMV	
<i>Megachileanthracina</i>	1.06	1.46	2.52	2.43	1.48	1.90± 0.65
<i>Megachilebicolor</i>	2.49	4.77	4.10	1.75	1.49	3.17± 1.45
<i>Megachilecephalotes</i>	0.66	0.80	0.28	0.54	0.16	0.56± 0.27
<i>Megachilelanata</i>	1.31	1.94	3.59	2.07	0.99	2.11± 1.00
<i>Megachilerelata</i>	2.48	3.56	4.20	2.25	0.48	2.84± 1.42
<i>Megachileumbripennis</i>	3.21	3.81	5.58	3.61	1.63	3.89± 1.41
Mean	2.80±1.00	2.72±1.55	3.38±1.81	2.11±1.00	1.04±0.61	2.22±1.42
S.Em±	0.41	0.63	0.74	0.41	0.25	0.26
CD at 5%	2.401	2.711	3.392	2.685	1.805	1.011


Table 2. The abundance of *Megachile* pollinators on mung bean fields in the blooming period at Pantnagar (2022)

<i>Megachile</i> species	Number of <i>Megachile</i> bees on different dates (bees/m ² /5 min)					Grand Mean
	37 th SMW	38 th SMV	39 th SMV	40 th SMV	41 th SMV	
<i>Megachileanthracina</i>	3.33	1.94	1.52	1.01	0.42	1.64± 1.10
<i>Megachilebicolor</i>	2.97	4.55	2.97	2.16	1.63	2.86± 1.10
<i>Megachilecephalotes</i>	0.24	0.97	1.35	0.83	0.94	0.86± 0.40
<i>Megachilelanata</i>	5.31	5.98	5.09	5.38	3.88	5.13± 0.77
<i>Megachilerelata</i>	2.40	4.28	2.50	2.12	1.79	2.62± 0.97
<i>Megachileumbripennis</i>	5.92	5.56	3.96	4.89	3.45	4.76± 1.05
Mean	3.36±2.06	3.87±2.00	2.89±1.44	2.73±1.95	2.01±1.38	2.98±1.77
S.Em±	0.84	0.82	0.59	0.80	0.56	0.32
CD at 5%	3.176	2.495	2.805	3.125	1.742	0.893

Table 3. Correlation between *Megachile* pollinators and weather parameters on mung bean flowers (2021-2022)

Megachile Species	T _{MAX}		T _{MIN}		MRH		ERH		RF		NRD		SSH		WV		EVP	
	2021	2022	2021	2022	2021	2022	2021	2022	2021	2022	2021	2022	2021	2022	2021	2022	2021	2022
<i>M. albifrons</i>	-0.574	-0.188	0.771	-0.381	0.361	-0.127	0.843	-0.075	0.661	0.565	0.799	-0.201	-	-0.305	0.496	-	0.156	-0.259
<i>M. anthracina</i>	0.048	0.531	0.030	0.783	0.185	-0.404	-0.292	0.378	-0.596	-0.585	-0.530	0.030	0.072	0.187	-0.419	0.368	-0.434	-0.276
<i>M. bicolor</i>	-0.459	0.382	0.680	0.809	-0.073	-0.173	0.308	0.742	0.000	0.069	-0.037	-0.378	-0.469	-0.413	0.348	-	0.213	-0.919*
<i>M. lanata</i>	0.064	0.803	0.455	0.950*	-0.196	-0.456	0.022	0.285	-0.327	-0.409	-0.384	-0.752	-0.040	0.120	0.083	-	0.055	-0.880*
<i>M. relata</i>	-0.193	0.244	0.849	0.709	-0.159	0.028	0.489	0.770	0.120	0.245	0.083	-0.414	-0.467	-0.546	0.473	-	0.306	-0.936*
<i>M. umbripennis</i>	0.014	0.649	0.708	0.935*	-0.220	-0.213	0.327	0.352	-0.038	-0.569	-0.091	-0.382	-0.256	0.177	0.343	-	0.237	-0.521

Figure 1: Correlation between *Megachile* pollinators and weather parameters on mung bean flowers (2021-2022)



UNDER PEER REVIEW