

# Effect of Plant Growth Regulators on Growth and Flower Yield of *Lisianthus* (*Eustoma grandiflorum*) Under Prayagraj Agro-Climatic Conditions

## ABSTRACT

The experiment was carried out under naturally ventilated polyhouse conditions, Department of Horticulture, SHUAT S, Prayagraj, during November, 2023 to April, 2024. The experiment was laid out in Completely Randomized Design (CRD) with three replications. There were ten treatments comprising of CCC, GA<sub>3</sub>, NAA at different concentrations viz., GA<sub>3</sub> (100, 200, 300) ppm and CCC (250, 500, 750) ppm and NAA (30, 45, 60) ppm along with control. The results revealed that foliar application of 300 ppm GA<sub>3</sub> for *Lisianthus*, significantly enhanced the treatment T<sub>3</sub> - GA<sub>3</sub> at 300 ppm was found in superior among other treatments in terms of plant height (20.66cm), number of leaves (33.06), plant spread (11.3cm), stem diameter (3.23cm), leaf area (15.43cm<sup>2</sup>), bud length (2.62cm), number of buds per plant (6.43), stalk length (5.70cm), Number of flowers per plant (6.43), number of branches per plant (3.12), chlorophyll content (66.10), minimum number of days for bud initiation (122.50), Flower yield per plant (6.43), flower yield per 250m<sup>2</sup> (40187.50), benefit cost ratio (4.08) per 250 m<sup>2</sup> was also observed in T<sub>3</sub> - GA<sub>3</sub> at 300 ppm.

*Keywords: Lisianthus, NAA, GA<sub>3</sub>, CCC, Polyhouse, PGR*

## 1. INTRODUCTION

*Eustoma grandiflorum* Shinn. Syn. *Lisianthus russelianus* Hook., (2n=36) belongs to family Gentianaceae. *Lisianthus* is also known as 'Texas Blue Bell' and Prairie Gentian (Halevy and Kofranek, [7]). *Eustoma* is named after the two Greek words Eu (beautiful, good, well), and stoma (mouth). Although delicate and soft-hued in style, *Lisianthus* flowers boast powerful symbolism. This floral species represents charisma, appreciation, and gratitude. It mainly used for beautification, bouquets, showy purposes and decorating walls, fences, gates, hedges. It is not only popular as a cut flower but also a bedding plant and a pot plant in the market (Grueber [5] and Harbaugh [8]). *Lisianthus* (*Eustoma grandiflorum* [Raf.] Shinn) is an ornamental species that has acquired great importance due to the attractiveness of its flowers, bright colours and long vase life. As cut flower, its acceptance in the international markets has increased in recent years and currently it is among the 10 most sold flowers in the world. It is

a plant native to the northern states of Mexico and southern United States of America (Domínguez-Ramírez [3]). Lisianthus is an annual or biannual species; the period from sowing to flowering can be divided into two stages. The first one is from seed germination up to the liners are ready for transplant; in this stage, the plants form a rosette with four leaf pairs. The second stage begins with the elongation of the stems and ends with flowering. Lisianthus has been ranked one of the lucrative and expensive cut flowers in the world. It has a great cosmopolitan demand mainly for its large and attractive flowers, long and hard stem, wide range of colors specially have purple, blue, lavender and has a long vase life which are the common traits of an ideal and best quality cut flower. Lisianthus, herbaceous annual growing to 15 to 60 cm tall, with bluish green, slightly succulent leaves. These are having large funnel-shaped flowers growing 32cm long straight stems, sometimes erect single stems, other times growing on branching stems that can rise to be eighteen feet tall. The flowers can grow up to two inches across and can be found in a variety of colors. Lisianthus are long-stemmed flowers in cymes, with often only a few openings at a time. Sepals on Lisianthus are only fused close to the base and are much smaller than petals Namratha et al. [15]. In Tamil Nadu, production of lisianthus is concentrated in limited area in the hill stations namely, Ooty and Kodaikanal. Lisianthus is a crop which has immense potential as cut flower and pot plant (katharai et al. [10]). Research work on the use of Gibberellic acid, CCC, NAA plays a vital role in improving the vegetative growth characters of the plants as they enhance the elongation and cell division by promoting the DNA synthesis in the cell. It reduced the juvenile phase due to increase in photosynthesis and respiration with enhanced CO<sub>2</sub> fixation in the plant Thirumal et al. [20] Gibberellic acid (GA<sub>3</sub>) is endogenous plant hormone, belongs to gibberellins group and takes part in various metabolic processes (Eason [4]) GA<sub>3</sub> is mostly known to enhance the production of hydrolyses ( $\alpha$ -amylase) in the endosperm of cereals (Kolumbina.et.al., [11])



Fig. Morphology of *Eustoma Grandiflorum*

## 2. MATERIAL AND METHODS

The present investigation was carried out under naturally ventilated polyhouse department of horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, in November 2023- April 2024. The Lisianthus seedling F<sub>1</sub> Rosie variety was planted on polythene bags. The bags were filled with a mixture of soil, vermicompost, Farm yard manure and cocopeat in the ratio of 1:1:1:1. Before planting the seedling in the polythene bags, soil was treated with fungicide, i.e. Copper oxychloride at 2%. During transplanting potting media was pressed firmly around the seedlings so that they do not get disturbed after irrigation. Also in this experiment, plant was sprayed with Gibberellic acid (GA<sub>3</sub>) (at 100, 200,

300 ppm), cycocel (CCC) (at 250, 500, 750 ppm) and Naphthalene Acetic Acid (NAA)( at 30, 60, 90 ppm). Foliar spraying was repeated for several periods 30, 60, 90, 120, 150 days of intervals, three plants were selected randomly from each treatment per replication and their observation were recorded at 30 days intervals. Statistical analysis was conducted using Fisher's analysis of variance (ANOVA) technique, in accordance with the guidelines.

**Table 1. Treatment details**

Sl.No.	Treatment	Notation
1	T <sub>0</sub>	Control
2	T <sub>1</sub>	GA3 at 100 ppm
3	T <sub>2</sub>	GA3 at 200 ppm
4	T <sub>3</sub>	GA3 at 300 ppm
5	T <sub>4</sub>	Cycocel at 250ppm
6	T <sub>5</sub>	Cycocel at 500ppm
7	T <sub>6</sub>	Cycocel at 750ppm
8	T <sub>7</sub>	NAA at 30 ppm
9	T <sub>8</sub>	NAA at 45 ppm
10	T <sub>9</sub>	NAA at 60 ppm

### 3. RESULT AND DISSCUSSION

Effect of plant growth regulators on different concentration of vegetative, flowering, yield and economic parameters are presented in table 2, 3, 4 and 5 respectively. The Data was taken from DEC-APR.

#### 3.1 Vegetative parameter

##### 3.1.1 Plant height (cm)

Maximum Plant height (cm) was observed with T<sub>3</sub> - GA<sub>3</sub> at 300ppm (20.66 cm) which was at par with T<sub>7</sub>- CCC at 750 ppm (18.19 cm) whereas, minimum was reported in T<sub>0</sub> - Control (13.86 cm). These are similar finding reported by Kathari [10] in lisianthus. and Aparna [21] reported GA<sub>3</sub> induce m-RNA synthesis pertaining to hydrolytic enzymes, which promotes mitotic activity in apical meristem and increase cell division and cell elongation, leading to increased length of internodes, in turn increases the plant height. Providing conducive environment inside the naturally ventilated polyhouse at night time helped in better plant height also.

### **3.1.2 Number of leaves per plant**

Maximum number of leaves per plant was observed with T<sub>3</sub> - GA<sub>3</sub>at 300ppm (33.06) which was at par with T<sub>9</sub> - NAA at 60 ppm (29.40) whereas, minimum was reported in T<sub>4</sub>- CCC at 250ppm (25.0). Similar trend was also observed previously in chrysanthemum Habiba et al. [6] and he reported GA<sub>3</sub> influencing the increased vegetative growth by increasing cell division and cell elongation is the reason behind the production of a greater number of leaves per plant

### **3.1.3 Plant Spread (cm<sup>2</sup>)**

Maximum Plant Spread (cm<sup>2</sup>) was observed with T<sub>3</sub> - GA<sub>3</sub> at 300 ppm (11.3 cm<sup>2</sup>) which was at par with T<sub>5</sub> - CCC at 500 ppm (9.54 cm<sup>2</sup>) whereas, minimum was reported in T<sub>0</sub> - Control (7.96 cm<sup>2</sup>). This might be attributed to the higher concentration of GA<sub>3</sub>, which is more efficient in cell multiplication and elongation of immature tissues, whereas lesser quantities were less favourable. Palei [16]and Doddagoudar [2] comparable results reported that the application of GA<sub>3</sub> plant growth regulators enhances cell division with rapid internode elongation and is confined in the sub-apical meristem, which increases the plant spread.

### **3.1.4 Number of shoots per plant**

Maximum number of shoots per plant was observed with T<sub>3</sub> - GA<sub>3</sub> at 300 ppm (3.12) which was at par with T<sub>2</sub> - GA<sub>3</sub> at 200 ppm (2.35) whereas, minimum was reported in T<sub>9</sub>- NAA at 60ppm (1.87). The increase in number of branches per plant application of GA<sub>3</sub> may be ascribed to increase cell division and cell enlargement, promotion of protein synthesis in the plant. Stimulation of branching may also be attributed to the breakage of apical dominance. (Benny *et.al.*) [1].

### **3.1.5 Chlorophyll content**

The maximum Chlorophyll content was observed with T<sub>3</sub> - GA<sub>3</sub> at 300 ppm (66.10  $\mu\text{mol m}^{-2}$ ) which was at par with T<sub>7</sub>- NAA at 30 ppm (64.61  $\mu\text{mol m}^{-2}$ ) whereas, minimum was reported in T<sub>1</sub>- GA<sub>3</sub> at 100ppm (60.70  $\mu\text{mol m}^{-2}$ ). Changes in chlorophyll content caused by growth regulators may be attributed to decrease chlorophyll breakdown and enhance the chlorophyll synthesis.

### **3.1.6 Stem diameter (cm)**

The maximum Stem diameter (cm) was observed with T<sub>3</sub> - GA<sub>3</sub> at 300ppm (3.23 cm) which was at par with T<sub>8</sub>- NAA at 45 ppm (2.53 cm) whereas, minimum was reported in T<sub>0</sub> - Control (1.85 cm). Increase in stem diameter is directly proportional to increase in plant height. The maximum plant height was obtained in T<sub>3</sub> which may have been led to maximum diameter of stem T<sub>3</sub>. With the increase in the concentration of GA<sub>3</sub> diameter increases due to a reflection of the stimulation of cambium and its immediate cell progeny. Similar Doddagoudar [2], reported comparable results in marigold.

### **3.1.7 Maximum Leaf area (cm<sup>2</sup>)**

The data on vegetative parameters are given in table 2. It is clear from the table that is significant difference among the treatments for number of leaves per plant of lisianthus. Kumar [12] reported that GA<sub>3</sub> promotes cell division and cell elongation, thereby increase the leaf length and leaf area and enhance sugar translocation.

## **3.2 Floral parameters**

### **3.2.1 Stalk length(cm)**

The maximum Stalk length (cm) was observed with T<sub>3</sub> - GA<sub>3</sub> at 300ppm (5.70 cm<sup>2</sup>) which was at par with T<sub>4</sub>- Cycocel at 250ppm (5.14 cm<sup>2</sup>) whereas, minimum was reported in T<sub>0</sub> - Control (3.92 cm<sup>2</sup>). GA<sub>3</sub> was the best for obtaining better growth of plants, maximum number of cut blooms with longer stalk as well as bigger flower size obtained. A similar trend was observed previously in China aster by Sailaja and Panchali [19].

### **3.2.2 Bud length (cm)**

The maximum Bud length (cm) was observed with T<sub>3</sub> - GA<sub>3</sub> at 300ppm (2.62 cm) which was at par with T<sub>2</sub>- GA<sub>3</sub> at 200 ppm and T<sub>1</sub> GA<sub>3</sub> at 100 ppm (2.45 cm) whereas, minimum was reported in T<sub>7</sub> - NAA at 30 ppm (2.08 cm). Justo et al.in carnation reported that the increase in the length of the flower bud in GA<sub>3</sub> treated plants is due to rapid cell elongation, increased cell

divisions and cell enlargement. Foliar application of GA<sub>3</sub> also significantly increased flower bud length.

### **3.2.3 Number of buds per plant**

The data on floral parameters are given in table 3. It is clear from the table that significant difference among the treatments for number of buds per plant of lisianthus. The maximum number of buds per plant was observed with T<sub>3</sub>- GA<sub>3</sub> at 300 ppm (6.43) which was at par with T<sub>8</sub> - NAA at 45 ppm (5.34) whereas, minimum was reported in T<sub>4</sub>- CCC at 250ppm (4.02). The effect of different growth regulators showed significant influence on the number of buds per plant. Similar results were recorded by Kumar et al. [13] in marigold.

### **3.2.4 Number of days for bud initiation**

The minimum Number of days for bud initiation was observed with T<sub>3</sub> - GA<sub>3</sub> at 300 ppm (122.50) which was at par with T<sub>5</sub> - CCC at 500 ppm (128.7) whereas, maximum was reported in T<sub>7</sub>- NAA at 30ppm (131.06). Holkar [9] reported that GA<sub>3</sub> application enhances food translocation for the development of floral primordia, which leads to early flowering. This is due to increase in photosynthesis and respiration along with enhanced fixation by GA<sub>3</sub> that led to flower bud initiation.

### **3.2.5 Number of flowers per plant**

The maximum number of flowers per plant was observed with T<sub>3</sub> - GA<sub>3</sub> at 300 ppm (6.43) which was at par with T<sub>8</sub> - NAA at 45 ppm (5.34) whereas, minimum was reported in T<sub>4</sub>- CCC at 250ppm (4.02). Increased number of flower is attributed to the production of large number of flower buds along with the fact that termination of vertical growth by pinching lead to more laterals/secondary branches at early stage of growth, which then had sufficient time to accumulate carbohydrates for proper flower bud differentiation producing a greater number of flowers per plant. GA<sub>3</sub> helps in breaking bud dormancy and acts as florigen initiating flowering. Similar results were also reported by Mithilesh Kumar [14] in Marigold.

### **3.3 Yield Parameters**

#### **3.3.1 Flower yield per plant**

The maximum flower yield per plant was observed with T<sub>3</sub> - GA<sub>3</sub> at 300 ppm (6.43) which was at par with T<sub>8</sub> - NAA at 45 ppm (5.34) whereas, minimum was reported in T<sub>4</sub>- CCC at 250ppm (4.02). Increase in flower yield per plant due to the fact that plant growth regulators stimulate vegetative growth and induced changes in vegetative morphology, and thereby accelerate growth parameters. These findings are also in accordance with those of Rakesh [17] in carnation.

#### **3.3.2 Flower yield per 250 m<sup>2</sup>**

The maximum flower yield per 250 m<sup>2</sup> was observed with T<sub>3</sub> - GA<sub>3</sub> at 300 ppm (40187.50) which was at par with T<sub>8</sub> - NAA at 45 ppm (33375.00) whereas, minimum was reported in T<sub>4</sub> - CCC at 250ppm (25125.00). GA<sub>3</sub> treatment enhance induction of flower bud break i.e., differentiation of floral primordia in the apical region which leads to increase production of flowers per plants and hence increase the flower yield. These results are also in close conformity with the findings of Ryagi [18] in Carnation.

#### **3.4 Economic Parameter**

The maximum gross return was observed with T<sub>3</sub> - GA<sub>3</sub> at 300 ppm (401875.0) which was at par with T<sub>8</sub> - NAA at 45 ppm (3.50) whereas, minimum was reported in T<sub>4</sub>- CCC at 250 ppm (251250.0). The maximum net return was observed with T<sub>3</sub> - GA<sub>3</sub> at 300 ppm (303412.00) which was at par with T<sub>8</sub> - NAA at 45 ppm (3.50) whereas, minimum was reported in T<sub>4</sub>- CCC at 250 ppm (148387.00). The maximum Benefit cost ratio was observed with T<sub>3</sub> - GA<sub>3</sub> at 300 ppm (4.08) which was at par with T<sub>8</sub> - NAA at 45 ppm (3.50) whereas, minimum was reported in T<sub>4</sub>- CCC at 250 ppm (2.44).

UNDER PEER REVIEW

**Table 2. Effect of different concentration of GA3, CCC and NAA on vegetative parameters of Lisianthus.**

<b>Treatment Notation</b>	<b>Treatments</b>	<b>Plant spread (cm<sup>2</sup>)</b>	<b>Plant height (cm)</b>	<b>Number of Leaves</b>	<b>Number of Shoots</b>	<b>Chlorophyll content (μmol m<sup>-2</sup>)</b>	<b>Stem Diameter (cm)</b>	<b>Leaf area (cm<sup>2</sup>)</b>
<b>T<sub>0</sub></b>	Control	7.96	13.86	26.29	2.18	61.43	1.85	13.34
<b>T<sub>1</sub></b>	GA3at 100 ppm	8.06	15.30	25.04	2.30	60.70	2.27	11.29
<b>T<sub>2</sub></b>	GA3at 200 ppm	8.09	16.28	26.26	2.35	60.86	2.14	13.35
<b>T<sub>3</sub></b>	GA3at 300 ppm	11.3	20.66	33.06	3.12	66.10	3.23	15.43
<b>T<sub>4</sub></b>	Cycocelat 250ppm	8.50	15.96	25.0	2.08	62.92	2.37	12.97
<b>T<sub>5</sub></b>	Cycocelat 500ppm	9.54	17.61	26.33	2.11	63.12	2.19	13.09
<b>T<sub>6</sub></b>	Cycocelat 750ppm	9.38	16.81	25.20	1.90	63.54	2.51	12.20
<b>T<sub>7</sub></b>	NAAat 30 ppm	9.5	18.19	27.53	1.94	64.61	2.47	13.38
<b>T<sub>8</sub></b>	NAAat 45 ppm	8.99	16.98	28.04	1.92	63.68	2.53	13.60
<b>T<sub>9</sub></b>	NAAat 60 ppm	9.1	16.21	29.40	1.87	63.65	2.42	13.93
	F-Test	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
	S.Ed (+)	0.14	0.90	0.30	0.16	0.53	0.25	0.92
	CD (5%)	0.30	0.89	0.64	0.33	1.11	0.54	1.93
	CV	1.95	6.57	1.38	8.97	1.03	13.21	8.52

**Table 3. Effect of different concentration of GA3, CCC and NAA on floral parameters of Lisianthus.**

Treatment Notation	Treatments	Stalk length(cm)	Bud length(cm)	No. of buds per plants	No. of days for bud initiation	No. of flowers per plant
T <sub>0</sub>	Control	3.92	2.13	4.49	4.49	4.49
T <sub>1</sub>	GA3at 100 ppm	4.36	2.43	4.99	4.99	4.99
T <sub>2</sub>	GA3at 200 ppm	3.77	2.45	5.31	5.31	5.31
T <sub>3</sub>	GA3at 300 ppm	5.70	2.62	6.43	6.43	6.43
T <sub>4</sub>	Cycocelat 250ppm	5.14	2.09	4.02	4.02	4.02
T <sub>5</sub>	Cycocelat 500ppm	4.69	2.30	4.45	4.45	4.45
T <sub>6</sub>	Cycocelat 750ppm	4.68	2.11	5.28	5.28	5.28
T <sub>7</sub>	NAAat 30 ppm	4.75	2.08	5.13	5.13	5.13
T <sub>8</sub>	NAAat 45 ppm	4.88	2.14	5.34	5.34	5.34
T <sub>9</sub>	NAAat 60 ppm	4.93	2.11	5.19	5.19	5.19
	F-Test	S	S	S	S	S
	S.Ed (+)	0.12	0.09	0.25	0.25	0.25
	CD (5%)	0.25	0.18	0.54	0.54	0.54
	CV	3.22	4.87	6.25	6.25	6.25

**Table 4. Effect of different concentration of GA3, CCC and NAA on yield parameters of Lisianthus.**

Treatment Notation	Treatments	Yield per plant	Yield per (250 m <sup>2</sup> )
T <sub>0</sub>	Control	4.49	28062.50
T <sub>1</sub>	GA3at 100 ppm	4.99	31187.50
T <sub>2</sub>	GA3at 200 ppm	5.31	33187.50
T <sub>3</sub>	GA3at 300 ppm	6.43	40187.50
T <sub>4</sub>	Cycocelat 250ppm	4.02	25125.00
T <sub>5</sub>	Cycocelat 500ppm	4.45	27812.50

<b>T<sub>6</sub></b>	Cycocelat 750ppm	5.28	33000.00
<b>T<sub>7</sub></b>	NAAat 30 ppm	5.13	32062.50
<b>T<sub>8</sub></b>	NAAat 45 ppm	5.34	33375.00
<b>T<sub>9</sub></b>	NAAat 60 ppm	5.19	32437.50
	F-Test	S	S
	S.Ed (+)	0.25	480.11
	CD (5%)	0.54	1017.78
	CV	6.25	13.05

**Table 5. Effect of different concentration of GA<sub>3</sub>, CCC and NAA on Economic parameters of Lisianthus.**

<b>Treatment Notation</b>	<b>Treatments combination</b>	<b>Total cost of cultivation</b>	<b>Gross Return</b>	<b>Net Return</b>	<b>Benefit cost ratio</b>
<b>T<sub>0</sub></b>	Control	94875.00	280625.0	185750.00	2.95
<b>T<sub>1</sub></b>	GA <sub>3</sub> at 100 ppm	96063.00	311875.0	215812.00	3.24
<b>T<sub>2</sub></b>	GA <sub>3</sub> at 200 ppm	97263.00	331875.0	214612.00	3.20
<b>T<sub>3</sub></b>	GA <sub>3</sub> at 300 ppm	98463.00	401875.0	303412.00	4.08
<b>T<sub>4</sub></b>	Cycocel at 250ppm	102863.00	251250.0	148387.00	2.44
<b>T<sub>5</sub></b>	Cycocel at 500ppm	107752.00	278125.0	170373.00	2.58
<b>T<sub>6</sub></b>	Cycocel at 750ppm	113461.00	330000.0	216539.00	2.90
<b>T<sub>7</sub></b>	NAA at 30 ppm	95231.00	320625.0	225394.00	3.36
<b>T<sub>8</sub></b>	NAA at 45 ppm	95355.00	333750.0	238395.00	3.50
<b>T<sub>9</sub></b>	NAA at 60 ppm	95434.00	324375.0	228941.00	3.39

## CONCLUSION

From the present investigation, it is concluded that among the different treatments, the treatment T<sub>3</sub> - GA<sub>3</sub> at 300 ppm was found in superior among other treatments in terms of Plant height, number of leaves , plant spread , stem diameter, leaf area, bud length, number of buds per plant, stalk length , Number of flowers per plant, number of branches per plant, chlorophyll content, minimum number of days for bud initiation, Flower yield per plant, flower yield per 250m<sup>2</sup>, benefit cost ratio per 250m<sup>2</sup> was also observed in T<sub>3</sub> - GA<sub>3</sub> at 300 ppm. GA<sub>3</sub> at 300 ppm could be recommended for improved plant growth and flowering of Lisianthus (*Eustoma grandiflorum*).

## REFERENCES

1. **Benny JC, Devi Singh DS, Fatmi U, Jose DA. (2018).** Effect of plant growth regulators, Gibberellic acid (GA<sub>3</sub>) and Naphthalene Acetic Acid (NAA) on growth and yield of carnation (*Dianthus caryophyllus* L.) under naturally ventilated polyhouse. *Plant Archives*, 17. 803-812.
2. **Doddagoudar SR, Vyakarnahal BS, Gouda SM. (2004).** Effect of mother plant nutrition and chemical spray on seed germination and seedling vigour of China aster cv. Kamini Karnataka, *Journal of agriculture sciences*, 17(4):701-704.
3. **Domínguez-Ramírez, A. (2002).** El Cultivo del Lisianthus (*Eustoma grandiflorum*). [www.uaaan.mx/postgrado/images/files/hort/simposio2/Ponencia07.pdf](http://www.uaaan.mx/postgrado/images/files/hort/simposio2/Ponencia07.pdf).
4. **Eason, J.R. (2002).** Sandersonia aurantiaca: an evaluation of postharvest pulsing solutions to maximize cut flower quality. *New Zealand J. Crop Hort. Sci.*, 30: 273-279.
5. **Grueber KL, Corr BE, Wilkins HF. (1985).** Evaluation of *Eustoma grandiflorum* as a bedding plant. *Minn. Stat e Flor. Bull*, 34(1):16-18.
6. **Habiba SU, Islam MS, Jamal Uddin AFM. (2012).** Influence of terminal bud pinching on growth and yield of chrysanthemum, (*Chrysanthemum indicum* L). *Journal of Bangladesh Academy of Sciences*, 36(2):251-255.
7. **Halevy, A.H., and Kofranek, A.M. (1984).** Evaluation of lisianthus as a new flower crop. *HortScience* 19, 845–847.
8. **Harbaugh BK. (2007).** Lisianthus. *Eustoma grandiflorum*. In: Anderson, N.O. (Ed.), *Flower breeding and genetics. Springer, Netherlands*, 645-663.
9. **Holkar PS, Chandrashekar SY, Hemanthkumar P, Basavalingaiah, Ganapathi M. (2018).** Effect of gibberellic acid on growth, flowering, flower quality and corm yield of gladiolus, *Trends in Biosciences*, 11(24):32113216.
10. **Kat hari Lakshmaiah, S Subramanian, M Ganga and P Jeyakumar (2019).** Optimization of pinching and GA<sub>3</sub> application to improve growth and flowering of lisianthus (*Eustoma grandiflorum*). *Journal of Pharmacognosy and Phytochemistry* 2019; 8(6): 614-616.
11. **Kolumbina, M., M. Blesing and D.J. Mares. (2006).**  $\alpha$ -Amylase and programmed cell death in aleurone of ripening wheat grains. *J. Exp. Bot.*, 57(4): 877-885.
12. **Kumar P, Singh A, Laishram N, Pandey RK, Dogra S, Jeelani MI, Sinha BK. (2019)** Effects of plant growth regulators on quality flower and seed production of marigold (*Tagetes erecta* L.)
13. **Kumar Suresh P, Bagawathi R, Rajive Kumar, Ronya T.(2008).** Effect of plant growth regulators on vegetative growth, flowering and corm production of gladiolous in Arunachal Pradesh. *J of Orn. Horti*, 10(2):90-95.
14. **Kumar, Mithilesh & Singh, AK & Kumar, Ashok. (2015).** Effect of plant growth regulators on growth and flowering characters of African marigold (*Tagetes erecta* ). *Current Advances in Agricultural Sciences*. 7. 85. 10.5958/2394-4471.2015.00020.9.
15. **Namrat ha G, Chandrashekar SY, Hemla Naik B, Shivaprasad M and Hanumantharaya L (2021).** Varietal evaluation of Lisianthus (*Eustoma grandiflorum* Shinn.) for morphological parameters under protected cultivation. *The Pharma Innovation Journal*, 10(12): 2160-2162.

16. **Palei S, Das AK, Dash DK. (2016).** Effect of plant growth regulators on growth, flowering and yield at tributaries of African marigold (*Tagetes erecta L.*), *Research Paper Agriculture*. 2(1):2454-9916.
17. **Rakesh., Singhrot, R. S. and Beniwal, B. S. (2003).** Effect of GA3 and pinching on growth and yield in Chrysanthemum. *Haryana J. hort. Sci.* 32(1-2): 61-63.
18. **Ryagi VY, Mantur SM, Reddy B S.(2007).** Effect of pinching on growth, yield and quality of flowers of Carnation varieties grown under polyhouse. *Karnataka J Agric. Sci*, 20(4):816-818.
19. **Sailaja SM, Panchbhai DM.(2014).** Effect of pinching on growth and quality characters of China aster varieties. *Asian J Hort*, 9(1):36-39.
20. **Thirumal Murugan V, K Manivannan and S Nanthakumar (2021).** Studies on the effect of plant growth regulators on growth, flowering and xanthophyll content of African marigold (*tagetes erecta L.*). *J Pharmacogn Phytochem*, 10(3):292-294.
21. **Veluru Aparna, Krishna Prakash, Neema M, Arora Ajay, Naveen Kumar P, Singh MC. (2018).** Effect of gibberellic acid on plant growth and flowering of Chrysanthemum cv. Thai Chen Queen under short day planting conditions. *International Journal of Agriculture Sciences*. ISSN: 0975-3710 & E-ISSN: 0975-9107. 10(11):6274- 6278.